

# Statistics for Data Science -1

## Lecture 5.3: Permutation formula-distinct objects

Usha Mohan

Indian Institute of Technology Madras

## Learning objectives

1. Understand basic principles of counting.
2. Concept of factorials.
3. Understand differences between counting with order (permutation) and counting without regard to order (combination).
4. Use permutations and combinations to answer real life applications.

## Permutations

Permutation when objects are distinct

Permutation when objects are distinct- repetitions not allowed

Permutation when objects are distinct- repetitions allowed

# Permutation

## Definition

A *permutation* is an ordered arrangement of all or some of  $n$  objects.

## Example

## Example

Take  $A, B, C$ - Possible arrangements- taking all at a time

## Example

Take  $A, B, C$ - Possible arrangements- taking all at a time

First place	Second place	Third place
A	B	C
A	C	B
B	A	C
B	C	A
C	A	B
C	B	A

## Example

Take  $A, B, C$ - Possible arrangements- taking two at a time



## Example

Take  $A, B, C$ - Possible arrangements- taking two at a time

First place	Second place
A	B
A	C
B	A
B	C
C	A
C	B

## Example

Take  $A, B, C, D$ - Possible arrangements- taking all at a time

# Example

Take  $A, B, C, D$ - Possible arrangements- taking all at a time

First place	Second place	Third place	Fourth place
A	B	C	D
A	B	D	C
A	C	B	D
A	C	D	B
A	D	B	C
A	D	C	B
B	A	C	D
B	A	D	C
B	C	A	D
B	C	D	A
B	D	A	C
B	D	C	A
C	A	B	D
C	A	D	B
C	B	A	D
C	B	D	A
C	D	A	B
C	D	B	A
D	A	B	C
D	A	C	B
D	B	A	C
D	B	C	A
D	C	A	B
D	C	B	A

## Example

Take  $A, B, C, D$ - Possible arrangements- taking two at a time

## Example

Take  $A, B, C, D$ - Possible arrangements- taking two at a time

First place	Second place
A	B
A	C
A	D
B	A
B	C
B	D
C	A
C	B
C	D
D	A
D	B
D	C

## Permutation formula

The number of possible permutations of  $r$  objects

## Permutation formula

The number of possible permutations of  $r$  objects from a collection of  $n$  **distinct** objects is given by the formula

## Permutation formula

The number of possible permutations of  $r$  objects from a collection of  $n$  **distinct** objects is given by the formula

$$n \times (n - 1) \times \dots \times (n - r + 1)$$

and is denoted by  ${}^n P_r$



## Permutation formula

The number of possible permutations of  $r$  objects from a collection of  $n$  **distinct** objects is given by the formula

$$n \times (n - 1) \times \dots \times (n - r + 1)$$

and is denoted by  ${}^n P_r$

$${}^n P_r =$$

## Permutation formula

The number of possible permutations of  $r$  objects from a collection of  $n$  **distinct** objects is given by the formula

$$n \times (n - 1) \times \dots \times (n - r + 1)$$

and is denoted by  ${}^n P_r$

$${}^n P_r = \frac{n!}{(n - r)!}$$

## Permutation formula

The number of possible permutations of  $r$  objects from a collection of  $n$  **distinct** objects is given by the formula

$$n \times (n - 1) \times \dots \times (n - r + 1)$$

and is denoted by  ${}^n P_r$

$${}^n P_r = \frac{n!}{(n - r)!}$$

► Special cases

1.  ${}^n P_0 =$

## Permutation formula

The number of possible permutations of  $r$  objects from a collection of  $n$  **distinct** objects is given by the formula

$$n \times (n - 1) \times \dots \times (n - r + 1)$$

and is denoted by  ${}^n P_r$

$${}^n P_r = \frac{n!}{(n - r)!}$$

### ► Special cases

1.  ${}^n P_0 = \frac{n!}{(n-0)!} = \frac{n!}{n!} = 1$  There is only one ordered arrangement of 0 objects.

## Permutation formula

The number of possible permutations of  $r$  objects from a collection of  $n$  **distinct** objects is given by the formula

$$n \times (n - 1) \times \dots \times (n - r + 1)$$

and is denoted by  ${}^n P_r$

$${}^n P_r = \frac{n!}{(n - r)!}$$

### ► Special cases

1.  ${}^n P_0 = \frac{n!}{(n-0)!} = \frac{n!}{n!} = 1$  There is only one ordered arrangement of 0 objects.
2.  ${}^n P_1 =$

## Permutation formula

The number of possible permutations of  $r$  objects from a collection of  $n$  **distinct** objects is given by the formula

$$n \times (n - 1) \times \dots \times (n - r + 1)$$

and is denoted by  ${}^n P_r$

$${}^n P_r = \frac{n!}{(n - r)!}$$

### ► Special cases

1.  ${}^n P_0 = \frac{n!}{(n-0)!} = \frac{n!}{n!} = 1$  There is only one ordered arrangement of 0 objects.
2.  ${}^n P_1 = \frac{n!}{(n-1)!} = n$ . There are  $n$  ways of choosing one object from  $n$  objects.

## Permutation formula

The number of possible permutations of  $r$  objects from a collection of  $n$  **distinct** objects is given by the formula

$$n \times (n - 1) \times \dots \times (n - r + 1)$$

and is denoted by  ${}^n P_r$

$${}^n P_r = \frac{n!}{(n - r)!}$$

### ► Special cases

1.  ${}^n P_0 = \frac{n!}{(n-0)!} = \frac{n!}{n!} = 1$  There is only one ordered arrangement of 0 objects.
2.  ${}^n P_1 = \frac{n!}{(n-1)!} = n$ . There are  $n$  ways of choosing one object from  $n$  objects.
3.  ${}^n P_n =$

## Permutation formula

The number of possible permutations of  $r$  objects from a collection of  $n$  **distinct** objects is given by the formula

$$n \times (n - 1) \times \dots \times (n - r + 1)$$

and is denoted by  ${}^n P_r$

$${}^n P_r = \frac{n!}{(n - r)!}$$

### ► Special cases

1.  ${}^n P_0 = \frac{n!}{(n-0)!} = \frac{n!}{n!} = 1$  There is only one ordered arrangement of 0 objects.
2.  ${}^n P_1 = \frac{n!}{(n-1)!} = n$ . There are  $n$  ways of choosing one object from  $n$  objects.
3.  ${}^n P_n = \frac{n!}{(n-n)!} = \frac{n!}{0!} = n!$ . We can arrange  $n$  distinct objects in  $n!$  ways- multiplication principle of counting.



## Permutation formula

The number of possible permutations of  $r$  objects from a collection of  $n$  **distinct** objects is given by the formula

$$n \times (n - 1) \times \dots \times (n - r + 1)$$

and is denoted by  ${}^n P_r$

$${}^n P_r = \frac{n!}{(n - r)!}$$

### ► Special cases

1.  ${}^n P_0 = \frac{n!}{(n-0)!} = \frac{n!}{n!} = 1$  There is only one ordered arrangement of 0 objects.
2.  ${}^n P_1 = \frac{n!}{(n-1)!} = n$ . There are  $n$  ways of choosing one object from  $n$  objects.
3.  ${}^n P_n = \frac{n!}{(n-n)!} = \frac{n!}{0!} = n!$ . We can arrange  $n$  distinct objects in  $n!$  ways- multiplication principle of counting.

## Example

Take  $A, B, C$ - Possible arrangements- taking all at a time

First place	Second place	Third place
A	B	C
A	C	B
B	A	C
B	C	A
C	A	B
C	B	A

## Example

Take  $A, B, C$ - Possible arrangements- taking all at a time

First place	Second place	Third place
A	B	C
A	C	B
B	A	C
B	C	A
C	A	B
C	B	A

$$n = 3, r = 3, {}^n P_r = \frac{n!}{(n-r)!} = \frac{3!}{0!} = 6$$

## Example

Take  $A, B, C$ - Possible arrangements- taking two at a time

First place	Second place
A	B
A	C
B	A
B	C
C	A
C	B

## Example

Take  $A, B, C$ - Possible arrangements- taking two at a time

First place	Second place
A	B
A	C
B	A
B	C
C	A
C	B

$$n = 3, r = 2, {}^n P_r = \frac{n!}{(n-r)!} = \frac{3!}{1!} = 6$$

# Example

Take A, B, C, D- Possible arrangements- taking all at a time

First place	Second place	Third place	Fourth place
A	B	C	D
A	B	D	C
A	C	B	D
A	C	D	B
A	D	B	C
A	D	C	B
B	A	C	D
B	A	D	C
B	C	A	D
B	C	D	A
B	D	A	C
B	D	C	A
C	A	B	D
C	A	D	B
C	B	A	D
C	B	D	A
C	D	A	B
C	D	B	A
D	A	B	C
D	A	C	B
D	B	A	C
D	B	C	A
D	C	A	B
D	C	B	A

# Example

Take A, B, C, D- Possible arrangements- taking all at a time

First place	Second place	Third place	Fourth place
A	B	C	D
A	B	D	C
A	C	B	D
A	C	D	B
A	D	B	C
A	D	C	B
B	A	C	D
B	A	D	C
B	C	A	D
B	C	D	A
B	D	A	C
B	D	C	A
C	A	B	D
C	A	D	B
C	B	A	D
C	B	D	A
C	D	A	B
C	D	B	A
D	A	B	C
D	A	C	B
D	B	A	C
D	B	C	A
D	C	A	B
D	C	B	A

$$n = 4, r = 4, {}^n P_r = \frac{n!}{(n-r)!} = \frac{4!}{0!} = 24$$

## Example

Take  $A, B, C, D$ - Possible arrangements- taking two at a time

First place	Second place
A	B
A	C
A	D
B	A
B	C
B	D
C	A
C	B
C	D
D	A
D	B
D	C



## Example

Take  $A, B, C, D$ - Possible arrangements- taking two at a time

First place	Second place
A	B
A	C
A	D
B	A
B	C
B	D
C	A
C	B
C	D
D	A
D	B
D	C

$$n = 4, r = 2, {}^n P_r = \frac{n!}{(n-r)!} = \frac{4!}{2!} = 12$$

## Example: application

- ▶ From a committee of 8 persons, in how many ways can we choose a chairman and a vice chairman assuming one person can not hold more than one position?

## Example: application

- ▶ From a committee of 8 persons, in how many ways can we choose a chairman and a vice chairman assuming one person can not hold more than one position?
- ▶  $8 \times 7 =$

## Example: application

- ▶ From a committee of 8 persons, in how many ways can we choose a chairman and a vice chairman assuming one person can not hold more than one position?
- ▶  $8 \times 7 = 56$

## Example: application

- Find the number of 4-digit numbers that can be formed using the digits 1, 2, 3, 4, 5 if no digit is repeated.

## Example: application

- ▶ Find the number of 4-digit numbers that can be formed using the digits 1, 2, 3, 4, 5 if no digit is repeated.
- ▶  $5 \times 4 \times 3 \times 2 \times 1 =$

## Example: application

- ▶ Find the number of 4-digit numbers that can be formed using the digits 1, 2, 3, 4, 5 if no digit is repeated.
- ▶  $5 \times 4 \times 3 \times 2 \times 1 = 120$

## Example: application

- ▶ Find the number of 4-digit numbers that can be formed using the digits 1, 2, 3, 4, 5 if no digit is repeated.
- ▶  $5 \times 4 \times 3 \times 2 \times 1 = 120$
- ▶ How many of these will be even?



## Example: application

- ▶ Find the number of 4-digit numbers that can be formed using the digits 1, 2, 3, 4, 5 if no digit is repeated.
- ▶  $5 \times 4 \times 3 \times 2 \times 1 = 120$
- ▶ How many of these will be even? 48

## Example: application

- ▶ Six people go to the cinema. They sit in a row with ten seats. Find how many ways can this be done if

## Example: application

- ▶ Six people go to the cinema. They sit in a row with ten seats. Find how many ways can this be done if
  - ▶ (i) they can sit anywhere:

## Example: application

- ▶ Six people go to the cinema. They sit in a row with ten seats. Find how many ways can this be done if
  - ▶ (i) they can sit anywhere:  ${}^{10}P_6 = 1,51,200$

## Example: application

- ▶ Six people go to the cinema. They sit in a row with ten seats. Find how many ways can this be done if
  - ▶ (i) they can sit anywhere:  ${}^{10}P_6 = 1,51,200$
  - ▶ (ii) all the empty seats are next to each other:

## Example: application

- ▶ Six people go to the cinema. They sit in a row with ten seats. Find how many ways can this be done if
  - ▶ (i) they can sit anywhere:  ${}^{10}P_6 = 1,51,200$
  - ▶ (ii) all the empty seats are next to each other:  ${}^7P_6 = 5,040$

## Example

## Example

Take  $A, B, C$ - Possible arrangements- taking all at a time



## Example

Take A, B, C- Possible arrangements- taking all at a time

First place	Second place	Third place
A	A	A
A	A	B
A	A	C
A	B	A
A	B	B
A	B	C
A	C	A
A	C	B
A	C	C
B	A	A
B	A	B
B	A	C
B	B	A
B	B	B
B	B	C
B	C	A
B	C	B
B	C	C
C	A	A
C	A	B
C	A	C
C	B	A
C	B	B
C	B	C
C	C	A
C	C	B
C	C	C

## Example

Take  $A, B, C$ - Possible arrangements- taking two at a time

## Example

Take  $A, B, C$ - Possible arrangements- taking two at a time

First place	Second place
A	A
A	B
A	C
B	A
B	B
B	C
C	A
C	B
C	C

## Permutation formula

## Permutation formula

The number of possible permutations of  $r$  objects from a collection of  $n$  **distinct** objects when repetition is allowed is given by the formula

## Permutation formula

The number of possible permutations of  $r$  objects from a collection of  $n$  **distinct** objects when repetition is allowed is given by the formula

$$n \times n \times \dots \times n$$

and is denoted by  $n^r$

## Example

Take  $A, B, C$ - Possible arrangements- taking all at a time.  $n = 3, r = 3, n^r = 27$

## Example

Take  $A, B, C$ - Possible arrangements- taking all at a time.  $n = 3, r = 3, n^r = 27$

First place	Second place	Third place
A	A	A
A	A	B
A	A	C
A	B	A
A	B	B
A	B	C
A	C	A
A	C	B
A	C	C
B	A	A
B	A	B
B	A	C
B	B	A
B	B	B
B	B	C
B	C	A
B	C	B
B	C	C
C	A	A
C	A	B
C	A	C
C	B	A
C	B	B
C	B	C
C	C	A
C	C	B
C	C	C



## Example

Take  $A, B, C$ - Possible arrangements- taking two at a time

First place	Second place
A	A
A	B
A	C
B	A
B	B
B	C
C	A
C	B
C	C

## Example

Take  $A, B, C$ - Possible arrangements- taking two at a time

First place	Second place
A	A
A	B
A	C
B	A
B	B
B	C
C	A
C	B
C	C

$$n = 3, r = 2, n^r = 9$$

## Section summary

1. The number of possible permutations of  $r$  objects from a collection of  $n$  **distinct** objects is given by the formula

$$n \times (n - 1) \times \dots \times (n - r + 1)$$

and is denoted by  ${}^n P_r = \frac{n!}{(n-r)!}$

2. The number of possible permutations of  $r$  objects from a collection of  $n$  **distinct** objects when **repetition** is allowed is given by the formula

$$n \times n \times \dots \times n$$

and is denoted by  $n^r$