



**IIT Madras**  
ONLINE DEGREE

# Sets: Examples

Madhavan Mukund

<https://www.cmi.ac.in/~madhavan>

Mathematics for Data Science 1  
Week 1

# Sets

- A **set** is a collection of items

# Sets

- A **set** is a collection of items
- Finite sets can be listed out explicitly
  - $\{\text{Kohli, Dhoni, Pujara}\}$ ,  $\{1,4,9,16,25\}$


# Sets

- A **set** is a collection of items
- Finite sets can be listed out explicitly
  - $\{\text{Kohli, Dhoni, Pujara}\}, \{1, 4, 9, 16, 25\}$
- Infinite sets cannot be listed out
  - $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$  not formal

# Sets

- A **set** is a collection of items
- Finite sets can be listed out explicitly
  - $\{\text{Kohli, Dhoni, Pujara}\}, \{1,4,9,16,25\}$
- Infinite sets cannot be listed out
  - $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$  not formal
- **Membership**  $x \in X$ , **Subset**  $X \subseteq Y$ 
  - $5 \in \mathbb{Z}$ ,  $\sqrt{2} \notin \mathbb{Q}$
  - $\text{Primes} \subseteq \mathbb{N}$ ,  $\mathbb{N} \subseteq \mathbb{Z}$ ,  $\mathbb{Z} \subseteq \mathbb{Q}$ ,  $\mathbb{Q} \subseteq \mathbb{R}$


Venn Diagram



# Sets

- A **set** is a collection of items
- Finite sets can be listed out explicitly
  - $\{\text{Kohli, Dhoni, Pujara}\}, \{1,4,9,16,25\}$
- Infinite sets cannot be listed out
  - $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$  not formal
- **Membership**  $x \in X$ , **Subset**  $X \subseteq Y$ 
  - $5 \in \mathbb{Z}, \sqrt{2} \notin \mathbb{Q}$
  - $\text{Primes} \subseteq \mathbb{N}, \mathbb{N} \subseteq \mathbb{Z}, \mathbb{Z} \subseteq \mathbb{Q}, \mathbb{Q} \subseteq \mathbb{R}$
- **Powerset** — set of subsets of a set
  - $X = \{a, b\}$ , powerset  $\{\emptyset, \{a\}, \{b\}, \{a, b\}\}$
  - Set with  $n$  elements has  $2^n$  subsets

Venn Diagram



## Set Comprehension

- Squares of the even integers

$$\{x^2 \mid x \in \mathbb{Z}, x \bmod 2 = 0\}$$

$$\{0, 4, 16, 36, 64, 100, 144, 196, 256, \dots\}$$



## Set Comprehension

- Squares of the even integers

$$\{x^2 \mid x \in \mathbb{Z}, x \bmod 2 = 0\}$$

- **Generate** Elements drawn from existing set

$$\{0, 4, 16, 36, 64, 100, 144, 196, 256, \dots\}$$

$$\dots \quad -2 \quad -1 \quad 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \quad \dots$$

# Set Comprehension

## ■ Squares of the even integers

$$\{x^2 \mid x \in \mathbb{Z}, x \bmod 2 = 0\}$$

- **Generate** Elements drawn from existing set
- **Filter** Select elements that satisfy a constraint

$$\{0, 4, 16, 36, 64, 100, 144, 196, 256, \dots\}$$

... -2 -1 0 1 2 3 4 5 ...

... -2 0 2 4 ...

# Set Comprehension

## ■ Squares of the even integers

$$\{x^2 \mid x \in \mathbb{Z}, x \bmod 2 = 0\}$$

- **Generate** Elements drawn from existing set
- **Filter** Select elements that satisfy a constraint
- **Transform** Modify selected elements

$$\{0, 4, 16, 36, 64, 100, 144, 196, 256, \dots\}$$

...    -2    -1    0    1    2    3    4    5

...    -2            0            2            4

...     $(-2)^2$              $0^2$              $2^2$              $4^2$

# Set Comprehension

## ■ Squares of the even integers

$$\{x^2 \mid x \in \mathbb{Z}, x \bmod 2 = 0\}$$

- **Generate** Elements drawn from existing set
- **Filter** Select elements that satisfy a constraint
- **Transform** Modify selected elements

$$\{0, 4, 16, 36, 64, 100, 144, 196, 256, \dots\}$$

...	-2	-1	0	1	2	3	4	5	...
...	-2		0		2		4		...
...	4		0		4		16		...

# Set Comprehension

## ■ Squares of the even integers

$$\{x^2 \mid x \in \mathbb{Z}, x \bmod 2 = 0\}$$

- **Generate** Elements drawn from existing set
- **Filter** Select elements that satisfy a constraint
- **Transform** Modify selected elements

## ■ More filters

$$\{0, 4, 16, 36, 64, 100, 144, 196, 256, \dots\}$$

...	-2	-1	0	1	2	3	4	5	...
...	-2		0		2		4		...
...	4		0		4		16		...

## Set Comprehension

### ■ Squares of the even integers

$$\{x^2 \mid x \in \mathbb{Z}, x \bmod 2 = 0\}$$

$$\{0, 4, 16, 36, 64, 100, 144, 196, 256, \dots\}$$

- **Generate** Elements drawn from existing set
- **Filter** Select elements that satisfy a constraint
- **Transform** Modify selected elements

...	-2	-1	0	1	2	3	4	5	...
...	-2		0		2		4		...
...	4		0		4		16		...

### ■ More filters

- Rationals in reduced form

$$\{p/q \mid p/q \in \mathbb{Q}, \gcd(p, q) = 1\}$$

# Set Comprehension

- Squares of the even integers

$\{x^2 \mid x \in \mathbb{Z}, x \bmod 2 = 0\}$

$\{0, 4, 16, 36, 64, 100, 144, 196, 256, \dots\}$

- **Generate** Elements drawn from existing set
- **Filter** Select elements that satisfy a constraint
- **Transform** Modify selected elements

...	-2	-1	0	1	2	3	4	5	...
...	-2		0		2		4		...
...	4		0		4		16		...

- More filters

- Rationals in reduced form  
 $\{p/q \mid p/q \in \mathbb{Q}, \gcd(p, q) = 1\}$
- Reals in interval  $[-1, 2)$   
 $\{r \mid r \in \mathbb{R}, -1 \leq r < 2\}$

## Set Comprehension ...

- Cubes of first 5 natural numbers

$$Y = \{n^3 \mid n \in \{0, 1, 2, 3, 4\}\}$$



## Set Comprehension ...

- Cubes of first 5 natural numbers

$$Y = \{n^3 \mid n \in \{0, 1, 2, 3, 4\}\}$$

- Cubes of first 500 natural numbers?

$$Y = \{n^3 \mid n \in \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, \dots, 498, 499\}\}$$

## Set Comprehension ...

- Cubes of first 5 natural numbers

$$Y = \{n^3 \mid n \in \{0, 1, 2, 3, 4\}\}$$

- Cubes of first 500 natural numbers?

$$Y = \{n^3 \mid n \in \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, \dots, 498, 499\}\}$$

- Use set comprehension to define first 500 natural numbers

$$X = \{n \mid n \in \mathbb{N}, n < 500\}$$

## Set Comprehension ...

- Cubes of first 5 natural numbers

$$Y = \{n^3 \mid n \in \{0, 1, 2, 3, 4\}\}$$

- Cubes of first 500 natural numbers?

$$Y = \{n^3 \mid n \in \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, \dots, 498, 499\}\}$$

- Use set comprehension to define first 500 natural numbers

$$X = \{n \mid n \in \mathbb{N}, n < 500\}$$

- Now, a more readable version

$$X = \{n \mid n \in \mathbb{N}, n < 500\}$$

$$Y = \{n^3 \mid n \in X\}$$

## Perfect squares

- Integers whose square root is also an integer

$$\{z \mid z \in \mathbb{Z}, \sqrt{z} \in \mathbb{Z}\}$$

## Perfect squares

- Integers whose square root is also an integer  
 $\{z \mid z \in \mathbb{Z}, \sqrt{z} \in \mathbb{Z}\}$
- All squares are positive, so this is the same as  
 $\{n \mid n \in \mathbb{N}, \sqrt{n} \in \mathbb{N}\}$

## Perfect squares

- Integers whose square root is also an integer  
 $\{z \mid z \in \mathbb{Z}, \sqrt{z} \in \mathbb{Z}\}$
- All squares are positive, so this is the same as  
 $\{n \mid n \in \mathbb{N}, \sqrt{n} \in \mathbb{N}\}$
- Alternatively, generate all the perfect squares  
 $\{n^2 \mid n \in \mathbb{N}\}$

## Perfect squares

- Integers whose square root is also an integer  
 $\{z \mid z \in \mathbb{Z}, \sqrt{z} \in \mathbb{Z}\}$
- All squares are positive, so this is the same as  
 $\{n \mid n \in \mathbb{N}, \sqrt{n} \in \mathbb{N}\}$
- Alternatively, generate all the perfect squares  
 $\{n^2 \mid n \in \mathbb{N}\}$
- Extend the definition to rationals
  - $\frac{9}{16} = \left(\frac{3}{4}\right)^2$  is a square,  $\frac{1}{2} \neq \left(\frac{p}{q}\right)^2$  for any  $p, q$  is not

## Perfect squares

- Integers whose square root is also an integer  
 $\{z \mid z \in \mathbb{Z}, \sqrt{z} \in \mathbb{Z}\}$
- All squares are positive, so this is the same as  
 $\{n \mid n \in \mathbb{N}, \sqrt{n} \in \mathbb{N}\}$
- Alternatively, generate all the perfect squares  
 $\{n^2 \mid n \in \mathbb{N}\}$
- Extend the definition to rationals
  - $\frac{9}{16} = \left(\frac{3}{4}\right)^2$  is a square,  $\frac{1}{2} \neq \left(\frac{p}{q}\right)^2$  for any  $p, q$  is not
  - $\{q \mid q \in \mathbb{Q}, \sqrt{q} \in \mathbb{Q}\},$



## Perfect squares

- Integers whose square root is also an integer  
 $\{z \mid z \in \mathbb{Z}, \sqrt{z} \in \mathbb{Z}\}$
- All squares are positive, so this is the same as  
 $\{n \mid n \in \mathbb{N}, \sqrt{n} \in \mathbb{N}\}$
- Alternatively, generate all the perfect squares  
 $\{n^2 \mid n \in \mathbb{N}\}$
- Extend the definition to rationals
  - $\frac{9}{16} = \left(\frac{3}{4}\right)^2$  is a square,  $\frac{1}{2} \neq \left(\frac{p}{q}\right)^2$  for any  $p, q$  is not
  - $\{q \mid q \in \mathbb{Q}, \sqrt{q} \in \mathbb{Q}\}$ , or  $\{q^2 \mid q \in \mathbb{Q}\}$

## Perfect squares

- Integers whose square root is also an integer  
 $\{z \mid z \in \mathbb{Z}, \sqrt{z} \in \mathbb{Z}\}$
- All squares are positive, so this is the same as  
 $\{n \mid n \in \mathbb{N}, \sqrt{n} \in \mathbb{N}\}$
- Alternatively, generate all the perfect squares  
 $\{n^2 \mid n \in \mathbb{N}\}$
- Extend the definition to rationals
  - $\frac{9}{16} = \left(\frac{3}{4}\right)^2$  is a square,  $\frac{1}{2} \neq \left(\frac{p}{q}\right)^2$  for any  $p, q$  is not
  - $\{q \mid q \in \mathbb{Q}, \sqrt{q} \in \mathbb{Q}\}$ , or  $\{q^2 \mid q \in \mathbb{Q}\}$
- Choose the generator as required