



**IIT Madras**  
ONLINE DEGREE

# Relations: Examples

Madhavan Mukund

<https://www.cmi.ac.in/~madhavan>

Mathematics for Data Science 1  
Week 1

# Relations

- $A \times B$  — Cartesian product, all pairs  $(a, b)$ ,  $a \in A$  and  $b \in B$

# Relations

- $A \times B$  — Cartesian product, all pairs  $(a, b)$ ,  $a \in A$  and  $b \in B$
- $A = \{1, 4, 7\}$ ,  $B = \{1, 16, 49\}$

# Relations

- $A \times B$  — Cartesian product, all pairs  $(a, b)$ ,  $a \in A$  and  $b \in B$
- $A = \{1, 4, 7\}$ ,  $B = \{1, 16, 49\}$ 
  - $A \times B = \{(1, 1), (1, 16), (1, 49), (4, 1), (4, 16), (4, 49), (7, 1), (7, 16), (7, 49)\}$

# Relations

- $A \times B$  — Cartesian product, all pairs  $(a, b)$ ,  $a \in A$  and  $b \in B$
- $A = \{1, 4, 7\}$ ,  $B = \{1, 16, 49\}$ 
  - $A \times B = \{(1, 1), (1, 16), (1, 49), (4, 1), (4, 16), (4, 49), (7, 1), (7, 16), (7, 49)\}$
  - $B \times A = \{(1, 1), (16, 1), (49, 1), (1, 4), (16, 4), (49, 4), (1, 7), (16, 7), (49, 7)\}$

# Relations

- $A \times B$  — Cartesian product, all pairs  $(a, b)$ ,  $a \in A$  and  $b \in B$
- $A = \{1, 4, 7\}$ ,  $B = \{1, 16, 49\}$ 
  - $A \times B = \{(1, 1), (1, 16), (1, 49), (4, 1), (4, 16), (4, 49), (7, 1), (7, 16), (7, 49)\}$
  - $B \times A = \{(1, 1), (16, 1), (49, 1), (1, 4), (16, 4), (49, 4), (1, 7), (16, 7), (49, 7)\}$
  - $B \times B = \{(1, 1), (1, 16), (1, 49), (16, 1), (16, 16), (16, 49), (49, 1), (49, 16), (49, 49)\}$

# Relations

- $A \times B$  — Cartesian product, all pairs  $(a, b)$ ,  $a \in A$  and  $b \in B$
- $A = \{1, 4, 7\}$ ,  $B = \{1, 16, 49\}$ 
  - $A \times B = \{(1, 1), (1, 16), (1, 49), (4, 1), (4, 16), (4, 49), (7, 1), (7, 16), (7, 49)\}$
  - $B \times A = \{(1, 1), (16, 1), (49, 1), (1, 4), (16, 4), (49, 4), (1, 7), (16, 7), (49, 7)\}$
  - $B \times B = \{(1, 1), (1, 16), (1, 49), (16, 1), (16, 16), (16, 49), (49, 1), (49, 16), (49, 49)\}$
- Can take Cartesian product of more than two sets



# Relations

- $A \times B$  — Cartesian product, all pairs  $(a, b)$ ,  $a \in A$  and  $b \in B$
- $A = \{1, 4, 7\}$ ,  $B = \{1, 16, 49\}$ 
  - $A \times B = \{(1, 1), (1, 16), (1, 49), (4, 1), (4, 16), (4, 49), (7, 1), (7, 16), (7, 49)\}$
  - $B \times A = \{(1, 1), (16, 1), (49, 1), (1, 4), (16, 4), (49, 4), (1, 7), (16, 7), (49, 7)\}$
  - $B \times B = \{(1, 1), (1, 16), (1, 49), (16, 1), (16, 16), (16, 49), (49, 1), (49, 16), (49, 49)\}$
- Can take Cartesian product of more than two sets
  - $A \times B \times A = \{(1, 1, 1), (1, 1, 4), (1, 1, 7), (1, 16, 1), (1, 16, 7), \dots, (7, 49, 1), (7, 49, 16), (7, 49, 49)\}$

# Relations

- $A \times B$  — Cartesian product, all pairs  $(a, b)$ ,  $a \in A$  and  $b \in B$
- $A = \{1, 4, 7\}$ ,  $B = \{1, 16, 49\}$ 
  - $A \times B = \{(1, 1), (1, 16), (1, 49), (4, 1), (4, 16), (4, 49), (7, 1), (7, 16), (7, 49)\}$
  - $B \times A = \{(1, 1), (16, 1), (49, 1), (1, 4), (16, 4), (49, 4), (1, 7), (16, 7), (49, 7)\}$
  - $B \times B = \{(1, 1), (1, 16), (1, 49), (16, 1), (16, 16), (16, 49), (49, 1), (49, 16), (49, 49)\}$
- Can take Cartesian product of more than two sets
  - $A \times B \times A = \{(1, 1, 1), (1, 1, 4), (1, 1, 7), (1, 16, 1), (1, 16, 7), \dots, (7, 49, 1), (7, 49, 16), (7, 49, 49)\}$
- A relation picks out certain tuples in the Cartesian product

# Relations

- $A \times B$  — Cartesian product, all pairs  $(a, b)$ ,  $a \in A$  and  $b \in B$
- $A = \{1, 4, 7\}$ ,  $B = \{1, 16, 49\}$ 
  - $A \times B = \{(1, 1), (1, 16), (1, 49), (4, 1), (4, 16), (4, 49), (7, 1), (7, 16), (7, 49)\}$
  - $B \times A = \{(1, 1), (16, 1), (49, 1), (1, 4), (16, 4), (49, 4), (1, 7), (16, 7), (49, 7)\}$
  - $B \times B = \{(1, 1), (1, 16), (1, 49), (16, 1), (16, 16), (16, 49), (49, 1), (49, 16), (49, 49)\}$
- Can take Cartesian product of more than two sets
  - $A \times B \times A = \{(1, 1, 1), (1, 1, 4), (1, 1, 7), (1, 16, 1), (1, 16, 7), \dots, (7, 49, 1), (7, 49, 16), (7, 49, 49)\}$
- A relation picks out certain tuples in the Cartesian product
  - $S \subseteq A \times B = \{(1, 1), (4, 16), (7, 49)\}$

# Relations

- $A \times B$  — Cartesian product, all pairs  $(a, b)$ ,  $a \in A$  and  $b \in B$
- $A = \{1, 4, 7\}$ ,  $B = \{1, 16, 49\}$ 
  - $A \times B = \{(1, 1), (1, 16), (1, 49), (4, 1), (4, 16), (4, 49), (7, 1), (7, 16), (7, 49)\}$
  - $B \times A = \{(1, 1), (16, 1), (49, 1), (1, 4), (16, 4), (49, 4), (1, 7), (16, 7), (49, 7)\}$
  - $B \times B = \{(1, 1), (1, 16), (1, 49), (16, 1), (16, 16), (16, 49), (49, 1), (49, 16), (49, 49)\}$
- Can take Cartesian product of more than two sets
  - $A \times B \times A = \{(1, 1, 1), (1, 1, 4), (1, 1, 7), (1, 16, 1), (1, 16, 7), \dots, (7, 49, 1), (7, 49, 16), (7, 49, 49)\}$
- A relation picks out certain tuples in the Cartesian product
  - $S \subseteq A \times B = \{(1, 1), (4, 16), (7, 49)\}$
  - $S = \{(a, b) \mid (a, b) \in A \times B, b = a^2\}$

# Examples of relations

- Divisibility

# Examples of relations

- Divisibility
  - Pairs of natural numbers  $(d, n)$  such that  $d|n$

# Examples of relations

- Divisibility
  - Pairs of natural numbers  $(d, n)$  such that  $d|n$
  - Pairs such as  $(7, 63), (17, 85), (3, 9), \dots$

# Examples of relations

## ■ Divisibility

- Pairs of natural numbers  $(d, n)$  such that  $d|n$
- Pairs such as  $(7, 63), (17, 85), (3, 9), \dots$
- $D = \{(d, n) \mid (d, n) \in \mathbb{N} \times \mathbb{N}, d|n\}$



# Examples of relations

## ■ Divisibility

- Pairs of natural numbers  $(d, n)$  such that  $d|n$
- Pairs such as  $(7, 63), (17, 85), (3, 9), \dots$
- $D = \{(d, n) \mid (d, n) \in \mathbb{N} \times \mathbb{N}, d|n\}$
- Can also extend to integer divisors

# Examples of relations

## ■ Divisibility

- Pairs of natural numbers  $(d, n)$  such that  $d|n$
- Pairs such as  $(7, 63), (17, 85), (3, 9), \dots$
- $D = \{(d, n) \mid (d, n) \in \mathbb{N} \times \mathbb{N}, d|n\}$
- Can also extend to integer divisors
- $E = \{(d, n) \mid (d, n) \in \mathbb{Z} \times \mathbb{N}, d|n\}$

# Examples of relations

## ■ Divisibility

- Pairs of natural numbers  $(d, n)$  such that  $d|n$
- Pairs such as  $(7, 63), (17, 85), (3, 9), \dots$
- $D = \{(d, n) \mid (d, n) \in \mathbb{N} \times \mathbb{N}, d|n\}$
- Can also extend to integer divisors
- $E = \{(d, n) \mid (d, n) \in \mathbb{Z} \times \mathbb{N}, d|n\}$
- Now  $(-7, 63), (-17, 85), (-3, 9), \dots$  are also in  $E$

# Examples of relations

## ■ Divisibility

- Pairs of natural numbers  $(d, n)$  such that  $d|n$
- Pairs such as  $(7, 63), (17, 85), (3, 9), \dots$
- $D = \{(d, n) \mid (d, n) \in \mathbb{N} \times \mathbb{N}, d|n\}$
- Can also extend to integer divisors
- $E = \{(d, n) \mid (d, n) \in \mathbb{Z} \times \mathbb{N}, d|n\}$
- Now  $(-7, 63), (-17, 85), (-3, 9), \dots$  are also in  $E$

## ■ Prime powers

# Examples of relations

## ■ Divisibility

- Pairs of natural numbers  $(d, n)$  such that  $d|n$
- Pairs such as  $(7, 63), (17, 85), (3, 9), \dots$
- $D = \{(d, n) \mid (d, n) \in \mathbb{N} \times \mathbb{N}, d|n\}$
- Can also extend to integer divisors
- $E = \{(d, n) \mid (d, n) \in \mathbb{Z} \times \mathbb{N}, d|n\}$
- Now  $(-7, 63), (-17, 85), (-3, 9), \dots$  are also in  $E$

## ■ Prime powers

- Pairs of natural numbers  $(p, n)$  such that  $p$  is prime and  $n = p^m$  for some natural number  $m$

# Examples of relations

## ■ Divisibility

- Pairs of natural numbers  $(d, n)$  such that  $d|n$
- Pairs such as  $(7, 63), (17, 85), (3, 9), \dots$
- $D = \{(d, n) \mid (d, n) \in \mathbb{N} \times \mathbb{N}, d|n\}$
- Can also extend to integer divisors
- $E = \{(d, n) \mid (d, n) \in \mathbb{Z} \times \mathbb{N}, d|n\}$
- Now  $(-7, 63), (-17, 85), (-3, 9), \dots$  are also in  $E$

## ■ Prime powers

- Pairs of natural numbers  $(p, n)$  such that  $p$  is prime and  $n = p^m$  for some natural number  $m$
- Examples:  $(3, 1), (5, 625), (7, 343), \dots$

# Examples of relations

## ■ Divisibility

- Pairs of natural numbers  $(d, n)$  such that  $d|n$
- Pairs such as  $(7, 63), (17, 85), (3, 9), \dots$
- $D = \{(d, n) \mid (d, n) \in \mathbb{N} \times \mathbb{N}, d|n\}$
- Can also extend to integer divisors
- $E = \{(d, n) \mid (d, n) \in \mathbb{Z} \times \mathbb{N}, d|n\}$
- Now  $(-7, 63), (-17, 85), (-3, 9), \dots$  are also in  $E$

## ■ Prime powers

- Pairs of natural numbers  $(p, n)$  such that  $p$  is prime and  $n = p^m$  for some natural number  $m$
- Examples:  $(3, 1), (5, 625), (7, 343), \dots$
- First define primes:  $P = \{p \mid p \in \mathbb{N}, \text{factors}(p) = \{1, p\}, p \neq 1\}$

# Examples of relations

## ■ Divisibility

- Pairs of natural numbers  $(d, n)$  such that  $d|n$
- Pairs such as  $(7, 63), (17, 85), (3, 9), \dots$
- $D = \{(d, n) \mid (d, n) \in \mathbb{N} \times \mathbb{N}, d|n\}$
- Can also extend to integer divisors
- $E = \{(d, n) \mid (d, n) \in \mathbb{Z} \times \mathbb{N}, d|n\}$
- Now  $(-7, 63), (-17, 85), (-3, 9), \dots$  are also in  $E$

## ■ Prime powers

- Pairs of natural numbers  $(p, n)$  such that  $p$  is prime and  $n = p^m$  for some natural number  $m$
- Examples:  $(3, 1), (5, 625), (7, 343), \dots$
- First define primes:  $P = \{p \mid p \in \mathbb{N}, \text{factors}(p) = \{1, p\}, p \neq 1\}$
- Prime powers:  $PP = \{(p, n) \mid (p, n) \in P \times \mathbb{N}, n = p^m \text{ for some } m \in \mathbb{N}\}$



# Beyond numbers

## Airline routes

- An airline flies to set of cities — e.g. Bangalore, Chennai, Delhi, Kolkata, ...

# Beyond numbers

## Airline routes

- An airline flies to set of cities — e.g. Bangalore, Chennai, Delhi, Kolkata, ...
- Let  $C$  denote the set of cities served by the airline

# Beyond numbers

## Airline routes

- An airline flies to set of cities — e.g. Bangalore, Chennai, Delhi, Kolkata, ...
- Let  $C$  denote the set of cities served by the airline
- Some cities are connected by direct flights

# Beyond numbers

## Airline routes

- An airline flies to set of cities — e.g. Bangalore, Chennai, Delhi, Kolkata, ...
- Let  $C$  denote the set of cities served by the airline
- Some cities are connected by direct flights
- $D \subseteq C \times C$

# Beyond numbers

## Airline routes

- An airline flies to set of cities — e.g. Bangalore, Chennai, Delhi, Kolkata, ...
- Let  $C$  denote the set of cities served by the airline
- Some cities are connected by direct flights
- $D \subseteq C \times C$
- Is  $D$  reflexive, irreflexive?

# Beyond numbers

## Airline routes

- An airline flies to set of cities — e.g. Bangalore, Chennai, Delhi, Kolkata, ...
- Let  $C$  denote the set of cities served by the airline
- Some cities are connected by direct flights
- $D \subseteq C \times C$
- Is  $D$  reflexive, irreflexive?
  - Hopefully irreflexive!

# Beyond numbers

## Airline routes

- An airline flies to set of cities — e.g. Bangalore, Chennai, Delhi, Kolkata, ...
- Let  $C$  denote the set of cities served by the airline
- Some cities are connected by direct flights
- $D \subseteq C \times C$
- Is  $D$  reflexive, irreflexive?
  - Hopefully irreflexive!
- Is  $D$  symmetric?

# Beyond numbers

## Airline routes

- An airline flies to set of cities — e.g. Bangalore, Chennai, Delhi, Kolkata, ...
- Let  $C$  denote the set of cities served by the airline
- Some cities are connected by direct flights
- $D \subseteq C \times C$
- Is  $D$  reflexive, irreflexive?
  - Hopefully irreflexive!
- Is  $D$  symmetric?
  - If there is a direct flight from Bangalore to Delhi, is there always a direct flight back from Delhi to Bangalore



# Beyond numbers

## Airline routes

- An airline flies to set of cities — e.g. Bangalore, Chennai, Delhi, Kolkata, ...
- Let  $C$  denote the set of cities served by the airline
- Some cities are connected by direct flights
- $D \subseteq C \times C$
- Is  $D$  reflexive, irreflexive?
  - Hopefully irreflexive!
- Is  $D$  symmetric?
  - If there is a direct flight from Bangalore to Delhi, is there always a direct flight back from Delhi to Bangalore
  - For bigger cities, yes

# Beyond numbers

## Airline routes

- An airline flies to set of cities — e.g. Bangalore, Chennai, Delhi, Kolkata, ...
- Let  $C$  denote the set of cities served by the airline
- Some cities are connected by direct flights
- $D \subseteq C \times C$
- Is  $D$  reflexive, irreflexive?
  - Hopefully irreflexive!
- Is  $D$  symmetric?
  - If there is a direct flight from Bangalore to Delhi, is there always a direct flight back from Delhi to Bangalore
  - For bigger cities, yes
  - For smaller cities, may have a triangular route  
Chennai  $\rightarrow$  Madurai  $\rightarrow$  Salem  $\rightarrow$  Chennai

# Tables as relations

## ■ Flying distances between cities

Source	Destination	Distance (km)
Bangalore	Chennai	290
Chennai	Delhi	1752
Delhi	Bangalore	1735
Delhi	Chennai	1752
...	...	...

# Tables as relations

- Flying distances between cities

Source	Destination	Distance (km)
Bangalore	Chennai	290
Chennai	Delhi	1752
Delhi	Bangalore	1735
Delhi	Chennai	1752
...	...	...

- Table is a relation:  $Dist \subseteq C \times C \times \mathbb{N}$

# Tables as relations

- Flying distances between cities

Source	Destination	Distance (km)
Bangalore	Chennai	290
Chennai	Delhi	1752
Delhi	Bangalore	1735
Delhi	Chennai	1752
...	...	...

- Table is a relation:  $Dist \subseteq C \times C \times \mathbb{N}$
- Some entries are useless: (Delhi,Delhi,0)

# Tables as relations

- Flying distances between cities

Source	Destination	Distance (km)
Bangalore	Chennai	290
Chennai	Delhi	1752
Delhi	Bangalore	1735
Delhi	Chennai	1752
...	...	...

- Table is a relation:  $Dist \subseteq C \times C \times \mathbb{N}$
- Some entries are useless: (Delhi,Delhi,0)
- Restrict to cities served by direct flights  
 $Dist = \{(a, b, d) \mid (a, b) \in D, d \text{ is distance from } a \text{ to } b\}$

# Tables as relations

- Flying distances between cities

Source	Destination	Distance (km)
Bangalore	Chennai	290
Chennai	Delhi	1752
Delhi	Bangalore	1735
Delhi	Chennai	1752
...	...	...

- Table is a relation:  $Dist \subseteq C \times C \times \mathbb{N}$
- Some entries are useless: (Delhi,Delhi,0)
- Restrict to cities served by direct flights  
 $Dist = \{(a, b, d) \mid (a, b) \in D, d \text{ is distance from } a \text{ to } b\}$
- Distances are symmetric, even if  $D$  is not

# Tables as relations

- Flying distances between cities

Source	Destination	Distance (km)
Bangalore	Chennai	290
Chennai	Delhi	1752
Delhi	Bangalore	1735
Delhi	Chennai	1752
...	...	...

- Table is a relation:  $Dist \subseteq C \times C \times \mathbb{N}$
- Some entries are useless: (Delhi,Delhi,0)
- Restrict to cities served by direct flights  
 $Dist = \{(a, b, d) \mid (a, b) \in D, d \text{ is distance from } a \text{ to } b\}$
- Distances are symmetric, even if  $D$  is not
- Save space by representing only one direction in the table



## Tables as relations ...

<b>Roll no</b>	<b>Name</b>	<b>Date of birth)</b>
A71396	Abhay Shah	03-07-2001
B82976	Payal Ghosh	18-06-1999
F98989	Jeremy Pinto	22-02-2003
C93986	Payal Ghosh	14-05-2000
...	...	...

## Tables as relations ...

<b>Roll no</b>	<b>Name</b>	<b>Date of birth)</b>
A71396	Abhay Shah	03-07-2001
B82976	Payal Ghosh	18-06-1999
F98989	Jeremy Pinto	22-02-2003
C93986	Payal Ghosh	14-05-2000
...	...	...

- Some columns are special — each student has a unique roll number

## Tables as relations ...

Roll no	Name	Date of birth)
A71396	Abhay Shah	03-07-2001
B82976	Payal Ghosh	18-06-1999
F98989	Jeremy Pinto	22-02-2003
C93986	Payal Ghosh	14-05-2000
...	...	...

- Some columns are special — each student has a unique roll number
  - Such a column is called a **key**

# Tables as relations ...

Roll no	Name	Date of birth)
A71396	Abhay Shah	03-07-2001
B82976	Payal Ghosh	18-06-1999
F98989	Jeremy Pinto	22-02-2003
C93986	Payal Ghosh	14-05-2000
...	...	...

- Some columns are special — each student has a unique roll number
  - Such a column is called a **key**
  - Name is not a key, in general

## Tables as relations ...

Roll no	Name	Date of birth)
A71396	Abhay Shah	03-07-2001
B82976	Payal Ghosh	18-06-1999
F98989	Jeremy Pinto	22-02-2003
C93986	Payal Ghosh	14-05-2000
...	...	...

- Some columns are special — each student has a unique roll number
  - Such a column is called a **key**
  - Name is not a key, in general
- Given the roll number, can retrieve the data for a student

# Tables as relations ...

Roll no	Name	Date of birth)
A71396	Abhay Shah	03-07-2001
B82976	Payal Ghosh	18-06-1999
F98989	Jeremy Pinto	22-02-2003
C93986	Payal Ghosh	14-05-2000
...	...	...

- Some columns are special — each student has a unique roll number
  - Such a column is called a **key**
  - Name is not a key, in general
- Given the roll number, can retrieve the data for a student
  - **Function** from Roll Numbers to (Name, Date of Birth)

# Tables as relations ...

Roll no	Name	Date of birth)
A71396	Abhay Shah	03-07-2001
B82976	Payal Ghosh	18-06-1999
F98989	Jeremy Pinto	22-02-2003
C93986	Payal Ghosh	14-05-2000
...	...	...

- Some columns are special — each student has a unique roll number
  - Such a column is called a **key**
  - Name is not a key, in general
- Given the roll number, can retrieve the data for a student
  - **Function** from Roll Numbers to (Name, Date of Birth)
  - **(key,value)** pairs

# Operations on relations

Roll No	Name	Date of birth)
A71396	Abhay Shah	03-07-2001
B82976	Payal Ghosh	18-06-1999
F98989	Jeremy Pinto	22-02-2003
C93986	Payal Ghosh	14-05-2000
...	...	...

Roll no	Subject	Grade)
A71396	English	B
B82976	Mathematics	A
C93986	Physics	B
B82976	Chemistry	A
...	...	...



# Operations on relations

Roll No	Name	Date of birth)
A71396	Abhay Shah	03-07-2001
B82976	Payal Ghosh	18-06-1999
F98989	Jeremy Pinto	22-02-2003
C93986	Payal Ghosh	14-05-2000
...	...	...

Roll no	Subject	Grade)
A71396	English	B
B82976	Mathematics	A
C93986	Physics	B
B82976	Chemistry	A
...	...	...

- Generate a table with roll numbers, names and grades

# Operations on relations

Roll No	Name	Date of birth)
A71396	Abhay Shah	03-07-2001
B82976	Payal Ghosh	18-06-1999
F98989	Jeremy Pinto	22-02-2003
C93986	Payal Ghosh	14-05-2000
...	...	...

Roll no	Subject	Grade)
A71396	English	B
B82976	Mathematics	A
C93986	Physics	B
B82976	Chemistry	A
...	...	...

- Generate a table with roll numbers, names and grades

Roll No	Name	Subject	Grade)
A71396	Abhay Shah	English	B
B82976	Payal Ghosh	Mathematics	A
B82976	Payal Ghosh	Chemistry	A
C93986	Payal Ghosh	Physics	B
...	...	...	...

# Operations on relations

Roll No	Name	Date of birth)
A71396	Abhay Shah	03-07-2001
B82976	Payal Ghosh	18-06-1999
F98989	Jeremy Pinto	22-02-2003
C93986	Payal Ghosh	14-05-2000
...	...	...

Roll no	Subject	Grade)
A71396	English	B
B82976	Mathematics	A
C93986	Physics	B
B82976	Chemistry	A
...	...	...

- Generate a table with roll numbers, names and grades
- **Join** the relations on Roll No

Roll No	Name	Subject	Grade)
A71396	Abhay Shah	English	B
B82976	Payal Ghosh	Mathematics	A
B82976	Payal Ghosh	Chemistry	A
C93986	Payal Ghosh	Physics	B
...	...	...	...

# Operations on relations

Roll No	Name	Date of birth)
A71396	Abhay Shah	03-07-2001
B82976	Payal Ghosh	18-06-1999
F98989	Jeremy Pinto	22-02-2003
C93986	Payal Ghosh	14-05-2000
...	...	...

Roll no	Subject	Grade)
A71396	English	B
B82976	Mathematics	A
C93986	Physics	B
B82976	Chemistry	A
...	...	...

- Generate a table with roll numbers, names and grades

- **Join** the relations on Roll No

- $\{(r, n, s, g) \mid$   
     $(r, n, d) \in \text{Students},$   
     $(r', s, g) \in \text{Grades},$   
     $r = r'\}$

Roll No	Name	Subject	Grade)
A71396	Abhay Shah	English	B
B82976	Payal Ghosh	Mathematics	A
B82976	Payal Ghosh	Chemistry	A
C93986	Payal Ghosh	Physics	B
...	...	...	...

# Summary

- A relation describes special tuples in a Cartesian product
- Data tables are essentially relations
- Combining information on tables can be described in terms of operations on relations