

Statistics for Data Science -1

Lecture 5.5: Combinations

Usha Mohan

Indian Institute of Technology Madras

Learning objectives

1. Understand basic principles of counting.
2. Concept of factorials.
3. Understand differences between counting with order (permutation) and counting without regard to order (combination).
4. Use permutations and combinations to answer real life applications.

Combinations

Applications: Permutations or combinations

Introduction

- ▶ Example: How many ways can we select two students from a group of three students?

Introduction

- ▶ Example: How many ways can we select two students from a group of three students?
 - ▶ Let A , B , and C be the three students.
 - ▶ We can chose AB , AC , or BC .
 - ▶ Note, when we talked of permutations, the order was important, i.e., AB was different from BA .
 - ▶ In this case, they are the same- order is not important.

Introduction

- ▶ Example: How many ways can we select two students from a group of three students?
 - ▶ Let A , B , and C be the three students.
 - ▶ We can choose AB , AC , or BC .
 - ▶ Note, when we talked of permutations, the order was important, i.e., AB was different from BA .
 - ▶ In this case, they are the same- order is not important.
- ▶ Each selection is called a **combination** of 3 different objects taken 2 at a time.

Introduction

- ▶ Example: How many ways can we select two students from a group of three students?
 - ▶ Let A , B , and C be the three students.
 - ▶ We can choose AB , AC , or BC .
 - ▶ Note, when we talked of permutations, the order was important, i.e., AB was different from BA .
 - ▶ In this case, they are the same- order is not important.
- ▶ Each selection is called a **combination** of 3 different objects taken 2 at a time.
- ▶ In this case, the concern is only which of the 2 objects are chosen and not in the order in which they are chosen.

Example

Consider A, B, C - Possible combinations- taking two at a time

First place	Second place
A	B
A	C
B	C

Example

Consider A, B, C - Possible combinations- taking two at a time

First place	Second place
A	B
A	C
B	C

- Note each combination gives rise to $2!$ arrangements.

Example

Consider A, B, C - Possible combinations- taking two at a time

First place	Second place
A	B
A	C
B	C

- ▶ Note each combination gives rise to $2!$ arrangements.
- ▶ All combinations give $3 \times 2 = 6$ arrangements.

Example

Consider A, B, C - Possible combinations- taking two at a time

First place	Second place
A	B
A	C
B	C

- ▶ Note each combination gives rise to $2!$ arrangements.
- ▶ All combinations give $3 \times 2 = 6$ arrangements.
- ▶ Number of combinations $\times 2! =$

Example

Consider A, B, C - Possible combinations- taking two at a time

First place	Second place
A	B
A	C
B	C

- ▶ Note each combination gives rise to $2!$ arrangements.
- ▶ All combinations give $3 \times 2 = 6$ arrangements.
- ▶ Number of combinations $\times 2! =$ Number of permutations

Combinations: Notation and formula

- ▶ In general, each combination of r objects from n objects can give rise to $r!$ arrangements.

Combinations: Notation and formula

- ▶ In general, each combination of r objects from n objects can give rise to $r!$ arrangements.
- ▶ The number of possible combinations of r objects

Combinations: Notation and formula

- ▶ In general, each combination of r objects from n objects can give rise to $r!$ arrangements.
- ▶ The number of possible combinations of r objects from a collection of n **distinct** objects is denoted by nC_r and is given by

Combinations: Notation and formula

- ▶ In general, each combination of r objects from n objects can give rise to $r!$ arrangements.
- ▶ The number of possible combinations of r objects from a collection of n **distinct** objects is denoted by nC_r and is given by

$${}^nC_r =$$

Combinations: Notation and formula

- ▶ In general, each combination of r objects from n objects can give rise to $r!$ arrangements.
- ▶ The number of possible combinations of r objects from a collection of n **distinct** objects is denoted by nC_r and is given by

$${}^nC_r = \frac{n!}{r!(n-r)!}$$

Combinations: Notation and formula

- ▶ In general, each combination of r objects from n objects can give rise to $r!$ arrangements.
- ▶ The number of possible combinations of r objects from a collection of n **distinct** objects is denoted by nC_r and is given by

$${}^nC_r = \frac{n!}{r!(n-r)!}$$

- ▶ Another common notation is $\binom{n}{r}$ which is also referred to as the **binomial coefficient**

Combinations: Notation and formula

- ▶ In general, each combination of r objects from n objects can give rise to $r!$ arrangements.
- ▶ The number of possible combinations of r objects from a collection of n **distinct** objects is denoted by nC_r and is given by

$${}^nC_r = \frac{n!}{r!(n-r)!}$$

- ▶ Another common notation is $\binom{n}{r}$ which is also referred to as the **binomial coefficient**

Some useful results

Some useful results

1. ${}^nC_r =$

Some useful results

$$1. {}^nC_r = \frac{n!}{r!(n-r)!} =$$

Some useful results

$$1. \quad {}^nC_r = \frac{n!}{r!(n-r)!} = \frac{n!}{(n-r)!r!} =$$

Some useful results

$$1. \quad {}^nC_r = \frac{n!}{r!(n-r)!} = \frac{n!}{(n-r)!r!} = {}^nC_{(n-r)}$$

Some useful results

1. ${}^nC_r = \frac{n!}{r!(n-r)!} = \frac{n!}{(n-r)!r!} = {}^nC_{(n-r)}$

In other words, selecting r objects from n objects is the same as rejecting $n - r$ objects from n objects.

Some useful results

1. ${}^nC_r = \frac{n!}{r!(n-r)!} = \frac{n!}{(n-r)!r!} = {}^nC_{(n-r)}$

In other words, selecting r objects from n objects is the same as rejecting $n - r$ objects from n objects.

2. ${}^nC_n = 1$ and ${}^nC_0 = 1$ for all values of n

Some useful results

1. ${}^nC_r = \frac{n!}{r!(n-r)!} = \frac{n!}{(n-r)!r!} = {}^nC_{(n-r)}$

In other words, selecting r objects from n objects is the same as rejecting $n - r$ objects from n objects.

2. ${}^nC_n = 1$ and ${}^nC_0 = 1$ for all values of n

3. ${}^nC_r = {}^{n-1}C_{r-1} + {}^{n-1}C_r; 1 \leq r \leq n$

Example: Choosing questions in an exam

- ▶ In an examination, a question paper consists of 12 questions divided into two parts i.e., Part I and Part II, containing 7 and 5 questions, respectively. A student is required to attempt 8 questions in all, selecting at least 3 from each part. In how many ways can a student select the questions ?

Example: Choosing questions in an exam

- ▶ In an examination, a question paper consists of 12 questions divided into two parts i.e., Part I and Part II, containing 7 and 5 questions, respectively. A student is required to attempt 8 questions in all, selecting at least 3 from each part. In how many ways can a student select the questions ?
- ▶ Solution: ${}^7C_3 {}^5C_5$

Example: Choosing questions in an exam

- ▶ In an examination, a question paper consists of 12 questions divided into two parts i.e., Part I and Part II, containing 7 and 5 questions, respectively. A student is required to attempt 8 questions in all, selecting at least 3 from each part. In how many ways can a student select the questions ?
- ▶ Solution: ${}^7C_3 {}^5C_5 + {}^7C_4 {}^5C_4$

Example: Choosing questions in an exam


- ▶ In an examination, a question paper consists of 12 questions divided into two parts i.e., Part I and Part II, containing 7 and 5 questions, respectively. A student is required to attempt 8 questions in all, selecting at least 3 from each part. In how many ways can a student select the questions ?
- ▶ Solution: ${}^7C_3 {}^5C_5 + {}^7C_4 {}^5C_4 + {}^7C_5 {}^5C_3$

Example: Choosing questions in an exam

- ▶ In an examination, a question paper consists of 12 questions divided into two parts i.e., Part I and Part II, containing 7 and 5 questions, respectively. A student is required to attempt 8 questions in all, selecting at least 3 from each part. In how many ways can a student select the questions ?
- ▶ Solution: ${}^7C_3 {}^5C_5 + {}^7C_4 {}^5C_4 + {}^7C_5 {}^5C_3 = 35 + 175 + 210 = 420$

Example: Game of cards

Lets consider the case of choosing four cards from a deck of 52 cards.



Example: Game of cards contd.

1. Total number of ways of choosing four cards from 52 cards =

Example: Game of cards contd.

1. Total number of ways of choosing four cards from 52 cards =
$${}^{52}C_4 = \frac{52!}{4!48!} =$$

Example: Game of cards contd.

1. Total number of ways of choosing four cards from 52 cards =
$${}^{52}C_4 = \frac{52!}{4!48!} = 2,70,725$$

Example: Game of cards contd.

1. Total number of ways of choosing four cards from 52 cards =
$${}^{52}C_4 = \frac{52!}{4!48!} = 2,70,725$$
2. All four cards are of the same suit

Example: Game of cards contd.

1. Total number of ways of choosing four cards from 52 cards =
 ${}^{52}C_4 = \frac{52!}{4!48!} = 2,70,725$
2. All four cards are of the same suit
 ${}^4C_1 \times {}^{13}C_4 =$

Example: Game of cards contd.

1. Total number of ways of choosing four cards from 52 cards =
$${}^{52}C_4 = \frac{52!}{4!48!} = 2,70,725$$
2. All four cards are of the same suit
$${}^4C_1 \times {}^{13}C_4 = 4 \times \frac{13!}{4!9!} =$$

Example: Game of cards contd.

1. Total number of ways of choosing four cards from 52 cards =
$${}^{52}C_4 = \frac{52!}{4!48!} = 2,70,725$$
2. All four cards are of the same suit
$${}^4C_1 \times {}^{13}C_4 = 4 \times \frac{13!}{4!9!} = 2860$$

Example: Game of cards contd.

1. Total number of ways of choosing four cards from 52 cards =
$${}^{52}C_4 = \frac{52!}{4!48!} = 2,70,725$$
2. All four cards are of the same suit
$${}^4C_1 \times {}^{13}C_4 = 4 \times \frac{13!}{4!9!} = 2860$$
3. Cards are of same colour

Example: Game of cards contd.

1. Total number of ways of choosing four cards from 52 cards =
$${}^{52}C_4 = \frac{52!}{4!48!} = 2,70,725$$
2. All four cards are of the same suit
$${}^4C_1 \times {}^{13}C_4 = 4 \times \frac{13!}{4!9!} = 2860$$
3. Cards are of same colour ${}^2C_1 \times {}^{26}C_4 =$

Example: Game of cards contd.

1. Total number of ways of choosing four cards from 52 cards =
 ${}^{52}C_4 = \frac{52!}{4!48!} = 2,70,725$
2. All four cards are of the same suit
 ${}^4C_1 \times {}^{13}C_4 = 4 \times \frac{13!}{4!9!} = 2860$
3. Cards are of same colour ${}^2C_1 \times {}^{26}C_4 = 2 \times \frac{26!}{4!22!} =$

Example: Game of cards contd.

1. Total number of ways of choosing four cards from 52 cards =
$${}^{52}C_4 = \frac{52!}{4!48!} = 2,70,725$$
2. All four cards are of the same suit
$${}^4C_1 \times {}^{13}C_4 = 4 \times \frac{13!}{4!9!} = 2860$$
3. Cards are of same colour ${}^2C_1 \times {}^{26}C_4 = 2 \times \frac{26!}{4!22!} = 2,99,00$

Example: Choosing a cricket team

- ▶ Select a cricket team of eleven from 17 players in which only 5 players can bowl. The requirement is the cricket team of 11 must include exactly 4 bowlers? How many ways can the selection be done?

Example: Choosing a cricket team

- ▶ Select a cricket team of eleven from 17 players in which only 5 players can bowl. The requirement is the cricket team of 11 must include exactly 4 bowlers? How many ways can the selection be done?
- ▶ Solution:
 - ▶ Total number of players available for selection: 17
Number of bowlers: 5

Example: Choosing a cricket team

- ▶ Select a cricket team of eleven from 17 players in which only 5 players can bowl. The requirement is the cricket team of 11 must include exactly 4 bowlers? How many ways can the selection be done?
- ▶ Solution:
 - ▶ Total number of players available for selection: 17
Number of bowlers: 5
 - ▶ Need four bowlers: This selection can be done in 5C_4 ways.

Example: Choosing a cricket team

- ▶ Select a cricket team of eleven from 17 players in which only 5 players can bowl. The requirement is the cricket team of 11 must include exactly 4 bowlers? How many ways can the selection be done?
- ▶ Solution:
 - ▶ Total number of players available for selection: 17
Number of bowlers: 5
 - ▶ Need four bowlers: This selection can be done in 5C_4 ways.
 - ▶ Remaining seven players can be selected from remaining twelve players in ${}^{12}C_7$ ways.

Example: Choosing a cricket team

- ▶ Select a cricket team of eleven from 17 players in which only 5 players can bowl. The requirement is the cricket team of 11 must include exactly 4 bowlers? How many ways can the selection be done?
- ▶ Solution:
 - ▶ Total number of players available for selection: 17
Number of bowlers: 5
 - ▶ Need four bowlers: This selection can be done in 5C_4 ways.
 - ▶ Remaining seven players can be selected from remaining twelve players in ${}^{12}C_7$ ways.
 - ▶ Total number of ways the selection can be done is ${}^5C_4 \times {}^{12}C_7 =$

Example: Choosing a cricket team

- ▶ Select a cricket team of eleven from 17 players in which only 5 players can bowl. The requirement is the cricket team of 11 must include exactly 4 bowlers? How many ways can the selection be done?
- ▶ Solution:
 - ▶ Total number of players available for selection: 17
Number of bowlers: 5
 - ▶ Need four bowlers: This selection can be done in 5C_4 ways.
 - ▶ Remaining seven players can be selected from remaining twelve players in ${}^{12}C_7$ ways.
 - ▶ Total number of ways the selection can be done is
 ${}^5C_4 \times {}^{12}C_7 = 5 \times 792 =$

Example: Choosing a cricket team

- ▶ Select a cricket team of eleven from 17 players in which only 5 players can bowl. The requirement is the cricket team of 11 must include exactly 4 bowlers? How many ways can the selection be done?
- ▶ Solution:
 - ▶ Total number of players available for selection: 17
Number of bowlers: 5
 - ▶ Need four bowlers: This selection can be done in 5C_4 ways.
 - ▶ Remaining seven players can be selected from remaining twelve players in ${}^{12}C_7$ ways.
 - ▶ Total number of ways the selection can be done is ${}^5C_4 \times {}^{12}C_7 = 5 \times 792 = 3960$ ways

Example: Drawing lines in a circle

- ▶ Given n points on a circle, how many lines can be drawn connecting these points?

Example: Drawing lines in a circle

- ▶ Given n points on a circle, how many lines can be drawn connecting these points?
- ▶ $n = 2$ points, one line can be drawn connecting the points



line segment: AB

Example: Drawing lines in a circle

- ▶ Given n points on a circle, how many lines can be drawn connecting these points?
- ▶ $n = 2$ points, one line can be drawn connecting the points



line segment: AB

- ▶ $n = 3$ points, three line can be drawn connecting the points



line segments: AB , AC , and BC

Example: Drawing lines in a circle

- ▶ Given n points on a circle, how many lines can be drawn connecting these points?
- ▶ $n = 2$ points, one line can be drawn connecting the points



line segment: AB

- ▶ $n = 3$ points, three line can be drawn connecting the points



line segments: AB , AC , and BC

- ▶ In general, given n points, number of line segments that can be drawn connecting the points is nC_2

Section summary

1. Notation and formula for selecting r objects from n objects.
2. Some useful combinatorial identities.