#### Statistics for Data Science -1

Lecture 7.5: Conditional Probability: Independent events-examples

Usha Mohan

Indian Institute of Technology Madras

### Learning objectives

1. Understand notion of conditional probability, i.e find the probability of an event given another event has occurred.

### Learning objectives

- 1. Understand notion of conditional probability, i.e find the probability of an event given another event has occurred.
- 2. Distinguish between independent and dependent events.

### Learning objectives

- 1. Understand notion of conditional probability, i.e find the probability of an event given another event has occurred.
- 2. Distinguish between independent and dependent events.
- 3. Solve applications of probability.

Independent events: example Rolling a dice Deck of cards

Experiment: Roll a dice twice

- Experiment: Roll a dice twice
- Sample space:

$$S = \left\{ \begin{array}{l} (1,1), (1,2), (1,3), (1,4), (1,5), (1,6), \\ (2,1), (2,2), (2,3), (2,4), (2,5), (2,6), \\ (3,1), (3,2), (3,3), (3,4), (3,5), (3,6), \\ (4,1), (4,2), (4,3), (4,4), (4,5), (4,6), \\ (5,1), (5,2), (5,3), (5,4), (5,5), (5,6), \\ (6,1), (6,2), (6,3), (6,4), (6,5), (6,6), \end{array} \right\}$$

- Experiment: Roll a dice twice
- Sample space:

$$S = \left\{ \begin{array}{l} (1,1), (1,2), (1,3), (1,4), (1,5), (1,6), \\ (2,1), (2,2), (2,3), (2,4), (2,5), (2,6), \\ (3,1), (3,2), (3,3), (3,4), (3,5), (3,6), \\ (4,1), (4,2), (4,3), (4,4), (4,5), (4,6), \\ (5,1), (5,2), (5,3), (5,4), (5,5), (5,6), \\ (6,1), (6,2), (6,3), (6,4), (6,5), (6,6), \end{array} \right\}$$

- ► Define the following events
  - $\triangleright$   $E_1$ : The first outcome is a 3
  - $\triangleright$   $E_2$ : Sum of outcomes is 8
  - ► E<sub>3</sub>: Sum of outcomes is 7

- Experiment: Roll a dice twice
- Sample space:

$$S = \left\{ \begin{array}{l} (1,1), (1,2), (1,3), (1,4), (1,5), (1,6), \\ (2,1), (2,2), (2,3), (2,4), (2,5), (2,6), \\ (3,1), (3,2), (3,3), (3,4), (3,5), (3,6), \\ (4,1), (4,2), (4,3), (4,4), (4,5), (4,6), \\ (5,1), (5,2), (5,3), (5,4), (5,5), (5,6), \\ (6,1), (6,2), (6,3), (6,4), (6,5), (6,6), \end{array} \right\}$$

- Define the following events
  - $ightharpoonup E_1$ : The first outcome is a 3
  - ► *E*<sub>2</sub>: Sum of outcomes is 8
  - E<sub>3</sub>: Sum of outcomes is 7
- $\triangleright$  Are events  $E_1$  and  $E_2$  independent?

- Experiment: Roll a dice twice
- Sample space:

$$S = \left\{ \begin{array}{l} (1,1), (1,2), (1,3), (1,4), (1,5), (1,6), \\ (2,1), (2,2), (2,3), (2,4), (2,5), (2,6), \\ (3,1), (3,2), (3,3), (3,4), (3,5), (3,6), \\ (4,1), (4,2), (4,3), (4,4), (4,5), (4,6), \\ (5,1), (5,2), (5,3), (5,4), (5,5), (5,6), \\ (6,1), (6,2), (6,3), (6,4), (6,5), (6,6), \end{array} \right\}$$

- Define the following events
  - $ightharpoonup E_1$ : The first outcome is a 3
  - ► *E*<sub>2</sub>: Sum of outcomes is 8
  - E<sub>3</sub>: Sum of outcomes is 7
- Are events E<sub>1</sub> and E<sub>2</sub> independent?
- $\triangleright$  Are events  $E_1$  and  $E_3$  independent?

$$P(E_1 \cap E_2) =$$

$$P(E_1 \cap E_2) = P(\{(3,5)\}) = \frac{1}{36}$$

$$P(E_1 \cap E_2) = P(\{(3,5)\}) = \frac{1}{36}$$

▶ 
$$P(E_1) =$$

$$P(E_1 \cap E_2) = P(\{(3,5)\}) = \frac{1}{36}$$

$$P(E_1) = P(\{(3,1),(3,2),(3,3),(3,4),(3,5),(3,6)\}) = \frac{6}{36}$$

$$P(E_1 \cap E_2) = P(\{(3,5)\}) = \frac{1}{36}$$

$$P(E_1) = P(\{(3,1),(3,2),(3,3),(3,4),(3,5),(3,6)\}) = \frac{6}{36}$$

$$ightharpoonup P(E_2) =$$

$$P(E_1 \cap E_2) = P(\{(3,5)\}) = \frac{1}{36}$$

- $P(E_1) = P(\{(3,1),(3,2),(3,3),(3,4),(3,5),(3,6)\}) = \frac{6}{36}$
- $P(E_2) = P(\{(2,6),(3,5),(4,4),(5,3),(6,2)\}) = \frac{5}{36}$

$$P(E_1 \cap E_2) = P(\{(3,5)\}) = \frac{1}{36}$$

- $P(E_1) = P(\{(3,1),(3,2),(3,3),(3,4),(3,5),(3,6)\}) = \frac{6}{36}$
- $P(E_2) = P(\{(2,6),(3,5),(4,4),(5,3),(6,2)\}) = \frac{5}{36}$
- Since  $\frac{1}{36} \neq \frac{6}{36} \times \frac{5}{36}$  we see that  $P(E_1 \cap E_2) \neq P(E_1) \times P(E_2)$ , so events

$$P(E_1 \cap E_2) = P(\{(3,5)\}) = \frac{1}{36}$$

- $P(E_1) = P(\{(3,1),(3,2),(3,3),(3,4),(3,5),(3,6)\}) = \frac{6}{36}$
- $P(E_2) = P(\{(2,6),(3,5),(4,4),(5,3),(6,2)\}) = \frac{5}{36}$
- ▶ Since  $\frac{1}{36} \neq \frac{6}{36} \times \frac{5}{36}$  we see that  $P(E_1 \cap E_2) \neq P(E_1) \times P(E_2)$ , so events  $E_1$  and  $E_2$  are not independent.

$$P(E_1 \cap E_3) =$$

$$P(E_1 \cap E_3) = P(\{(3,4)\}) = \frac{1}{36}$$

$$P(E_1 \cap E_3) = P(\{(3,4)\}) = \frac{1}{36}$$

$$P(E_1) = P(\{(3,1),(3,2),(3,3),(3,4),(3,5),(3,6)\}) =$$

$$P(E_1 \cap E_3) = P(\{(3,4)\}) = \frac{1}{36}$$

$$P(E_1) = P(\{(3,1),(3,2),(3,3),(3,4),(3,5),(3,6)\}) = \frac{6}{36}$$

$$P(E_1 \cap E_3) = P(\{(3,4)\}) = \frac{1}{36}$$

$$P(E_1) = P(\{(3,1),(3,2),(3,3),(3,4),(3,5),(3,6)\}) = \frac{6}{36}$$

$$ightharpoonup P(E_3) =$$

$$P(E_1 \cap E_3) = P(\{(3,4)\}) = \frac{1}{36}$$

$$P(E_1) = P(\{(3,1),(3,2),(3,3),(3,4),(3,5),(3,6)\}) = \frac{6}{36}$$

$$P(E_3) = P(\{(1,6),(2,5),(3,4),(4,3),(5,2),(6,1)\}) = \frac{6}{36}$$

$$P(E_1 \cap E_3) = P(\{(3,4)\}) = \frac{1}{36}$$

- $P(E_1) = P(\{(3,1),(3,2),(3,3),(3,4),(3,5),(3,6)\}) = \frac{6}{36}$
- $P(E_3) = P(\{(1,6),(2,5),(3,4),(4,3),(5,2),(6,1)\}) = \frac{6}{36}$
- ▶ Since  $\frac{1}{36} = \frac{6}{36} \times \frac{6}{36}$  we see that  $P(E_1 \cap E_3) = P(E_1) \times P(E_3)$ , so


Rolling a dice

# Are $E_1$ and $E_3$ independent?-solution


$$P(E_1 \cap E_3) = P(\{(3,4)\}) = \frac{1}{36}$$

- $P(E_1) = P(\{(3,1),(3,2),(3,3),(3,4),(3,5),(3,6)\}) = \frac{6}{36}$
- $P(E_3) = P(\{(1,6),(2,5),(3,4),(4,3),(5,2),(6,1)\}) = \frac{6}{36}$
- Since  $\frac{1}{36} = \frac{6}{36} \times \frac{6}{36}$  we see that  $P(E_1 \cap E_3) = P(E_1) \times P(E_3)$ , so events  $E_1$  and  $E_3$  are independent.

#### Example: deck of cards




#### Example: deck of cards



- ► Define the following events
  - $\triangleright$   $E_1$ : A face card is selected.
  - $\triangleright$   $E_2$ : A king is selected.
  - $\triangleright$   $E_3$ : A heart is selected.


└ Deck of cards

### Example: deck of cards



- Define the following events
  - $\triangleright$   $E_1$ : A face card is selected.
  - $\triangleright$   $E_2$ : A king is selected.
  - $\triangleright$   $E_3$ : A heart is selected.
- ightharpoonup Are  $E_1$  and  $E_2$  independent?

#### Example: deck of cards



- Define the following events
  - ► *E*<sub>1</sub>: A face card is selected.
  - ► *E*<sub>2</sub>: A king is selected.
  - $\triangleright$   $E_3$ : A heart is selected.
- $\triangleright$  Are  $E_1$  and  $E_2$  independent?
- $\triangleright$  Are  $E_2$  and  $E_3$  independent?

$$P(E_1 \cap E_2) = P(\{KH, KC, KS, KD\}) = \frac{4}{52}$$

$$P(E_1 \cap E_2) = P(\{KH, KC, KS, KD\}) = \frac{4}{52}$$

► 
$$P(E_1) = P(\{JH, JC, JS, JD, KH, KC, KS, KD, QH, QC, QS, QD\}) = \frac{12}{52}$$

$$P(E_1 \cap E_2) = P(\{KH, KC, KS, KD\}) = \frac{4}{52}$$

- ►  $P(E_1) = P(\{JH, JC, JS, JD, KH, KC, KS, KD, QH, QC, QS, QD\}) = \frac{12}{52}$
- ►  $P(E_2) = P(\{KH, KC, KS, KD\}) = \frac{4}{52}$

$$P(E_1 \cap E_2) = P(\{KH, KC, KS, KD\}) = \frac{4}{52}$$

- ►  $P(E_1) = P(\{JH, JC, JS, JD, KH, KC, KS, KD, QH, QC, QS, QD\}) = \frac{12}{52}$
- $P(E_2) = P(\{KH, KC, KS, KD\}) = \frac{4}{52}$
- Since  $\frac{4}{52} \neq \frac{12}{52} \times \frac{4}{52}$  we see that  $P(E_1 \cap E_2) \neq P(E_1) \times P(E_2)$ , so events  $E_1$  and  $E_2$  are not independent.

▶  $E_2 \cap E_3$  is the event that a king and a heart is selected which is the event a kingheart is selected.

$$P(E_1 \cap E_2) = P(\{KH\}) = \frac{1}{52}$$

► 
$$P(E_2) = P(\{KH, KC, KS, KD\}) = \frac{4}{52}$$

└ Deck of cards

# Are $E_2$ and $E_3$ independent?-solution

▶  $E_2 \cap E_3$  is the event that a king and a heart is selected which is the event a kingheart is selected.

$$P(E_1 \cap E_2) = P(\{KH\}) = \frac{1}{52}$$

- ►  $P(E_2) = P(\{KH, KC, KS, KD\}) = \frac{4}{52}$
- ►  $P(E_3) = P({AH, 2H, 3H, 4H, 5H, 6H, 7H, 8H, 9H, 10H, JH, KH, QH}) = \frac{13}{52}$

▶  $E_2 \cap E_3$  is the event that a king and a heart is selected which is the event a kingheart is selected.

$$P(E_1 \cap E_2) = P(\{KH\}) = \frac{1}{52}$$

- ►  $P(E_2) = P(\{KH, KC, KS, KD\}) = \frac{4}{52}$
- ►  $P(E_3) = P({AH, 2H, 3H, 4H, 5H, 6H, 7H, 8H, 9H, 10H, JH, KH, QH}) = \frac{13}{52}$
- ▶ Since  $\frac{1}{52} = \frac{4}{52} \times \frac{13}{52}$  we see that  $P(E_2 \cap E_3) = P(E_2) \times P(E_3)$ , so events  $E_2$  and  $E_3$  are independent.

# Section summary

Examples of independent and dependent events.