

## IIT Madras ONLINE DEGREE

## Mathematics for Data Science 1 Week-03 Tutorial-04

(Refer Slide Time: 0:15)

Find the equation of a line which is perpendicular to line 
$$y - 5x = 0$$
 and if  $\sqrt{38}$  mits

$$y = 5 \times M_1 = 5$$

$$M_2 = -\frac{1}{2} \times M_1 = 5$$

$$y = -\frac{1}{2} \times K_1 = \frac{1}{2}$$

$$\frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

$$\frac{1}{2} \times K_1 + \frac{1}{2} \times K_2 = \frac{1}{2}$$

For our fourth question, we want the equation of a line which is perpendicular to this line, and is at this distance from the origin. So, from y-5x=0, we get y=5x, so therefore, the slope  $m_1$  is 5. And if our line is perpendicular to it, then our line  $m_2$  must be  $-1/m_1$ , which is equal to -1/5. So, we know that our line is some  $y=-\frac{x}{5}+C$ .

If we kind of simplify it, we are going to get 5y+x=C, this C is not the same thing as the previous C, I have just used that as C because it is an arbitrary constant, which is yet to be determined, otherwise it should have been 5C. Anyway, now we have to find the value of this C in this equation. For that, we are going to use the next bit of information that is given to us, which is the distance from the origin.

So, this line has this distance from the origin. So the distance from a point formula is  $\frac{|ax_1 + by_1 + c|}{\sqrt{a^2 + b^2}}$ , where  $(x_1, y_1)$  is the point from which we are measuring the distance for this

line. So, in our case,  $(x_1, y_1)$  is (0, 0) because we are doing from the origin. So in our case, we get modulus of  $\frac{|-c|}{\sqrt{a^2+b^2}}$ , So, modulus of |-c| is just the same thing as |c|.

And root of  $\sqrt{a^2+b^2}$ , in our case comes out to be  $\sqrt{25+1}$ ,, that is  $\sqrt{26}$ . So we have  $\frac{|c|}{\sqrt{26}}$ , this is given out to be  $\frac{1}{\sqrt{26}}$ , which would imply |c|=1, and that would imply  $c\pm 1$ . So, we get two answers.


(Refer Slide Time: 2:49)

IIT Madras

$$\frac{\left| a_{y_1} + b_{y_1} + c \right|}{\left| a_{y_2} + b_{y_3} \right|} = \frac{\left| -c \right|}{\left| a_{y_3} + b_{y_4} \right|} = \frac{\left| -c \right|}{\left| a_{y_4} + b_{y_5} + b_{y_5} \right|} = \frac{\left| -c \right|}{\left| a_{y_4} + b_{y_5} + b$$

What are the two answers? One is for c being +1, we have 5y + x = 1. And in the other case, we get 5y + x = -1 for the other choice. So, how does this happen, what is actually happening here to try to plot our lines?

(Refer Slide Time: 3:08)



So, we have two lines, which we were looking at, which is 5y + x = 1 and 5y + x = -1 and from this we get the intercepts to be, the intercept form of this would be  $\frac{y}{1/5} + \frac{x}{1} = 1$ . And in this case, we get  $\frac{y}{-1/5} + \frac{x}{-1} = 1$ . So, in one case, we have a y intercept of 1/5. So, let us assume this is 1/5, then x intercept is 1 which is 5 times of that, so that so it must be somewhere here, so this would be (1, 0) and this is (0, 1/5).

And our line is going through these two points, giving us something like this. Let us call this  $l_1$  and where do we get the  $\frac{1}{\sqrt{26}}$  distance from the origin, we get it when we measure it perpendicularly from the origin. Now, let us look at the other equation. So -1/5, so, this should be exactly below this this way and this is -1, so this would be exactly opposite in this way at the same distance.

So now we have these two points, so we can also construct this line, which goes this way. And as you can see, they are both parallel and exactly opposite to that  $\frac{1}{\sqrt{26}}$  you get this distance which is again perpendicular distance and it is also at  $\frac{1}{\sqrt{26}}$ . So, we have two lines which satisfy our requirements, one is 5y+x=1, the other is 5y+x=-1.