

RIVERSIDE GENERAL HOSPITAL
DISCHARGE INSTRUCTIONS

Patient: Margaret Ellison Age: 71 MRN: 000-TEST-0001
Admitted: 03/02 Discharged: 03/07
Attending: Dr. R. Okafor, Cardiology

REASON FOR HOSPITAL STAY

You were admitted because of worsening shortness of breath and leg swelling caused by congestive heart failure (your heart was not pumping as well as it should).
Fluid was removed with IV medicine and your breathing improved.

YOUR MEDICINES

NEW medicines

1. Furosemide (Lasix) 40 mg tablet - take 1 tablet by mouth every morning. This is a water pill to remove extra fluid. You may urinate more often. Take it early in the day.
2. Metoprolol succinate ER 25 mg tablet - take 1 tablet by mouth once daily. Do not crush or chew. Do not stop suddenly.

CHANGED medicines

3. Lisinopril: DECREASE from 20 mg to 10 mg once daily. Your kidney blood test was a little high, so your dose was lowered.

CONTINUE these home medicines

4. Atorvastatin 40 mg - 1 tablet at bedtime.
5. Aspirin 81 mg - 1 tablet daily with food.

STOP taking

6. Ibuprofen (Advil, Motrin) and naproxen (Aleve). These can make heart failure worse and hurt your kidneys. For pain you may use acetaminophen (Tylenol), no more than 3,000 mg in one day.

FOLLOW-UP

- Cardiology clinic (Dr. Okafor): within 7 days. Call 555-0142 to schedule.
- Blood test (kidney function and potassium): in 1 week, before your clinic visit.
- Primary care (Dr. Hale): within 2 weeks.

DAILY CARE AT HOME

- Weigh yourself every morning after using the bathroom and before eating. Write it down.
- Limit sodium (salt) to 2,000 mg per day. Do not add salt to food.
- Limit fluids to 1.5 liters (about 6 cups) per day.
- Walk for short periods as you feel able. Rest if you get short of breath.

CALL YOUR DOCTOR if you:

- Gain more than 3 pounds in 1 day or 5 pounds in 1 week
- Have more swelling in your feet, ankles or belly
- Feel dizzy or lightheaded, or have a fast heartbeat
- Need more pillows to sleep or wake up short of breath

CALL 911 if you have:

- Chest pain or pressure that does not go away
- Severe trouble breathing, fainting, or coughing up pink, foamy mucus