

SUMMIT ORTHOPEDIC HOSPITAL

Discharge Instructions - Total Hip Replacement (left)

Patient: Helen Park      Surgery date: 06/12      Going home: 06/14      Ref: TEST-3391

BLOOD CLOT PREVENTION

Apixaban (Eliquis) 2.5 mg - 1 tablet by mouth twice daily for 35 days after surgery.

This is to prevent blood clots. Do not skip doses. Tell any dentist or doctor that you take it.

PAIN CONTROL

- Acetaminophen 1,000 mg every 8 hours on a schedule for the first 2 weeks, then as needed.

Do not take more than 3,000 mg in 24 hours.

- Oxycodone 5 mg: 1 tablet every 6 hours ONLY IF pain is not controlled by acetaminophen.

This may cause drowsiness and constipation. No driving, no alcohol while taking it.

Number of tablets: 20. No refills.

- Celecoxib was NOT prescribed. Avoid ibuprofen and naproxen while on apixaban.

BOWEL CARE

- Docusate 100 mg twice daily while taking oxycodone. Add senna 8.6 mg at bedtime if no bowel movement in 2 days.

WOUND CARE

Keep the dressing clean and dry. You may shower on 06/20 (7 days); no baths, pools or hot tubs for 6 weeks. Do not put creams or ointments on the incision. Stitches/staples are removed at the 2-week visit.

ACTIVITY (hip precautions for 6 weeks)

- Do NOT bend your hip past 90 degrees (no low chairs or toilets without a raised seat).

- Do NOT cross your legs or ankles.

- Do NOT twist your body toward the operated leg.

- Walk with the walker as taught by physical therapy; short walks 4-5 times a day.

- No driving until cleared at your follow-up visit.

FOLLOW-UP

Surgeon Dr. Moreno, 2 weeks after surgery (06/26). Home physical therapy begins within 2 days after you get home; the agency will call you.

CALL THE OFFICE (555-0188) if: temperature over 101.5 F, redness, warmth or drainage at the incision, increasing pain, or calf pain/swelling in either leg.

CALL 911 if: sudden shortness of breath, chest pain, coughing blood, signs of stroke, bleeding that will not stop, or if you fall and cannot get up or your leg looks shortened or turned.