

HARBOR VIEW SURGICAL CENTER

Discharge Instructions - Laparoscopic Gallbladder Removal

Patient: Robert Chen Age: 64 Ref: TEST-5120
Surgery date: 09/22 Going home: 09/23
Surgeon: Dr. L. Ibarra

PAIN AND NAUSEA MEDICINES (new)

1. Tylenol (acetaminophen) 500 mg - take 2 tablets by mouth every 6 hours for pain.
2. Percocet (oxycodone/acetaminophen) 5/325 mg - take 1 tablet by mouth every 6 hours
ONLY IF your pain is severe. It can cause drowsiness. Do not drive.
3. Motrin IB (ibuprofen) 200 mg - take 1 tablet every 8 hours with food if you have swelling.
4. Ondansetron 4 mg - take 1 tablet every 8 hours only if you feel sick to your stomach.
5. Docusate 100 mg - take 1 capsule twice daily with meals to keep your bowels moving.

HOME MEDICINES

- Warfarin 5 mg: continue 1 tablet at bedtime. Your clinic will check your INR blood test.
- Lisinopril 10 mg: continue 1 tablet every morning.

STOP taking

- Advil (ibuprofen) and aspirin until your surgeon says it is safe.

FOLLOW-UP

- Surgeon, Dr. Ibarra: 2 weeks after surgery.
- INR blood test: in 3 days.

CARE AT HOME

- You may shower after 48 hours. Do not soak in a bath for 2 weeks.
- No lifting more than 10 pounds for 4 weeks.

CALL 911 if: chest pain, trouble breathing, or vomiting blood.

CALL THE OFFICE if: fever over 101 F, yellow skin or eyes, severe belly pain, or bleeding from the wounds.