

Clinical Epigenetics

The differential expression of mRNAs and long noncoding RNAs between ectopic and eutopic endometrium provides new insights into adenomyosis

--Manuscript Draft--

Manuscript Number:	CLEP-D-15-00065
Full Title:	The differential expression of mRNAs and long noncoding RNAs between ectopic and eutopic endometrium provides new insights into adenomyosis
Article Type:	Research
Abstract:	<p>Abstract</p> <p>Background: Adenomyosis, defined as ectopic endometrial tissue within the myometrium, can often be misdiagnosed as multiple uterine leiomyomata or endometrial thickening. We therefore performed a combined mRNA and long noncoding (lnc)RNA microarray and bioinformatic analysis of eutopic and ectopic endometrium in women with adenomyosis to better understand its pathogenesis and help in the development of a semi-invasive diagnostic test.</p> <p>Results: A total of 586 mRNAs were increased and 305 mRNAs decreased in ectopic endometrium of adenomyosis compared with eutopic endometrium, while 388 lncRNA transcripts were up-regulated and 188 down-regulated in ectopic compared with paired eutopic endometrial tissue.</p> <p>Conclusion: mRNA and lncRNA expression was comparable in eutopic and ectopic endometria. Bioinformatic analysis suggested a series of metabolic and molecular abnormalities in adenomyosis, which have many similarities with endometriosis. Furthermore, our study constitutes the first known report of lncRNA expression patterns in human adenomyosis ectopic and eutopic endometrial tissue.</p>

**The differential expression of mRNAs and long noncoding
RNAs between ectopic and eutopic endometrium provides
new insights into adenomyosis**

Cheng Zhou^{a,#}, Ting Zhang^{b,#}, Ji Zhou^a, Xiaobei Ni^a, Ran Huo^{a*}, Zhonghua
Shi^{a,c*}

^aState Key Laboratory of Reproductive Medicine, Department of Histology and
Embryology, Nanjing Medical University, Nanjing, China

^bWuxi Maternity and Child Health Hospital, Wuxi city ,China

^cState Key Laboratory of Reproductive Medicine, Nanjing Maternity and Child
Health Hospital, Nanjing Medical University, Nanjing, China

[#]These authors contributed equally to this work.

*Correspondence:

Ran Huo (Email: huoran@njmu.edu.cn) or Zhonghua Shi (Email:
jesse_1982@163.com)

Other authors' email addresses:

Cheng Zhou:510417094@qq.com; Ting Zhang:zhangting040715@163.com;

Ji Zhou: cherryzhouji@163.com; Xiaobei Ni: nixiaobei@163.com

Abstract

Background: Adenomyosis, defined as ectopic endometrial tissue within the myometrium, can often be misdiagnosed as multiple uterine leiomyomata or endometrial thickening. We therefore performed a combined mRNA and long noncoding (lnc)RNA microarray and bioinformatic analysis of eutopic and ectopic endometrium in women with adenomyosis to better understand its pathogenesis and help in the development of a semi-invasive diagnostic test.

Results: A total of 586 mRNAs were increased and 305 mRNAs decreased in ectopic endometrium of adenomyosis compared with eutopic endometrium, while 388 lncRNA transcripts were up-regulated and 188 down-regulated in ectopic compared with paired eutopic endometrial tissue.

Conclusion: mRNA and lncRNA expression was comparable in eutopic and ectopic endometria. Bioinformatic analysis suggested a series of metabolic and molecular abnormalities in adenomyosis, which have many similarities with endometriosis. Furthermore, our study constitutes the first known report of lncRNA expression patterns in human adenomyosis ectopic and eutopic endometrial tissue.

Keywords:

Adenomyosis, microarray, long noncoding RNA, lymphangiogenesis

Background

Adenomyosis' characteristics is the presence of ectopic endometrial mucosa

1 within the myometrium and the invagination of endometrium at a depth of at
2
3 least 2.5 mm below the basal layer of the endometrium. It is a common benign
4
5 pathology that is asymptomatic in one third of cases. Clinical signs, consisting
6
7 mainly of menometrorrhagia, dysmenorrhoea, and pelvic pain, are non-specific,
8
9 often leading to the misdiagnosis of multiple uterine leiomyomata or
10
11 endometrial thickening, both of which have different prognoses and
12
13 treatments[1].
14
15
16
17
18
19
20
21

22 The pathogenesis of adenomyosis is poorly defined. Its diagnosis is confirmed
23
24 by hypertrophy of smooth muscle caused by the invagination of endometrium
25
26 [2], and it is often associated with hormone-dependent pelvic lesions such as
27
28 myoma, deep pelvic endometriosis, polyps, or endometrial hyperplasia[3, 4].
29
30
31
32
33
34
35

36 Endometriosis is defined as the presence of endometrial glands and stroma in
37
38 ectopic locations, mainly the pelvic peritoneum, ovaries, and rectovaginal
39
40 septum. The clinical signs of endometriosis include dysmenorrhea,
41
42 dyspareunia, chronic pelvic pain, irregular uterine bleeding, and/or infertility[5].
43
44
45
46

47 A number of similarities exist between adenomyosis and endometriosis, both
48
49 of which were originally described as adenomyoma. The two conditions have
50
51 been recognized as distinct entities since the retrograde menstruation theory
52
53 became widely accepted to explain the pathogenesis of endometriosis,
54
55 because it does not explain adenomyosis[6]. However, magnetic resonance
56
57
58
59
60
61
62
63
64
65

1 imaging and laparoscopy showed that the two conditions often co-exist, at
2
3 least in some subgroups[7].One study reported that 27% of women with
4
5 endometriosis had concomitant adenomyosis[8]. Moreover, adenomyosis is
6
7 thought to be a significant factor of infertility in endometriosis, probably
8
9 because of changes in the uterine transport of spermatozoa[9].
10
11
12
13
14
15
16

17 Investigations into the pathophysiology of endometriosis have revealed
18
19 several well-sustained molecular hallmarks of this disease, such as genetic
20
21 predisposition, estrogen dependence, progesterone resistance, and
22
23 inflammation[10]. At the same time, eutopic and ectopic endometria of
24
25 adenomyosis were reported to show a series of metabolic and molecular
26
27 abnormalities which include increase angiogenesis and proliferation, decrease
28
29 apoptosis, allow local production of estrogens, create progesterone resistance,
30
31 and impair cytokine expression[11]. However, the mechanism responsible for
32
33 the survival and subsequent implantation of the translocated endometrium
34
35 remains unknown. Moreover, the origin of adenomyosis and endometriosis is
36
37 still controversial, which not only influencing our understanding of uterine
38
39 function and pathophysiology, but also of clinical management and treatment.
40
41 Long noncoding RNAs (lncRNAs) are a class of ncRNAs up to 200 nucleotides
42
43 in length. Through observations of the transcriptional activation ripple effect,
44
45 they were long disregarded as transcriptional noise because of their low
46
47 expression levels[12]. However, recent accumulating evidence suggests that
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 lncRNAs function in biological processes through diverse mechanisms[13],
2
3 and many studies have reported their roles in genetic imprinting, immune
4
5 responses, cell differentiation, human diseases, and other biological
6
7 processes[14-16]. It has also been suggested that their expression can be cell
8
9 specific and regulated by common transcription factors[17, 18].
10
11
12
13
14
15
16

17 This study was performed to establish the differential mRNA and lncRNA
18
19 expression in the eutopic and ectopic endometria of adenomyosis, with the aim
20
21 of furthering our understanding of the pathogenesis of adenomyosis and
22
23 developing a semi-invasive diagnostic test. We collected eutopic and ectopic
24
25 endometria from women with adenomyosis and conducted microarray and
26
27 bioinformatic analyses.
28
29
30
31
32

33 **Results**

34 **HE staining of surgical specimens**

35
36 Normal endometria contained a few uterine glands, numerous matrix cells,
37
38 reticular fibers, and blood vessels. Adenomyosis was diagnosed when ectopic
39
40 endometrial mucosa were observed within the myometrium. See Figure 1.
41
42
43
44
45
46

47 **Differentially expressed mRNAs and lncRNAs**

48
49 A total of 891 mRNAs (see Additional file 1:Table S1) and 576 lncRNAs (see
50
51 Additional file 2:Table S2) were identified by microarray analysis after quantile
52
53 normalization and data filtering steps. Of the mRNAs, 586 were up-regulated
54
55 and 305 were down-regulated (fold-change >2, $P < 0.05$) in ectopic compared
56
57
58
59
60
61
62
63
64
65

1 with eutopic endometria, respectively. Of the lncRNAs, 388 were up-regulated,
2
3 and 188 were down-regulated (fold-change >2, $P < 0.05$), respectively.
4
5

6 **Quantitative real-time PCR validation**

7
8 The expression levels of three randomly chosen differentially expressed
9 mRNAs (*NRP2*, *TGFBR1*, and *ITGB3*) and three lncRNAs (uc004dwe.2,
10
11 ENST00000454594, and NR_003521) were verified by quantitative
12
13 (q)RT-PCR. The relative fold-changes in expression, as detected by qPCR,
14
15
16
17
18
19
20 were consistent with the microarray data. See Figure 2.
21
22

23 **Gene Ontology (GO) and pathway analysis**

24
25 Genes showing greater than a 2-fold change in expression level were
26
27 analyzed by GO. Those genes up-regulated in ectopic compared with eutopic
28
29 endometrial samples were found to be involved in a variety of pathways
30
31 including vasculogenesis, cell–matrix adhesion, cell differentiation, and the
32
33 regulation of cell proliferation, apoptosis, and muscle contraction. By contrast,
34
35
36
37
38 those genes down-regulated in ectopic compared with eutopic endometrial
39
40
41
42 samples were involved in immune response pathways, and the positive
43
44
45 regulation of the I-kappaB kinase/NF-kappaB cascade. See Figure 3a,b.
46
47
48
49

50 The major pathways thought to involve mRNAs up-regulated in ectopic versus
51
52 eutopic endometria include the adhesion junction, MAPK signaling, and
53
54
55 cytokine–cytokine receptor interaction, while mRNAs down-regulated in
56
57
58
59 ectopic versus eutopic endometria are thought to be involved in pathways of
60
61
62
63
64
65

androgen and estrogen metabolism, among others. See Figure 3c,d.

Signal network analysis

Ingenuity pathway analysis was used to identify pathways of significantly differentially expressed genes. A network of genes was built according to the relationships among the genes, proteins, and compounds of the KEGG database[19, 20]. This identified many candidate genes and their interactions that may play an important role in the pathogenesis and development of adenomyosis. See Figure 3e.

lncRNA and mRNA coexpression profiles and lncRNA function prediction

To associate lncRNAs with the regulated expression of target mRNAs, we superimposed lncRNA target predictions onto the lncRNA–mRNA correlation network. The resulting network was defined as an lncRNA–mRNA regulatory network, with direct connections between an lncRNA and an mRNA represented by solid lines. A total of 24 lncRNAs were co-expressed with many different mRNAs in the co-expression profiles. See Figure 4. This enabled us to predict the functions of lncRNAs through the GO and KEGG pathway annotations of their co-expressed mRNAs.

Discussion

The establishment of adenomyosis requires several key steps, including epithelial–mesenchymal transition (EMT), survival and growth of ectopic tissue

1 in the myometrium, myometrial hypertrophy and hyperplasia, and induction of
2
3 an immunosuppressive microenvironment[21]. We used microarray analysis to
4
5 show that 891 mRNA and 576 lncRNA transcripts are dysregulated in the
6
7 ectopic endometrium of adenomyosis compared with paired eutopic
8
9 endometrium. Our GO analysis data show that the different genes between
10
11 eutopic and ectopic endometrium of adenomyosis are involved in pathways of
12
13 cell–matrix adhesion, cell differentiation, vasculogenesis, the positive
14
15 regulation of cell proliferation and the I-kappaB kinase/NF- kappaB cascade,
16
17 and the regulation of apoptosis, immune responses, EMT, and muscle
18
19 contraction, it's worth noting that,all of above have previously been reported to
20
21 participate in the pathogenesis and pathophysiology of endometriosis[22-24]. It
22
23 has been hypothesized that adenomyosis and endometriosis share a common
24
25 pathogenesis[25-27], which is supported by our data, although more research
26
27 is needed to confirm this hypothesis.
28
29
30
31
32
33
34
35
36
37
38
39
40
41

42 Our findings support those of a number of other studies about endometriosis
43
44 and adenomyosis. For example, Mechsner reported that epithelial and stromal
45
46 cells in endometriosis develop from persistent coelomic epithelial cells by
47
48 metaplasia [28], which is in line with our GO analysis including of cell
49
50 differentiation and the positive regulation of cell proliferation . Similarly, our
51
52 results support the theory that persistent estrogenic stimulation leads to
53
54 EMT[29], myometrial smooth muscle cell hypertrophy, and hyperplasia of
55
56
57
58
59
60
61
62
63
64
65

adenomyosis[30]. A previous study suggested that uterine myometrial growth can be modulated by prolonged mechanical stretch and contraction increasing the intrauterine pressure and leading to smooth muscle cell hypertrophy and hyperplasia of adenomyosis [31]. And, the most widely accepted hypothesis is that adenomyosis is caused by defects of the inner myometrial layer of the uterus[32], and that invagination of the stroma together with microenvironmental factors stimulate smooth muscle cell growth[21].

Indeed, many of the genes shown to be differentially expressed by our microarray data have been reported to be associated with the development of endometriosis. These include *CDH1*, *SGPP2*, *MET*, and *SLC2A1*, which demonstrate 2.5- to 4.7-fold reductions in the ectopic versus eutopic endometria of women with adenomyosis in our study, with the observed reduction in *CDH1* and *SGPP2* expression being consistent with the literature[33, 34]. Moreover, the expression of *VEGFA*, *PDGFA*, *KDR*, *ACTN4*, *CD44*, and *JAM2* was increased 2.3- to 3.2-fold in ectopic versus eutopic endometria of women with adenomyosis, with the up-regulation of *VEGFA*, *PDGFA*, *KDR*, and *ACTN4* being in line with earlier reports[35-38].

To the best of our knowledge, this study presents the first mRNA and lncRNA microarray analysis of eutopic and ectopic endometria in women with adenomyosis. Our findings show that mRNA expression differs between the

two endometrial types, with ectopic endometria more closely associated with pathways of focal adhesion, tight junctions, regulation of the actin cytoskeleton, the calcium signaling pathway, the transforming growth factor-beta signaling pathway, androgen and estrogen metabolism, the cell cycle, EMT, positive regulation of mesenchymal cell proliferation, and regulation of muscle contraction, all of which are related to the development of adenomyosis[21].

Although our GO and pathway analysis reveal many similarities between the development of adenomyosis and endometriosis, adenomyosis may also have some unique molecular mechanisms, and some differentially expressed lncRNAs may play a role in its pathogenesis and development by affecting their coexpressed mRNAs. For example, the most widely accepted theory of retrograde menstruation of endometriosis isn't suitable to explain adenomyosis. The detection of endometrial stromal cells within lymph nodes in a baboon model of induced endometriosis [39] and in 6–7% of women at lymphadenectomy [40] supports the possibility of the lymphatic dissemination of endometrial cells.

It's a great encouraging detection of the high over-expression of the neuropilin 2 gene (*NRP2*) in ectopic versus eutopic endometria which was verified by our qRT-PCR. This gene demonstrates tissue-specific distribution in glandular and surface epithelia, venous and lymphatic endothelia, and a leukocyte

subpopulation, with low levels reported in endometrial stromal cells[41].

NRP receptors and tyrosine kinase receptors such as vascular endothelial cell growth factor (VEGFR) are cell surface receptor families of the VEGF family that are important mediators of angiogenesis and lymphangiogenesis[42]. Our GO analysis also confirmed that the differentially expressed genes are more related to many pathways including vasculogenesis. NRP2 is important in modulating developmental lymphangiogenesis and mediating VEGF-C-induced lymphatic sprouting[43]. The previous observation of endometrial tissue in intramyometrial lymphatics [44] together with our microarray findings of NRP2' high over-expression in ectopic endometrium indicate that lymphatic vessels are a possible route for the invagination or dissemination of the basal endometrium in adenomyosis.

Our present study findings suggest that 24 lncRNAs are involved in the establishment of adenomyosis. Although the functions of many lncRNAs are poorly understood, it is of interest that our lncRNA microarray analysis showed that lncRNA uc004dwe.2 is over-expressed in ectopic compared with eutopic endometria. This is in line with our real-time PCR result and is highly correlated with the expression of NRP2, suggesting that lncRNA uc004dwe.2 may play a role in the pathophysiology of adenomyosis through modulating the function of NRP2.

Conclusions

In conclusion, our study is the first to investigate genome-wide lncRNA expression patterns in human adenomyosis tissue, and identified several mRNAs and lncRNAs differentially expressed between paired, autologous, ectopic and eutopic endometria in women with adenomyosis. Several metabolic and molecular abnormalities detected in this study have previously been reported to participate in biological pathways related to endometriosis. The lncRNAs identified may participate in the pathophysiology of adenomyosis through diverse mechanisms. Further research is required in this area because of the limited current understanding of lncRNA functions in adenomyosis.

Methods

Patients and samples

Endometrial samples were collected from women of childbearing years undergoing hysterectomies for adenomyosis. Patients using the oral contraceptive pill (combined or progesterone only), those on chronic medication, or those having undergone surgery in the 6 months prior to sample collection were excluded. Specimens of ectopic endometria were isolated away from areas of obvious uterine adenomyoma. Eutopic endometria were full-thickness, 1-cm biopsies of the uterine wall. Harvested samples were divided into two parts: one half was snap-frozen in liquid nitrogen and prepared

for RNA extraction. The other half was fixed and used for morphological examination. Sectioning and hematoxylin and eosin staining of surgical specimens were carried out by the pathological department of the Wuxi Maternity and Child Health Hospital (Jiangsu Province, China).

All patients had signed a written informed consent prior to recruitment and the study protocol was approved by the Medical Ethics Committee of Wuxi Maternity and Child Health Hospital.

Microarray analysis

mRNA and lncRNA microarray analysis was undertaken in three paired samples of uterine adenomyoma versus matched eutopic endometrial tissues using the Agilent (California, USA) microarray platform. Sample preparation and microarray hybridization were performed according to the manufacturer's standard protocols with minor modifications. Briefly, total RNA was extracted using TRIzol reagent (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions and purified to remove rRNA using the NucleoSpin® RNA Clean-up kit (Macherey-Nagel, Cat # 740.948.250). Then, each sample was amplified and transcribed into fluorescent (Cy5 and Cy3-dCTP) labeled cRNA along the entire length of the transcripts without 3' bias using a reaction with random primers. Labeled cRNAs were hybridized onto the Agilent Human LncRNA+mRNA Array V2.0 (4×180K format), which contains probes interrogating about 39,000 human lncRNAs and about 32,000

human mRNAs assembled from databases such as RefSeq, Ensembl, the Chen Runsheng Laboratory (Institute of Biophysics, Chinese Academy of Science), and other related literature. The arrays were scanned on the Agilent Scanner G2565CA after washing the slides.

Agilent Feature Extraction software version 10.7 (Arraystar, USA) was used to analyze the acquired array images. Quantile normalization and subsequent data processing were performed using the GeneSpring GX v11.5.1 software package (Arraystar). Microarray data were selected using threshold values of >2 and <-2 fold-change under false discovery rate (FDR) protection ($P < .05$). Hierarchical clustering was performed to show distinguishable expression patterns of lncRNAs and mRNAs.

RNA labeling, and microarray hybridization and analysis were performed by CapitalBio (Beijing, China).

Isolation of total RNA and Quantitative Real-time PCR

Total RNA was isolated from ectopic and eutopic endometria tissues as described above. The RNA yield was determined using a NanoDrop 2000c spectrophotometer (Thermo Scientific, Rockford, IL, USA), and its integrity was evaluated using agarose gel electrophoresis stained with ethidium bromide. Reverse transcription was carried out in a GeneAmp PCR System 9700 (Applied Biosystems) for 15 min at 37°C, followed by heat inactivation for

5 s at 85°C.

Real-time PCR was performed using a StepOnePlus Real-Time PCR Instrument (Applied Biosystems) in a 20-μL reaction including 2-μL of template DNA (100 ng/μL), 10 μL 2xSYBR Green Master Mix (Vazyme), 0.8 μL forward primer (5 μM), 0.8 μL reverse primer (5 μM), 0.4 μL 50xROX Reference Dye 1, and 6 μL nuclease-free water. Reactions were incubated in a 96-well optical plate (Applied Biosystems) at 95°C for 6 min, followed by 40 cycles of 95°C for 30 s, 60°C for 30 s, and 72°C for 30 s. Each sample was run in triplicate.

Melting curve analysis was performed to validate the specific generation of the expected PCR product. Primer sequences were synthesized by Generay Biotech (Shanghai, China) on the basis of the noncoding RNA sequences obtained from the National Center for Biotechnology Information (NCBI) database. Primer sequences are listed in Table I. lncRNA and mRNA expression levels were normalized to those of glyceraldehyde 3-phosphate dehydrogenase and calculated using the $2^{-\Delta\Delta C_t}$ method[45].

Bioinformatics analysis

All mRNA and lncRNAs microarray data were given by Capital Bio Corp. (Beijing, China). Raw microarray signals underwent \log_2 transformation after being normalized using Agilent GeneSpring software. Differential expression of mRNA or lncRNA was defined by an absolute value of fold-change (FC) >2 (eutopic endometrium=1) and $P < 0.05$ (unpaired t -test).

Gene ontology (GO) analysis of the main function of the differentially expressed genes was conducted according to the GO database, which provides the key functional classifications for the NCBI [46]. Fisher's exact test and the χ^2 test were used to classify the GO category, and the FDR[47] was calculated to correct the *P*-value: the smaller the FDR, the smaller the error in judging the *P*-value. *P*-values were computed for the GOs of all differential genes. Enrichment provided a measure of the significance of the function: as the enrichment increased, the corresponding function became more specific, which helped identify those GOs with a more concrete function description in adenomyosis. Within the significant category, the enrichment *Re* was given by:

$$Re = (n_f / n) / (N_f / N),$$

where n_f is the number of differential genes within the particular category, n is the total number of genes within the same category, N_f is the number of differential genes in the entire microarray, and N is the total number of genes in the microarray [48].

Similarly, ingenuity pathway analysis was used to identify pathways involving significantly differentially expressed genes according to the Kyoto Encyclopedia of Genes and Genomes (KEGG). Fisher's exact test and the χ^2 test were used as statistical tests, and the threshold of significance was defined by the *P*-value and FDR. The enrichment *Re* was calculated using the equation mentioned in a previous study [49].

Co-expression network and functional prediction of lncRNAs

Because most lncRNAs in current databases have not yet been functionally annotated, the prediction of their functions relies on the functional annotations of their co-expressed mRNAs, i.e., neighboring protein-coding genes, which may influence each other during biogenesis [50]. We constructed co-expression networks of genes with specific expression levels according to their normalized signal intensities. The Pearson's correlation coefficient was calculated for each pair of genes, and significantly correlated pairs were chosen to construct the network.

Availability of supporting data

The microarray data sets supporting the results of this article are included within the article and its additional files.

Additional files

Additional file 1:Table S1.Fold-changes of differentially expressed mRNAs in ectopic endometria compared with eutopic endometria.

Additional file 2 :Table S2. Fold-changes of differentially expressed lncRNAs in ectopic endometria compared with eutopic endometria.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

CZ and RH carried out the mRNA and lncRNA expression analyses. ZS performed the statistical analysis of microarray data.CZ and JZ performed the statistics and drafted the manuscript. XN and TZ participated in the study design and helped to draft

the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We thank the Shanghai FudanBio Technology Co. for performing bioinformatic analysis.

References

1. Levy G, Dehaene A, Laurent N, Lernout M, Collinet P, Lucot JP, Lions C, Poncelet E. An update on adenomyosis. *Diagn Interv Imaging*. 2013; 94:3-25.
2. Siegler AM, Camilien L. Adenomyosis. *J Reprod Med*. 1994; 39:841-853.
3. Nisolle M, Donnez J. Peritoneal endometriosis, ovarian endometriosis, and adenomyotic nodules of the rectovaginal septum are three different entities. *Fertil Steril*. 1997; 68:585-596.
4. Koninckx PR, Martin DC. Deep endometriosis: a consequence of infiltration or retraction or possibly adenomyosis externa? *Fertil Steril*. 1992; 58:924-928.
5. Eskenazi B, Warner ML. Epidemiology of endometriosis. *Obstet Gynecol Clin North Am*. 1997; 24:235-258.
6. Corwin EJ. Endometriosis: pathophysiology, diagnosis, and treatment. *Nurse Pract*. 1997; 22:35-38, 40-32, 45-36, passim; quiz 56-37.
7. Benagiano G, Brosens I, Habiba M. Structural and molecular features of the endomyometrium in endometriosis and adenomyosis. *Hum Reprod Update*. 2014; 20:386-402.
8. Bazot M, Fiori O, Darai E. Adenomyosis in endometriosis--prevalence and impact on fertility. Evidence from magnetic resonance imaging. *Hum Reprod*. 2006;21:1101-1102; author reply 1102-1103.
9. Kunz G, Beil D, Huppert P, Noe M, Kissler S, Leyendecker G. Adenomyosis in endometriosis--prevalence and impact on fertility. Evidence from magnetic resonance imaging. *Hum Reprod*. 2005; 20:2309-2316.
10. Burney RO, Giudice LC. Pathogenesis and pathophysiology of endometriosis. *Fertil Steril*. 2012; 98:511-519.
11. Benagiano G, Brosens I. The endometrium in adenomyosis. *Womens Health (Lond Engl)*. 2012; 8:301-312.
12. Ebisuya M, Yamamoto T, Nakajima M, Nishida E. Ripples from neighbouring transcription. *Nat Cell Biol*. 2008; 10:1106-1113.
13. Wang KC, Chang HY. Molecular mechanisms of long noncoding RNAs. *Mol Cell*. 2011;43:904-914.
14. Taft RJ, Pang KC, Mercer TR, Dinger M, Mattick JS. Non-coding RNAs: regulators of disease. *J Pathol*. 2010; 220:126-139.
15. Tupy JL, Bailey AM, Dailey G, Evans-Holm M, Siebel CW, Misra S, Celniker SE, Rubin GM. Identification of putative noncoding polyadenylated transcripts in *Drosophila melanogaster*. *Proc Natl Acad Sci U S A*. 2005; 102:5495-5500.
16. Wilusz JE, Sunwoo H, Spector DL. Long noncoding RNAs: functional surprises from the RNA world. *Genes Dev*. 2009; 23:1494-1504.

17. Tsai MC, Manor O, Wan Y, Mosammaparast N, Wang JK, Lan F, Shi Y, Segal E, Chang HY. Long noncoding RNA as modular scaffold of histone modification complexes. *Science*. 2010; 329:689-693.
18. Mercer TR, Dinger ME, Mattick JS. Long non-coding RNAs: insights into functions. *Nat Rev Genet*. 2009; 10:155-159.
19. Jansen R, Lan N, Qian J, Gerstein M. Integration of genomic datasets to predict protein complexes in yeast. *J Struct Funct Genomics*. 2002; 2:71-81.
20. Wei Z, Li M. Genome-wide linkage and association analysis of rheumatoid arthritis in a Canadian population. *BMC Proc*. 2007; 1 Suppl 1:S19.
21. Koike N, Tsunemi T, Uekuri C, Akasaka J, Ito F, Shigemitsu A, Kobayashi H. Pathogenesis and malignant transformation of adenomyosis (review). *Oncol Rep*. 2013; 29:861-867.
22. Ponce C, Torres M, Galleguillos C, Sovino H, Boric MA, Fuentes A, Johnson MC. Nuclear factor kappaB pathway and interleukin-6 are affected in eutopic endometrium of women with endometriosis. *Reproduction*. 2009; 137:727-737.
23. Herington JL, Bruner-Tran KL, Lucas JA, Osteen KG. Immune interactions in endometriosis. *Expert Rev Clin Immunol*. 2011; 7:611-626.
24. Gonzalez-Ramos R, Defrere S, Devoto L. Nuclear factor-kappaB: a main regulator of inflammation and cell survival in endometriosis pathophysiology. *Fertil Steril*. 2012; 98:520-528.
25. Leyendecker G, Herbertz M, Kunz G, Mall G. Endometriosis results from the dislocation of basal endometrium. *Hum Reprod*. 2002; 17:2725-2736.
26. Leyendecker G, Kunz G, Noe M, Herbertz M, Mall G. Endometriosis: a dysfunction and disease of the archimetra. *Hum Reprod Update*. 1998; 4:752-762.
27. Leyendecker G, Kunz G, Wildt L, Beil D, Deininger H. Uterine hyperperistalsis and dysperistalsis as dysfunctions of the mechanism of rapid sperm transport in patients with endometriosis and infertility. *Hum Reprod*. 1996; 11:1542-1551.
28. Mechsner S, Bartley J, Infanger M, Loddenkemper C, Herbel J, Ebert AD. Clinical management and immunohistochemical analysis of umbilical endometriosis. *Arch Gynecol Obstet*. 2009; 280:235-242.
29. Chen YJ, Li HY, Huang CH, Twu NF, Yen MS, Wang PH, Chou TY, Liu YN, Chao KC, Yang MH. Oestrogen-induced epithelial-mesenchymal transition of endometrial epithelial cells contributes to the development of adenomyosis. *J Pathol*. 2010;222:261-270.
30. Jaffer S, Shynlova O, Lye S. Mammalian target of rapamycin is activated in association with myometrial proliferation during pregnancy. *Endocrinology*. 2009; 150:4672-4680.
31. Ferenczy A. Pathophysiology of adenomyosis. *Hum Reprod Update*. 1998; 4:312-322.
32. Parrott E, Butterworth M, Green A, White IN, Greaves P. Adenomyosis--a result of disordered stromal differentiation. *Am J Pathol*. 2001; 159:623-630.
33. Panda H, Pelakh L, Chuang TD, Luo X, Bukulmez O, Chegini N. Endometrial miR-200c is altered during transformation into cancerous states and targets the expression of ZEBs, VEGFA, FLT1, IKKbeta, KLF9, and FBLN5. *Reprod Sci*. 2012; 19:786-796.
34. Santulli P, Marcellin L, Noel JC, Borghese B, Fayt I, Vaiman D, Chapron C, Mehats C. Sphingosine pathway deregulation in endometriotic tissues. *Fertil Steril*. 2012;

- 97:904-911.
35. Takehara M, Ueda M, Yamashita Y, Terai Y, Hung YC, Ueki M. Vascular endothelial growth factor A and C gene expression in endometriosis. *Hum Pathol.* 2004; 35:1369-1375.
 36. Yu CQ, Yu J, Han J, Zhou QL, Shen W. [Regulatory mechanism of malignant behavior of endometriosis mediated by puerarin]. *Zhong Xi Yi Jie He Xue Bao.* 2009; 7:41-47.
 37. Wang HB, Lang JH, Leng JH, Zhu L, Liu ZF, Sun DW. [Expression of vascular endothelial growth factor receptors in the ectopic and eutopic endometrium of women with endometriosis]. *Zhonghua Yi Xue Za Zhi.* 2005; 85:1555-1559.
 38. Honda H, Barrueto FF, Gogusev J, Im DD, Morin PJ. Serial analysis of gene expression reveals differential expression between endometriosis and normal endometrium. Possible roles for AXL and SHC1 in the pathogenesis of endometriosis. *Reprod Biol Endocrinol.* 2008;6:59.
 39. Hey-Cunningham AJ, Fazleabas AT, Braundmeier AG, Markham R, Fraser IS, Berbic M. Endometrial stromal cells and immune cell populations within lymph nodes in a nonhuman primate model of endometriosis. *Reprod Sci.* 2011; 18:747-754.
 40. Javert CT. The spread of benign and malignant endometrium in the lymphatic system with a note on coexisting vascular involvement. *Am J Obstet Gynecol.* 1952;64:780-806.
 41. Hey-Cunningham AJ, Markham R, Fraser IS, Berbic M. Dysregulation of vascular endothelial growth factors and their neuropilin receptors in the eutopic endometrium of women with endometriosis. *Reprod Sci.* 2013; 20:1382-1389.
 42. Ferrara N, Gerber HP, LeCouter J. The biology of VEGF and its receptors. *Nat Med.* 2003; 9:669-676.
 43. Xu Y, Yuan L, Mak J, Pardanaud L, Caunt M, Kasman I, Larrivee B, Del Toro R, Suchting S, Medvinsky A et al. Neuropilin-2 mediates VEGF-C-induced lymphatic sprouting together with VEGFR3. *J Cell Biol.* 2010; 188:115-130.
 44. Sahin AA, Silva EG, Landon G, Ordonez NG, Gershenson DM. Endometrial tissue in myometrial vessels not associated with menstruation. *Int J Gynecol Pathol.* 1989; 8:139-146.
 45. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods.* 2001; 25:402-408.
 46. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT et al. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet.* 2000; 25:25-29.
 47. Dupuy D, Bertin N, Hidalgo CA, Venkatesan K, Tu D, Lee D, Rosenberg J, Svrikapa N, Blanc A, Carnec A et al. Genome-scale analysis of in vivo spatiotemporal promoter activity in *Caenorhabditis elegans*. *Nat Biotechnol.* 2007; 25:663-668.
 48. Schlitt T, Palin K, Rung J, Dietmann S, Lappe M, Ukkonen E, Brazma A. From gene networks to gene function. *Genome Res.* 2003; 13:2568-2576.
 49. Yi M, Horton JD, Cohen JC, Hobbs HH, Stephens RM. WholePathwayScope: a comprehensive pathway-based analysis tool for high-throughput data. *BMC Bioinformatics.* 2006; 7:30.
 50. Guil S, Esteller M. Cis-acting noncoding RNAs: friends and foes. *Nat Struct Mol Biol.* 2012;19:1068-1075.

FIGURE LEGENDS

Figure 1 Hematoxylin and eosin staining of eutopic and ectopic endometria in adenomyosis. (a) Normal uterine endometrium. (b) Ectopic endometrium within the myometrium. G, gland.

Figure 2 Relative fold-changes in expression between ectopic and eutopic endometria, as detected by quantitative PCR (light gray bars), were in agreement with microarray data (dark gray bars).

Figure 3 Bioinformatic analysis for differentially expressed mRNAs and lncRNAs. (a) GO analysis of mRNAs up-regulated in ectopic endometria. (b) GO analysis of mRNAs down-regulated in ectopic endometria. (c) Top pathway annotations of up-regulated mRNAs may involved in. (d) Top pathway annotations of down-regulated mRNAs may involved in. (e) The interaction network of differentially expressed genes (Signal-net). Circles represent important functional genes in adenomyosis (red: up-regulated genes; blue: down-regulated genes). Solid arrow represents activation; dotted arrow represents indirect effect; solid line represents compound or binding/association.

Figure 4 lncRNA–mRNA core network consisting of 24 lncRNAs and their

mRNAs with the most credibility.

Table 1 Primers used in real-time PCR.

ID	Forward	Reverse
GAPDH	CAGCCGCATCTTCTTTTGCG	AGTTAAAAGCAGCCCTGGTGA
UC004dwe.2	GATCCCAGCCAGCCTAGAAG	AAATGCTGGAAAGGCCCTCAT
ENST00000454594	CGTTCGTTGTACAGAGGAGGA	CTGCAAGTGGGAGCGGAAA
NR_003521	GCTGCGTCAGGTAAGCGAG	CCGGCTCCTCCCCATCAG
NRP2	CAAGACAGGCTCTGAAGATTGC	GGGTTTGGCCAGGATGGTAA
TGFBR1	TTCCGTGAGGCAGAGATT	GCAATACAGCAAGTTCCATT
ITGB3	AAGAGCCAGAGTGTCCTCAAG	GTGCCCCGGTACGTGATATT

Figure 1




Figure 2




Figure 3





Figure 4



Supplementary Material

[Click here to download Supplementary Material: Additional file 1.xls](#)

Supplementary Material

[Click here to download Supplementary Material: Additional file 2.xlsx](#)