

Bashir Al Hussein Al Awamlh, M.D.

New York Presbyterian
New York, NY

Jim C. Hu, M.D., M.P.H.

Jonathan E. Shoag, M.D.

Weill Cornell Medicine
New York, NY
jch9011@med.cornell.edu

and Others

A complete list of authors is available with the full text of this letter at NEJM.org.

Supported by the Frederick J. and Theresa Dow Wallace Fund of the New York Community Trust (Drs. Hu and Shoag), by an award (to Dr. Shoag) from the Damon Runyon Cancer Research Foundation, and by a grant (1-UL1-TR002384-01) from the Clinical and Translational Science Center at Weill Cornell Medicine.

Disclosure forms provided by the authors are available with the full text of this letter at NEJM.org.

1. Butler S, Muralidhar V, Chavez J, et al. Active surveillance for low-risk prostate cancer in black patients. *N Engl J Med* 2019; 380:2070-2.
2. Lu-Yao GL, McLerran D, Wasson J, Wennberg JE. An assessment of radical prostatectomy: time trends, geographic variation, and outcomes. *JAMA* 1993;269:2633-6.
3. Broughman JR, Basak R, Nielsen ME, et al. Prostate cancer patient characteristics associated with a strong preference to preserve sexual function and receipt of active surveillance. *J Natl Cancer Inst* 2018;110:420-5.
4. Stanford JL, Feng Z, Hamilton AS, et al. Urinary and sexual function after radical prostatectomy for clinically localized prostate cancer: the Prostate Cancer Outcomes Study. *JAMA* 2000;283:354-60.

DOI: 10.1056/NEJMc1912868




Figure 1. Odds of Undergoing Active Surveillance for Low-Risk Prostate Cancer among Black Men, as Compared with Men of Other Races.

Shown are the adjusted odds of undergoing active surveillance or definitive treatment (radiation therapy or radical prostatectomy) for low-risk prostate cancer (defined as a prostate-specific antigen level of <10 ng per milliliter, a clinical stage of T1 to T2a, and a Gleason score of 6) among black men, as compared with men of other races, with or without adjustment for geographic region during the period from 2010 through 2015. I bars indicate 95% confidence intervals.

effect on the selection of active surveillance as compared with definitive treatment. Given the presence of such confounding factors, further study is needed to ascertain the true effect of geography and race on the choice of management for low-risk prostate cancer in the United States.

Volanesorsen, Familial Chylomicronemia Syndrome, and Thrombocytopenia

TO THE EDITOR: In the APPROACH trial (Aug. 8 issue),¹ volanesorsen — an antisense oligonucleotide targeting *APOC3* messenger RNA — reduced circulating triglycerides in patients with familial chylomicronemia syndrome, but thrombocytopenia developed in 76% of treated patients. This unexpected adverse event raises a key question regarding inhibition of apolipoprotein C-III synthesis as a therapeutic strategy: is thrombocytopenia an on-target effect of reduced apolipoprotein C-III activity or an off-target effect of volanesorsen?

We therefore asked whether heterozygous carriers of an inactivating mutation in *APOC3* are at increased risk for thrombocytopenia. Among 42,503 participants in the UK Biobank, 235 (0.6%)

were found to carry any of four previously described inactivating mutations in *APOC3*.² As expected, carriers had significantly lower levels of triglycerides than noncarriers (median, 71 mg per deciliter and 129 mg per deciliter, respectively; $P < 0.001$), as well as a lower risk of hypertriglyceridemia (odds ratio, 0.05; 95% CI, 0.03 to 0.11; $P < 0.001$) (Fig. 1A and 1B). We found no significant difference between carriers and noncarriers in platelet count (median, 239,000 and 238,000 per microliter, respectively; $P = 0.21$) or in the prevalence of thrombocytopenia (1.3% and 1.8%, respectively; $P = 0.63$) (Fig. 1C and 1D). These results suggest that the effect of volanesorsen is medication- or class-specific rather than an inherent property of apolipoprotein C-III inhibition.




Figure 1. Inactivating Mutations in *APOC3* and the Risk of Hypertriglyceridemia or Thrombocytopenia.

Gene sequencing of specimens from 42,503 participants in the UK Biobank identified an inactivating mutation in *APOC3* in 235 (0.6%). Panel A shows the distribution of triglyceride levels in carriers and noncarriers. The vertical dashed line indicates a triglyceride level of 150 mg per deciliter, the cutoff value for hypertriglyceridemia. Panel B shows the percentage of participants with hypertriglyceridemia. In a logistic-regression model with adjustment for age, sex, fasting duration at time of blood collection, and genetic ancestry, carriers had a significantly lower risk of hypertriglyceridemia than noncarriers. Panel C shows the distribution of platelet counts in carriers and noncarriers. In a linear regression model with adjustment for age, sex, fasting duration at time of blood collection, and genetic ancestry, there was no significant difference in platelet counts between carriers and noncarriers. The vertical dashed line indicates 140,000 per microliter, the cutoff value for thrombocytopenia. Panel D shows the percentage of participants with thrombocytopenia. In a logistic-regression model with adjustment for age, sex, fasting duration at time of blood collection, and genetic ancestry, there was no evidence of a higher risk of thrombocytopenia among carriers than among noncarriers. To convert the values for triglycerides to millimoles per liter, multiply by 0.01129.

Sumeet A. Khetarpal, M.D., Ph.D.

Massachusetts General Hospital
Boston, MA

Minxian Wang, Ph.D.

Broad Institute of MIT and Harvard
Cambridge, MA

Amit V. Khera, M.D.

Massachusetts General Hospital
Boston, MA
avkhera@mgh.harvard.edu

Drs. Khetarpal and Wang contributed equally to this letter.

Dr. Khera reports being supported by an institutional grant from the Broad Institute of MIT and Harvard (BroadIgnite), grant 1K08HG010155 from the National Human Genome Research Institute, a Hassenfeld Scholar Award from Massachusetts General Hospital, having served as a consultant to Color Genomics and Navitor Pharmaceuticals, having received honoraria from Illumina, having received grant support from the Novartis Institute for Biomedical Research and IBM Research, and having a patent application entitled “Genetic Risk Predictor” (20190017119) owned by the General Hospital Corporation. No other potential conflict of interest relevant to this letter was reported.

1. Witztum JL, Gaudet D, Freedman SD, et al. Volanesorsen and triglyceride levels in familial chylomicronemia syndrome. *N Engl J Med* 2019;381:531-42.
2. The TG and HDL Working Group of the Exome Sequencing Project, National Heart, Lung, and Blood Institute. Loss-of-function mutations in *APOC3*, triglycerides, and coronary disease. *N Engl J Med* 2014;371:22-31.

DOI: 10.1056/NEJMc1912350

THE AUTHORS REPLY: We previously conducted a thorough database assessment and found no antisense oligonucleotide class-specific effect on platelets.¹ Patients with familial chylomicronemia syndrome, who often have triglyceride levels greater than 2000 mg per deciliter, have an increased incidence of both thrombocytopenia and thrombocytosis.² As noted by Khetarpal and colleagues, in the APPROACH trial, thrombocytopenia developed in 24% of participants with familial chylomicronemia syndrome who received placebo and in 76% of those who received volanesorsen, suggesting that there was a volanesorsen-disease interaction in the trial participants, an effect we have not seen to this degree in other populations who have received the same dose (unpublished data). Thrombocytopenia is dose-dependent, and core management of familial chylomicronemia syndrome in patients who receive volanesorsen includes monitoring for this

effect and early intervention to mitigate and maintain patients on treatment. We are currently investigating a galactosamine-conjugated *APOC3* oligonucleotide that, in a phase 1-2a study conducted over the course of 3 months, achieved similar levels of triglyceride reduction at lower doses, with no reported changes in platelet counts.³

Joseph L. Witztum, M.D.

University of California, San Diego
La Jolla, CA
jwitztum@ucsd.edu

Richard S. Geary, Ph.D.

Ionis Pharmaceuticals
Carlsbad, CA

Louis O'Dea, Ph.D.

Akcea Therapeutics
Cambridge, MA

Since publication of their article, the authors report no further potential conflict of interest.

1. Crooke ST, Baker BF, Witztum JL, et al. The effects of 2'-O-methoxyethyl containing antisense oligonucleotides on platelets in human clinical trials. *Nucleic Acid Ther* 2017;27:121-9.

2. Gaudet D, Baass A, Tremblay A, et al. Natural history (up to 15 years) of platelet count in 84 patients with familial hyperchylomicronemia due to lipoprotein lipase deficiency. *J Clin Lipidol* 2017;11:797-8. abstract.

3. Alexander VJ, Xia S, Hurh E, et al. N-acetyl galactosamine-conjugated antisense drug to *APOC3* mRNA, triglycerides and atherogenic lipoprotein levels. *Eur Heart J* 2019;40:2785-96.

DOI: 10.1056/NEJMc1912350

Gender Parity in Clinical PrEP Trials

TO THE EDITOR: The Perspective article by Goldstein and Walensky (in this issue of the *Journal*; originally published online on Oct. 30)¹ addresses the approval by the Food and Drug Administration (FDA) of a new human immunodeficiency virus (HIV)-prevention indication for emtricitabine and tenofovir alafenamide (F/TAF) that unfortunately excludes cisgender (nontransgender) women because of a lack of clinical data. Approval was based on a single noninferiority trial, DISCOVER, that compared F/TAF with emtricitabine and tenofovir disoproxil fumarate (F/TDF), primarily in men who have sex with men. When DISCOVER was designed, the FDA encouraged the sponsor to conduct a second trial involving women.² DISCOVER excluded women because of a lack of consensus among stakeholders regarding a scientifically rigorous design to assess efficacy in women in the era of F/TDF, not because

of a lack of forethought or disregard for diversity. A noninferiority trial such as DISCOVER that used F/TDF as a control in women was not considered to be scientifically appropriate because previous trials had not shown a consistently positive treatment effect of F/TDF in women³; constancy of a control's treatment effect is necessary to justify the noninferiority margin and to ensure an interpretable study result.⁴ Other designs, such as superiority to placebo or to F/TDF, were considered to be unethical or unlikely to succeed, respectively. In fact, although F/TAF was found to be noninferior to F/TDF in DISCOVER, it was not superior.

Going forward, the sponsor will conduct a trial involving women in high-incidence settings with a primary efficacy comparison between women receiving F/TAF and two external control groups (without F/TDF).⁵ Although the trial will