

Journal of Thoracic Oncology

Silencing NKD2 by promoter region hypermethylation promotes esophageal cancer progression by activating Wnt signaling

--Manuscript Draft--

Manuscript Number:	JTO-D-15-01271
Full Title:	Silencing NKD2 by promoter region hypermethylation promotes esophageal cancer progression by activating Wnt signaling
Article Type:	Original Article
Order of Authors:	Baoping Cao, M.D., Ph.D. Weili Yang, M.D., Ph.D. Yongshuai Jin, M.D. Meiying Zhang, Ph.D. Tao He, M.D. Qimin Zhan, M.D., Ph.D. James G Herman, M.D. Guanglin Zhong, M.D. Mingzhou Guo, M.D., Ph.D.
Manuscript Region of Origin:	CHINA
Abstract:	<p>Introduction: Naked cuticle homolog 2 (NKD2) was found frequently methylated in human breast and gastric cancer. The epigenetic changes and mechanisms of NKD2 in human esophageal cancer remain unclear. Methods: Nine esophageal cancer cell lines and 154 cases of primary esophageal cancer samples were analyzed using methylation specific PCR, immunohistochemistry, western blot and a xenograft mouse model. Results: Loss of NKD2 expression and complete methylation were found in KYSE150 and TE1 cells. Reduced expression and partial methylation were observed in KYSE30, KYSE70, KYSE410, KYSE140 and COLO680 cells. High level expression and unmethylation were detected in KYSE450 and TE8 cells. Re-expression of NKD2 was induced by 5-aza-2'-deoxycytidine in NKD2 unexpressed or reduced cells. NKD2 was methylated in 53.2% (82/154) of human primary esophageal cancer samples, and promoter region hypermethylation was associated with reduced expression of NKD2 significantly ($p<0.01$). NKD2 methylation was associated with TNM stage and lymph node metastasis ($p<0.01$). The results suggest that NKD2 is regulated by promoter region methylation and methylation of NKD2 may serve as a prognostic marker in esophageal cancer. Our further studies demonstrate that NKD2 suppresses cell proliferation, colony formation, cell invasion and migration, as well as induces G1/S check point arrest in esophageal cancer cells. NKD2 suppressed xenograft tumor growth and inhibited Wnt signaling in human esophageal cancer cells. Conclusions: NKD2 is frequently methylated in human esophageal cancer, and the expression of NKD2 is regulated by promoter region methylation. NKD2 suppresses esophageal cancer progression by inhibiting Wnt signaling both in vitro and in vivo.</p>
Keywords:	NKD2, DNA methylation, Wnt signaling, esophageal cancer

Silencing NKD2 by promoter region hypermethylation promotes
esophageal cancer progression by activating Wnt signaling

*Running title: NKD2 suppresses esophageal cancer growth by inhibiting
Wnt signaling*

Baoping Cao^{1#}, Weili Yang^{1,2#}, Yongshuai Jin¹, Meiying Zhang^{1,2},
Tao He¹, Qimin Zhan³, James G. Herman⁴, Guanglin Zhong^{5*} and
Mingzhou Guo^{1*}.

Authors' Affiliations:

1. Department of Gastroenterology & Hepatology, Chinese PLA General
Hospital, #28 Fuxing Road, Beijing 100853, China
2. Medical College of NanKai University, #94 Weijin Road, Tianjin
300071, China
3. State Key Laboratory of Molecular Oncology, Cancer Institute and
Hospital, Chinese Academy of Medical Sciences & Peking Union
Medical College, Beijing 100021, P.R. China
4. The Hillman Cancer Center, University of Pittsburgh Cancer Institute,
5117 Centre Avenue, Suite 2.18/Research, Pittsburgh, PA 15213, USA
5. Department of Internal Medicine, Chinese PLA General Hospital, #28
Fuxing Road, Beijing 100853, China.

Baoping Cao and Weili Yang contributed equally to this work.

*Correspondence to:

1 Mingzhou Guo: Department of Gastroenterology & Hepatology, Chinese
2
3 PLA General Hospital, #28 Fuxing Road, Beijing 100853, China. Phone:
4
5 +86-10-66937651; Fax: +86-10-68180325; E-mail: mzguo@hotmail.com
6
7

8
9 Guanglin Zhong, M.D., Department of Internal Medicine, Chinese PLA
10
11 General Hospital, #28 Fuxing Road, Beijing 100853, China. E-mail:
12
13 guanglin1963@aliyun.com
14
15

16
17 E-Mails:
18
19

20 Baoping Cao: goddenlove@163.com
21

22 Weili Yang: yangweilipuyang@163.com
23

24
25 Yongshuai Jin: jinyongshuai2012@126.com
26

27
28 Meiying Zhang: zhangmeiying.1988@163.com
29

30
31 Tao He: hetao_1981@163.com
32

33
34 Qimin Zhan: zhanqimin@pumc.edu.cn
35

36
37 James G. Herman: Hermanj3@upmc.edu
38

39
40 Guanglin Zhong: guanglin1963@aliyun.com
41

42
43 Mingzhou Guo: mzguo@hotmail.com
44

45 **Acknowledgments**

46
47

48
49 This work was supported by the following grants: National Basic
50
51 Research Program of China (973 Program No. 2012CB934002,
52
53 2015CB553904); National High-tech R&D Program of China (863
54
55 Program No. SS2012AA020314, SS2012AA020821, SS2012AA020303);
56
57
58
59
60 National Key Scientific Instrument Special Programme of China (Grant
61
62
63
64
65

No. 2011YQ03013405) and National Science Foundation of China (NSFC No.81402345, 81121004, 81161120432, 81490753 and 81401950).

ABSTRACT

Introduction: Naked cuticle homolog 2 (NKD2) was found frequently methylated in human breast and gastric cancer. The epigenetic changes and mechanisms of NKD2 in human esophageal cancer remain unclear. **Methods:** Nine esophageal cancer cell lines and 154 cases of primary esophageal cancer samples were analyzed using methylation specific PCR, immunohistochemistry, western blot and a xenograft mouse model. **Results:** Loss of NKD2 expression and complete methylation were found in KYSE150 and TE1 cells. Reduced expression and partial methylation were observed in KYSE30, KYSE70, KYSE410, KYSE140 and COLO680 cells. High level expression and unmethylation were detected in KYSE450 and TE8 cells. Re-expression of NKD2 was induced by 5-aza-2'-deoxycytidine in NKD2 unexpressed or reduced cells. NKD2 was methylated in 53.2% (82/154) of human primary esophageal cancer samples, and promoter region hypermethylation was associated with reduced expression of NKD2 significantly ($p<0.01$). NKD2 methylation was associated with TNM stage and lymph node metastasis ($p<0.01$). The results suggest that NKD2 is regulated by

promoter region methylation and methylation of NKD2 may serve as a prognostic marker in esophageal cancer. Our further studies demonstrate that NKD2 suppresses cell proliferation, colony formation, cell invasion and migration, as well as induces G1/S check point arrest in esophageal cancer cells. NKD2 suppressed xenograft tumor growth and inhibited Wnt signaling in human esophageal cancer cells. **Conclusions:** NKD2 is frequently methylated in human esophageal cancer, and the expression of NKD2 is regulated by promoter region methylation. NKD2 suppresses esophageal cancer progression by inhibiting Wnt signaling both *in vitro* and *in vivo*.

Keywords: NKD2, DNA methylation, Wnt signaling, esophageal cancer

Introduction

Esophageal cancer is the eighth most common cancer and the sixth leading cause of cancer-related death worldwide ¹. The overall 5-year survival remains below 15% ². Poor outcomes in patients with esophageal cancer are related to diagnosis at advanced (metastatic) stages and the propensity for metastases ³. Esophageal squamous cell carcinoma (ESCC) is the predominant histological type of esophageal carcinoma worldwide ⁴. Tobacco use and alcohol consumption are risk factors for ESCC, and the combination of tobacco and alcohol consumption further increases the risk of ESCC. Mutations in enzymes that metabolize alcohol have been associated with increased risk of ESCC ⁵⁻⁷. Genetic and epigenetic

alterations are involved in esophageal carcinogenesis⁸. Aberrant expression of components in Wnt signaling pathway are found in many types of cancers, including esophageal cancer, and Wnt signaling pathway plays an important role in cancer progression^{9, 10}.

The naked cuticle (NKD) family includes *Drosophila* naked cuticle and its two vertebrate orthologs, NKD1 and NKD2. NKD1 is located in human chromosome 16q12.1, which has frequent loss of heterozygosity in human breast and hepatocellular carcinoma^{11, 12}. NKD2 is located in chromosome 5p15.3. Loss of heterozygosity has been frequently found in these regions in multiple tumors¹³⁻¹⁵. In both zebrafish and mice, NKD inhibits canonical and non-canonical Wnt signaling¹⁶⁻¹⁸. The C-terminus of NKD2 is highly disordered, while the N-terminal region of NKD2 contains most of the functional domain, including myristoylation, an EF-hand motif, a Dishevelled binding region, and a vesicle recognition and membrane targeting motif¹⁹⁻²¹. NKD2 binds to multiple proteins and may function as a switch protein through its several functional motifs²². Both NKD1 and NKD2 have been proposed to interact with Dishevelled through their EF-hand-like motif. In addition, NKD2 has been reported to bind to Dishevelled through its TGF α binding region^{21, 22}. NKD2 was reported to suppress tumor growth and metastasis in osteosarcoma through negative regulation of Wnt signaling²³. Our previous study found that methylation of NKD2 promotes breast cancer growth by activating

Wnt signaling²⁴. The methylation status and the function of NKD2 in esophageal cancer have yet to be elucidated. Therefore in this study we investigated the epigenetic changes and functions of NKD2 in human ESCC.

Materials and Methods

Human tissue samples and cell lines

Fifteen cases of human normal esophageal mucosa and 154 cases of human esophageal cancer samples were collected from the Chinese PLA General Hospital in Beijing. The median age of the cancer patients is 62.1 years old (range 46-87), and the ratio of males/females is 3.05:1. All cancer samples were classified according to TNM staging (AJCC 2010), including 5 cases of stage I, 99 cases of stage II and 50 cases of stage III. All samples were collected following the guidelines approved by the Institutional Review Board of the Chinese PLA General Hospital with written informed consent from patients (Reference No. 20090701-015).

Nine esophageal cancer cell lines (KYSE450, KYSE30, KYSE150, KESE70, TE8, KYSE410, TE1, KYSE140 and COLO680) were previously established from primary esophageal cancer and maintained in 90% RPMI 1640 (Invitrogen, Carlsbad, CA) supplemented with 10% fetal bovine serum.

5-aza-2'-deoxycytidine treatment

Esophageal cancer cell lines were split to a low density (30%

1 confluence) 12 hours before treatment. Cells were treated with
2
3 5-aza-2'-deoxycytidine (5-AZA, Sigma, St. Louis, MO) at a
4
5 concentration of 2 μ M. Growth medium conditioned with 5-AZA at a
6
7 concentration of 2 μ M was exchanged every 24 hours for a total of 96
8
9 hours of treatment.
10
11
12

13 **RNA isolation and semi-quantitative reverse transcription PCR**

14
15
16
17 Total RNA was isolated by Trizol reagent (Life Technologies,
18
19 Gaithersburg, MD). First strand cDNA was synthesized according to the
20
21 manufacturer's instructions (Invitrogen, Carlsbad, CA). PCR primers for
22
23 NKD2 are 5'-ACAGGAGGTTGTCTGCACACG-3' (F) and
24
25 5'-GACTTGAGGAACTGCTTCTCC-3' (R). The primer sets for NKD2
26
27 were designed to span intronic sequences between adjacent exons in order
28
29 to control for genomic DNA contamination. Semi-quantitative reverse
30
31 transcription PCR (RT-PCR) was amplified for 33 cycles. GAPDH was
32
33 used as an internal control.
34
35
36
37
38
39
40
41

42 **Bisulfite modification, methylation-specific PCR and bisulfite** 43 **sequencing**

44
45
46
47 DNA was prepared by the proteinase K method. Bisulfite treatment
48
49 was carried out as previously described^{25, 26}. Methylation-specific PCR
50
51 (MSP) primers were designed according to genomic sequences around
52
53 transcriptional start sites (TSS) and synthesized to detect unmethylated
54
55 (U) and methylated (M) alleles. Bisulfite sequencing (BSSQ) was
56
57
58
59
60
61
62
63
64
65

performed as previously described⁹. BSSQ products were amplified by primers flanking the targeted regions including MSP products. The MSP primers are as follows: NKD2-M-Forward 5'-GAGGTCGATGCGTTGCGGGTAGC-3' and NKD2-M-Reverse 5'-CGACGACCCGACTCCCTCTAAACG-3'; NKD2-U-Forward 5'-GTTGAGGTTGATGTGTTGTGGGTAGT-3' and NKD2-U-Reverse 5'-CAACAACAACCCAACTCCCTCTAAACA-3'. The bisulfite sequencing primers are 5'-GTTGGTGGGGTTTTAGGTTGG-3' (F) and 5'-AACTAAATTCTAAAACCRAAACC-3' (R).

Immunohistochemistry

Immunohistochemistry (IHC) was performed in human esophageal cancer samples and paired adjacent tissue samples. The NKD2 antibody was diluted 1:500 (Novus Biology, CO, USA). The staining intensity and extent of the staining area were scored using the German semi-quantitative scoring system as described previously^{27, 28}.

Plasmid construction

Human full-length NKD2 CDS (GenBank accession number NM_033120) was amplified and subcloned as described previously²⁴. The primers used were 5'-GAGGATCCGCCACCATGGGGAACTGCAGTCGAAG-3' (F) and 5'-GATCTCGAGCTAGGACGGGTGGAAGTGGT-3' (R). NKD2 expressing lentiviral or empty vectors were packaged using the

1 ViraPower™ lentiviral expression system (Invitrogen, San Diego, CA,
2 USA). Lentivirus was added to the growing medium of KYSE150 and
3
4 TE1 cells, and NKD2 stably expressed cells were selected by blasticidin
5
6 (Invitrogen, San Diego, CA, USA) at a concentration of 2μg/ml.
7
8
9

10 11 **Cell viability detection**

12
13
14 Cells were plated into 96-well plates at 2×10^3 cells/well, and the cell
15
16 viability was measured by MTT assay (KeyGEN Biotech, Nanjing, China)
17
18 at 0, 24, 48 and 72h. Absorbance was measured on a microplate reader
19
20 (Thermo Multiskan MK3, MA, USA) at a wavelength of 490 nm.
21
22
23
24

25 26 **Colony formation assay**

27
28 NKD2 unexpressed and stably expressed cells were seeded at 500
29
30 cells per well in 6-well culture plates in triplicate. The complete growth
31
32 medium conditioned with blasticidin at 2ug/ml was exchanged every 72
33
34 hours. After 2 weeks, cells were fixed with 75% ethanol for 30 min and
35
36 stained with 0.2% crystal violet (Beyotime, Nanjing, China) for
37
38 visualization and counting.
39
40
41
42
43

44 45 **Flow cytometry**

46
47 NKD2 unexpressed and re-expressed KYSE150 and TE1 cells were
48
49 starved 12 hours for synchronization, and the cells were re-stimulated
50
51 with 10% FBS for 24 hours. Cells were fixed with 70% ethanol and
52
53 treated using the Cell Cycle Detection Kit (KeyGen Biotech, Nanjing,
54
55 China). The cells were then sorted by a FACS Caliber flow cytometer
56
57
58
59
60
61
62
63
64
65

(BD Biosciences, Mansfield, CA). The cell phase distribution was analyzed by the Modfit software (Verity Software House, ME, USA).

Transwell assay

NKD2 unexpressed and re-expressed KYSE150 and TE1 cells were suspended in serum-free medium. Cells (2×10^5) were placed into the upper chamber of an 8 μ m pore size transwell apparatus (Corning, NY, USA) and incubated for 20 hours. Cells that migrated to the lower surface of the membrane were stained with crystal violet and counted in three independent high-power fields ($\times 200$). For invasion analysis, NKD2 unexpressed and re-expressed KYSE150 and TE1 cells (2×10^5) were seeded into the upper chamber of a transwell apparatus coated with extracellular matrix gel (ECM gel, BD Biosciences, San Jose, CA) and incubated for 36 hours. Cells that invaded into the lower membrane surface were stained with crystal violet and counted in three independent high-power fields ($\times 200$).

SiRNA knockdown technique

Selected siRNAs targeting NKD2 and the RNAi negative control duplex were used in this study. The sequences of the siRNAs targeting NKD2 and the RNAi negative control are as follows: NKD2-F: 5'-GGGAUUGAGAACUACACGUTT-3' , NKD2-R: 5'-ACGUGUAGUUCUCAAUCCCTT-3' , Negative Control-F: 5'-UUCUCCGAACGUGUCACGUTT-3' and Negative Control-R:

1 5'-ACGUGACACGUUCGGAGAATT-3'. The RNAi oligonucleotide
2
3 and RNAi negative control duplex were transfected into KYSE450 cells,
4
5 which expressed high levels of NKD2.
6
7

8 **In vivo tumorigenicity**

9

10 NKD2 stably expressed and unexpressed KYSE150 cells (4×10^6
11
12 cells in 0.2 ml phosphate-buffered saline) were subcutaneously injected
13
14 into the dorsal flank of 5-week-old female BALB/c nude mice. The tumor
15
16 size was measured every 3 days for 24 days beginning 3 days after
17
18 implantation. The tumor volumes were calculated according to the
19
20 following formula: $V = L \times W^2/2$, where V, volume (mm^3); L, biggest
21
22 diameter (mm); W, smallest diameter (mm). All procedures were
23
24 approved by the Animal Ethics Committee of the Chinese PLA General
25
26 Hospital.
27
28
29
30
31
32
33
34
35

36 **Western blot**

37

38 Protein samples from esophageal cancer cells were collected and
39
40 western blot was performed as described previously ²⁹. Antibodies were
41
42 diluted according to manufacturer's instructions. The primary antibodies
43
44 were as follows: NKD2 (Cell Signaling Technology, Danvers, MA),
45
46 MMP2, MMP7, MMP9, cyclin D1, c-myc, p- β -catenin, β -catenin and
47
48 β -actin (Bioworld Technology, MN, USA).
49
50
51
52
53
54
55

56 **Statistical analysis**

57

58 SPSS 17.0 software (IBM, NY, USA) was used for data analysis. All
59
60
61
62
63
64
65

1 data were presented as means \pm standard deviation (SD) and analyzed
2
3 using the Student's t test. The Chi-squared test and the Fisher's exact test
4
5 were used to analyze the association of NKD2 methylation status with
6
7 clinic-pathologic factors and the association of NKD2 expression with
8
9 methylation status. The value of $p < 0.05$ was considered to be
10
11 statistically significant.
12
13
14
15

16 RESULTS

17 NKD2 expression is regulated by promoter region methylation in 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65

The expression of NKD2 was detected by semi-quantitative RT-PCR in human esophageal cancer cell lines. As shown in Figure 1A, loss of NKD2 expression was found in KYSE150 and TE1 cells. Reduced expression of NKD2 was observed in KYSE30, KYSE70, KYSE410, KYSE140 and COLO680 cells. High level expression of NKD2 was detected in KYSE450 and TE8 cells. The methylation status of the NKD2 promoter was examined by MSP. Complete methylation was found in KYSE150 and TE1 cells. Partial methylation was detected in KYSE30, KYSE70, KYSE410, KYSE140 and COLO680 cells. Unmethylation was observed in KYSE450 and TE8 cells (Figure 1B). These results demonstrate that loss of expression or reduced expression of NKD2 correlated with promoter region methylation in human esophageal cancer cells. To further reveal the methylation density and validate the MSP

1 results, BSSQ technique was used. As shown in Figure 1C, NKD2 was
2
3 completely methylated in KYSE150 and TE1 cells, partially methylated
4
5 in KYSE410 cells, and unmethylated in KYSE450 cells and normal
6
7 esophageal mucosa. The results are consistent with MSP results (Figure
8
9 1C). To further analyze NKD2 expression is regulated by promoter
10
11 region methylation, KYSE450, KYSE30, KYSE150, KYSE70, TE8,
12
13 KYSE410, TE1, KYSE140 and COLO680 cells were treated with 5-AZA,
14
15 a demethylating reagent. Restoration of NKD2 expression was induced
16
17 by 5-AZA in KYSE150 and TE1 cells. Increased expression of NKD2
18
19 was observed in KYSE30, KYSE70, KYSE410, KYSE140 and
20
21 COLO680 cells treated with 5-AZA, while, no expression changes were
22
23 found in KYSE450 and TE8 cells before and after 5-AZA treatment
24
25 (Figure 1A). These results suggest that the expression of NKD2 is
26
27 regulated by promoter region methylation in human esophageal cancer
28
29 cells.
30
31
32
33
34
35
36
37
38
39
40
41

42 **NKD2 is frequently methylated in primary human esophageal cancer**

43
44 To further explore the methylation status of NKD2 in primary
45
46 human esophageal cancer, the methylation status was examined by MSP
47
48 in 154 cases of esophageal cancer tissue samples and 15 cases of normal
49
50 esophageal mucosa from non-cancerous patients. NKD2 was methylated
51
52 in 53.2% (82/154) of primary esophageal cancer samples, and no
53
54 methylation was detected in normal esophageal mucosa (Figure 2A and
55
56
57
58
59
60
61
62
63
64
65

1 B). As shown in table 1, NKD2 methylation was associated with TNM
2 stage and lymph node metastasis significantly (both $p < 0.01$), but no
3 association was found between NKD2 methylation and age, gender,
4 tumor size and differentiation (all $p > 0.05$). To further validate that
5 NKD2 expression is regulated by promoter region methylation, 30 cases
6 of available matched esophageal cancer and adjacent tissue paraffin
7 samples were evaluated by IHC. NKD2 staining was observed mainly in
8 the cytoplasm of the esophagus. NKD2 is highly expressed in adjacent
9 tissue samples and reduced in primary cancer tissue samples (Figure 2C
10 and D). Reduced expression of NKD2 is associated with the promoter
11 region hypermethylation ($p < 0.01$, Figure 2E). These results demonstrate
12 that NKD2 is regulated by promoter region methylation in primary
13 esophageal cancer.

14 **Restoration of NKD2 expression suppresses cell proliferation and** 15 **induces G1/S arrest in esophageal cancer cells**

16 To evaluate the effects of NKD2 on cell proliferation, the cell
17 viability was detected by MTT and colony formation assays. The OD
18 value was 0.892 ± 0.027 vs. 0.763 ± 0.024 ($p < 0.05$) in KYSE150 cells
19 and 0.551 ± 0.024 vs 0.438 ± 0.011 ($p < 0.001$) in TE1 cells before and
20 after restoration of NKD2 expression (Figure 3A). The results
21 demonstrated that NKD2 inhibited esophageal cancer cell viability. The
22 effect of NKD2 on cell proliferation was evaluated by colony formation

1 assay. The clone numbers were 135.3 ± 6.8 vs. 57.7 ± 4.0 ($p < 0.001$) in
2
3 KYSE150 cells and 58.3 ± 4.7 vs. 29.7 ± 3.5 ($p < 0.001$) in TE1 cells
4
5 before and after restoration of NKD2 expression (Figure 3B). These
6
7 results suggest that NKD2 suppresses esophageal cancer cell growth.
8
9

10
11 To further understand the mechanism of NKD2 in esophageal cancer
12 development, the role of NKD2 in cell cycle was analyzed by flow
13 cytometry. In KYSE150 cells, the cell phase distribution before and after
14 re-expression of NKD2 was as follows: G0/1 phase: $33.46 \pm 0.58\%$ vs.
15 $41.82 \pm 1.73\%$, S phase: $46.17 \pm 2.21\%$ vs. $37.15 \pm 1.46\%$, and G2/M
16 phase: $20.37 \pm 2.21\%$ vs. $21.03 \pm 0.31\%$. The G0/1 phase is increased and
17 the S phase is reduced significantly after re-expression of NKD2 (all $p <$
18 0.01).
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

35 In TE1 cells, the cell phase distribution before and after re-expression
36 of NKD2 was as follows: G0/1 phase: $44.13 \pm 2.60\%$ vs. $61.73 \pm 1.28\%$,
37 S phase: $44.21 \pm 3.88\%$ vs. $22.93 \pm 1.77\%$, and G2/M phase: $11.67 \pm$
38 1.38% vs. $15.34 \pm 0.82\%$ (Figure 3C). The G0/1 phase is increased and
39 the S phase is reduced significantly after re-expression of NKD2 in TE1
40 cells (all $p < 0.001$). These results suggest that NKD2 induced G1/S check
41 point arrest in esophageal cancer cells.
42
43
44
45
46
47
48
49
50
51
52
53
54

55 **Restoration of NKD2 expression inhibits cell migration and invasion**
56
57 **in human esophageal cancer cells**
58
59
60
61
62
63
64
65

1 The transwell assay in the absence of ECM gel coating was
2
3 employed to explore the effect of NKD2 on cell migration. The number
4
5 of migrated cells for each high power field under the microscope was
6
7
8
9 105.7 ± 5.1 vs. 63.0 ± 4.0 in KYSE150 cells and 147.0 ± 6.6 vs. $52.3 \pm$
10
11 5.7 in TE1 cells before and after restoration of NKD2 expression. The
12
13 cell number was reduced significantly after re-expression of NKD2 in
14
15 esophageal cancer cells (all $p < 0.001$, Figure 4A). These results
16
17 demonstrate that NKD2 inhibits esophageal cancer cell migration.
18
19
20
21

22 Next, the transwell assay with ECM coating was employed to
23
24 evaluate the effect of NKD2 on cell invasion. The number of invasive
25
26 cells for each high power field under the microscope was 114.7 ± 4.5 vs.
27
28 79.7 ± 4.5 in KYSE150 cells and 137.0 ± 4.0 vs. 65.0 ± 2.7 in TE1 cells
29
30 before and after restoration of NKD2 expression. The cell number was
31
32 reduced significantly after re-expression of NKD2 in KYSE150 and TE1
33
34 cells (all $p < 0.001$, Figure 4B). These results suggest that NKD2 impedes
35
36 esophageal cancer cell invasion.
37
38
39
40
41
42
43
44

45 To further understand the mechanism of NKD2 in esophageal cancer
46
47 migration and invasion, the expression levels of MMP2, MMP7 and
48
49 MMP9 were detected by western blot. As shown in Figure 4C, the
50
51 expression levels of MMP2, MMP7 and MMP9 were reduced after
52
53 re-expression of NKD2 in KYSE150 and TE1 cells. The inhibitory role of
54
55 NKD2 on MMP2, MMP7 and MMP9 expression was further validated by
56
57
58
59
60
61
62
63
64
65

1 knocking down NKD2 in KYSE450 cells. Taken together, the above
2
3 results suggest that NKD2 suppresses esophageal cancer cell migration
4
5 and invasion.
6
7

8 **NKD2 inhibits Wnt/ β -catenin signaling in esophageal cancer**

9

10
11 NKD2 has been reported to negatively regulate canonical Wnt
12
13 signaling in multiple tumors ^{23, 24}. To determine whether the canonical
14
15 Wnt signaling pathway is regulated by NKD2 in human esophageal
16
17 cancer, the key components in downstream of Wnt signaling pathway
18
19 were detected by western blotting. The level of β -catenin was reduced
20
21 and the level of phospho- β -catenin was increased after re-expression of
22
23 NKD2 in KYSE150 and TE1 cells. The expression of Wnt signaling
24
25 targeting genes, c-myc and cyclinD1, was reduced after re-expression of
26
27 NKD2 in KYSE150 and TE1 cells (Figure 5A). These results
28
29 demonstrate that NKD2 inhibits Wnt signaling in human esophageal
30
31 cancer. To further validate the role of NKD2 on the Wnt signaling
32
33 pathway, siRNA knockdown technique was employed. The expression of
34
35 β -catenin, c-myc and cyclinD1 was increased, and the level of
36
37 phospho- β -catenin was reduced after knockdown of NKD2 in KYSE450
38
39 cells (Figure 5B). These results suggest that NKD2 represses esophageal
40
41 cancer cell proliferation by inhibiting Wnt signaling.
42
43
44
45
46
47
48
49
50
51
52
53
54

55 **NKD2 suppresses tumor growth in esophageal cancer cell xenograft** 56 57 **mice** 58 59 60 61 62 63 64 65

To further validate the effects of NKD2 in esophageal cancer *in vivo*, NKD2 unexpressed and re-expressed KYSE150 cell xenograft mouse models were employed (Figure 6A). The volume of xenograft tumors was $345.12 \pm 18.42 \text{ mm}^3$ in NKD2 unexpressed KYSE150 cells and $96.78 \pm 17.29 \text{ mm}^3$ in NKD2 re-expressed KYSE150 cells. The tumor volume is smaller in NKD2 re-expressed KYSE150 cell xenografts compared to NKD2 unexpressed KYSE150 cell xenografts ($p < 0.001$, Figure 6A and B). The tumor weight was $220.62 \pm 28.51 \text{ mg}$ in NKD2 unexpressed KYSE150 cell xenografts and $22.35 \pm 5.19 \text{ mg}$ in NKD2 re-expressed KYSE150 cell xenografts. The tumor weight is lower in NKD2 expressed KYSE150 cell xenografts compared to NKD2 unexpressed KYSE150 cell xenografts ($p < 0.001$, Figure 6C). To further validate NKD2 inhibits Wnt signaling *in vivo*, the expression of NKD2 and the levels of phospho- β -catenin was detected by IHC staining. The levels of phospho- β -catenin were increased in NKD2 expressing esophageal cancer cell xenografts (Figure 6D). These results suggest that NKD2 suppresses esophageal cancer cell growth by inhibiting Wnt signaling *in vivo*.

Discussion

The main risk factors for ESCC are cigarette smoking and alcohol consumption³⁰. Genome-wide association analysis has demonstrated that gene-environment interaction promotes development of ESCC³¹. Other

1 studies have found that 2% of esophageal cancer and 11% of head and
2
3 neck cancer patients develop a second cancer due to field cancerization ³².
4
5
6 ³³. These studies support the idea that environment plays an important
7
8 role in ESCC. Whole-genome and whole-exome sequencing in Chinese
9
10 patients with ESCC revealed eight mutated genes, including six known
11
12 tumor-associated genes (*TP53*, *RB1*, *CDKN2A*, *PIK3CA*, *NOTCH*, and
13
14 *NFE2L2*) and two novel genes (*ADAM29* and *FAM135B*) ³⁴. Additional
15
16 genes were found frequently methylated in ESCC in previous studies ^{9, 27},
17
18
19
20
21
22 ³⁵⁻⁴⁴. Despite recent advances in treatment strategies, there has been no
23
24 significant improvement in overall survival rate for advanced and
25
26 metastatic disease ⁴⁵. New strategies are necessary for early detection and
27
28 to improve treatment options in ESCC. Aberrant epigenetic changes can
29
30 be induced by environmental factors, and epigenetic changes are
31
32 reversible under certain circumstances ^{46, 47}. Therefore, more effective
33
34 therapeutic strategies based on epigenetics are developing.
35
36
37
38
39
40
41

42 In this study, we demonstrated that NKD2 is frequently methylated
43
44 in human ESCC and the expression of NKD2 is regulated by promoter
45
46 region methylation. NKD2 methylation is associated with TNM stage and
47
48 lymph node metastasis, suggesting that NKD2 methylation may serve as
49
50 a poor prognostic marker in human ESCC. Further study found that
51
52 NKD2 inhibits esophageal cancer cell proliferation, colony formation and
53
54 induces G1/S check point arrest. In addition, NKD2 suppressed
55
56
57
58
59
60
61
62
63
64
65

1 esophageal cancer cell migration and invasion. These results suggest that
2
3 NKD2 is involved in esophageal cancer progression and metastasis. The
4
5 role of NKD2 in suppression of esophageal cancer growth was validated
6
7
8 by an esophageal cancer cell xenograft model *in vivo*. We further
9
10 explored the mechanism by which NKD2 suppresses esophageal cancer
11
12 progression and metastasis. NKD2 impedes ESCC metastasis by
13
14 down-regulating MMP2, MMP7 and MMP9 expression, and suppresses
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

At the time of diagnosis, more than 50% of esophageal cancer patients have been metastasized ⁴⁸. Though there are many approaches to treat metastatic disease, the overall survival time remains poor. Understanding the molecular events in ESCC may improve therapeutic strategies ^{49, 50}. Our finding provide more clues for epigenetic-based personalized medicine in esophageal cancer.

Conclusion

NKD2 is frequently methylated in human esophageal cancer and the expression of NKD2 is regulated by promoter region methylation. Methylation of NKD2 is associated with TNM stage and lymph node metastasis. NKD2 suppresses human esophageal cancer growth by inhibiting Wnt signaling.

List of abbreviations

5-AZA, 5-aza-2'-deoxycytidine; BSSQ, bisulfite sequencing;

GAPDH, glyceraldehyde-3-phosphate dehydrogenase; ESCC, esophageal squamous cell carcinoma; IHC, immunohistochemistry; IVD, in vitro methylated DNA; ECM gel, extracellular matrix gel; MMP, matrix metalloproteinase; MSP, methylation specific polymerase chain reaction; NKD, naked cuticle; NL, normal lymphocyte DNA; RT-PCR, reverse-transcription polymerase chain reaction; TGF α , transforming growth factor α ; TSS, transcription start sites.

Competing interests:

The authors declare no conflict of interest.

Authors' contributions

BC and WY performed experiments, analyzed data and wrote the manuscript. YJ, MZ, TH and QZ provided feedback and experimental advice. JGH and GZ provided experimental advice and manuscript editing. MG conceived the study design, supervised the experiments and edited the manuscript. All authors approved the final version of the submitted manuscript.

References

1. Ferlay J, Soerjomataram I, Dikshit R, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 2015;136:E359-386.
2. Enzinger PC, Mayer RJ. Esophageal cancer. *N Engl J Med* 2003;349:2241-2252.
3. Pennathur A, Farkas A, Krasinskas AM, et al. Esophagectomy for T1 esophageal cancer: outcomes in 100 patients and implications for endoscopic therapy. *Ann Thorac Surg* 2009;87:1048-1054; discussion 1054-1045.
4. Pennathur A, Gibson MK, Jobe BA, et al. Oesophageal carcinoma. *Lancet* 2013;381:400-412.
5. De Stefani E, Barrios E, Fierro L. Black (air-cured) and blond (flue-cured) tobacco and cancer risk. III: Oesophageal cancer. *Eur J Cancer* 1993;29A:763-766.

6. Lee CH, Wu DC, Lee JM, et al. Carcinogenetic impact of alcohol intake on squamous cell carcinoma risk of the oesophagus in relation to tobacco smoking. *Eur J Cancer* 2007;43:1188-1199.
7. Vaughan TL, Davis S, Kristal A, et al. Obesity, alcohol, and tobacco as risk factors for cancers of the esophagus and gastric cardia: adenocarcinoma versus squamous cell carcinoma. *Cancer Epidemiol Biomarkers Prev* 1995;4:85-92.
8. Ahrens TD, Werner M, Lassmann S. Epigenetics in esophageal cancers. *Cell Tissue Res* 2014;356:643-655.
9. Jia Y, Yang Y, Zhan Q, et al. Inhibition of SOX17 by microRNA 141 and methylation activates the WNT signaling pathway in esophageal cancer. *J Mol Diagn* 2012;14:577-585.
10. He G, Guan X, Chen X, et al. Expression and Splice Variant Analysis of Human TCF4 Transcription Factor in Esophageal Cancer. *J Cancer* 2015;6:333-341.
11. Argos M, Kibriya MG, Jasmine F, et al. Genomewide scan for loss of heterozygosity and chromosomal amplification in breast carcinoma using single-nucleotide polymorphism arrays. *Cancer Genet Cytogenet* 2008;182:69-74.
12. Sheu JC, Lin YW, Chou HC, et al. Loss of heterozygosity and microsatellite instability in hepatocellular carcinoma in Taiwan. *Br J Cancer* 1999;80:468-476.
13. Arias-Pulido H, Narayan G, Vargas H, et al. Mapping common deleted regions on 5p15 in cervical carcinoma and their occurrence in precancerous lesions. *Mol Cancer* 2002;1:3.
14. Xu SF, Peng ZH, Li DP, et al. Refinement of heterozygosity loss on chromosome 5p15 in sporadic colorectal cancer. *World J Gastroenterol* 2003;9:1713-1718.
15. Lu Y, Yu Y, Zhu Z, et al. Identification of a new target region by loss of heterozygosity at 5p15.33 in sporadic gastric carcinomas: genotype and phenotype related. *Cancer Lett* 2005;224:329-337.
16. Van Raay TJ, Coffey RJ, Solnica-Krezel L. Zebrafish Naked1 and Naked2 antagonize both canonical and non-canonical Wnt signaling. *Dev Biol* 2007;309:151-168.
17. Katoh M. Molecular cloning, gene structure, and expression analyses of NKD1 and NKD2. *Int J Oncol* 2001;19:963-969.
18. Yan D, Wallingford JB, Sun TQ, et al. Cell autonomous regulation of multiple Dishevelled-dependent pathways by mammalian Nkd. *Proc Natl Acad Sci U S A* 2001;98:3802-3807.
19. Rousset R, Mack JA, Wharton KA, Jr., et al. Naked cuticle targets dishevelled to antagonize Wnt signal transduction. *Genes Dev* 2001;15:658-671.
20. Zeng W, Wharton KA, Jr., Mack JA, et al. naked cuticle encodes an inducible antagonist of Wnt signalling. *Nature* 2000;403:789-795.
21. Li C, Franklin JL, Graves-Deal R, et al. Myristoylated Naked2 escorts transforming growth factor alpha to the basolateral plasma membrane of polarized epithelial cells. *Proc Natl Acad Sci U S A* 2004;101:5571-5576.
22. Hu T, Krezel AM, Li C, et al. Structural studies of human Naked2: a biologically active intrinsically unstructured protein. *Biochem Biophys Res Commun* 2006;350:911-915.
23. Zhao S, Kurenbekova L, Gao Y, et al. NKD2, a negative regulator of Wnt signaling, suppresses tumor growth and metastasis in osteosarcoma. *Oncogene* 2015.
24. Dong Y, Cao B, Zhang M, et al. Epigenetic silencing of NKD2, a major component of Wnt signaling, promotes breast cancer growth. *Oncotarget* 2015.
25. Cao B, Yang Y, Pan Y, et al. Epigenetic silencing of CXCL14 induced colorectal cancer migration and invasion. *Discov Med* 2013;16:137-147.
26. Herman JG, Graff JR, Myohanen S, et al. Methylation-specific PCR: a novel PCR assay for

methylation status of CpG islands. *Proc Natl Acad Sci U S A* 1996;93:9821-9826.

27. Jia Y, Yang Y, Brock MV, et al. Methylation of TFPI-2 is an early event of esophageal carcinogenesis. *Epigenomics* 2012;4:135-146.

28. Yan W, Wu K, Herman JG, et al. Epigenetic regulation of DACH1, a novel Wnt signaling component in colorectal cancer. *Epigenetics* 2013;8:1373-1383.

29. Yu Y, Yan W, Liu X, et al. DACT2 is frequently methylated in human gastric cancer and methylation of DACT2 activated Wnt signaling. *Am J Cancer Res* 2014;4:710-724.

30. Rustgi AK, El-Serag HB. Esophageal carcinoma. *N Engl J Med* 2014;371:2499-2509.

31. Wu C, Kraft P, Zhai K, et al. Genome-wide association analyses of esophageal squamous cell carcinoma in Chinese identify multiple susceptibility loci and gene-environment interactions. *Nat Genet* 2012;44:1090-1097.

32. Chuang SC, Hashibe M, Scelo G, et al. Risk of second primary cancer among esophageal cancer patients: a pooled analysis of 13 cancer registries. *Cancer Epidemiol Biomarkers Prev* 2008;17:1543-1549.

33. Chuang SC, Scelo G, Tonita JM, et al. Risk of second primary cancer among patients with head and neck cancers: A pooled analysis of 13 cancer registries. *Int J Cancer* 2008;123:2390-2396.

34. Song Y, Li L, Ou Y, et al. Identification of genomic alterations in oesophageal squamous cell cancer. *Nature* 2014;509:91-95.

35. Yun T, Liu Y, Gao D, et al. Methylation of CHFR sensitizes esophageal squamous cell cancer to docetaxel and paclitaxel. *Genes Cancer* 2015;6:38-48.

36. Wu L, Herman JG, Brock MV, et al. Silencing DACH1 promotes esophageal cancer growth by inhibiting TGF-beta signaling. *PLoS One* 2014;9:e95509.

37. Lu D, Ma J, Zhan Q, et al. Epigenetic silencing of RASSF10 promotes tumor growth in esophageal squamous cell carcinoma. *Discov Med* 2014;17:169-178.

38. Jiang S, Linghu E, Zhan Q, et al. Methylation of ZNF331 Promotes Cell Invasion and Migration in Human Esophageal Cancer. *Curr Protein Pept Sci* 2015;16:322-328.

39. Guo M, Ren J, House MG, et al. Accumulation of promoter methylation suggests epigenetic progression in squamous cell carcinoma of the esophagus. *Clin Cancer Res* 2006;12:4515-4522.

40. Guo M, Ren J, Brock MV, et al. Promoter methylation of HIN-1 in the progression to esophageal squamous cancer. *Epigenetics* 2008;3:336-341.

41. Guo M, House MG, Suzuki H, et al. Epigenetic silencing of CDX2 is a feature of squamous esophageal cancer. *Int J Cancer* 2007;121:1219-1226.

42. Guo M, House MG, Akiyama Y, et al. Hypermethylation of the GATA gene family in esophageal cancer. *Int J Cancer* 2006;119:2078-2083.

43. Chen XY, He QY, Guo MZ. XAF1 is frequently methylated in human esophageal cancer. *World J Gastroenterol* 2012;18:2844-2849.

44. Brock MV, Gou M, Akiyama Y, et al. Prognostic importance of promoter hypermethylation of multiple genes in esophageal adenocarcinoma. *Clin Cancer Res* 2003;9:2912-2919.

45. Mohamed A, El-Rayes B, Khuri FR, et al. Targeted therapies in metastatic esophageal cancer: advances over the past decade. *Crit Rev Oncol Hematol* 2014;91:186-196.

46. Herman JG, Baylin SB. Gene silencing in cancer in association with promoter hypermethylation. *N Engl J Med* 2003;349:2042-2054.

47. Baylin SB, Esteller M, Rountree MR, et al. Aberrant patterns of DNA methylation, chromatin formation and gene expression in cancer. *Hum Mol Genet* 2001;10:687-692.

- 1 48. Wu AH, Wan P, Bernstein L. A multiethnic population-based study of smoking, alcohol and body
2 size and risk of adenocarcinomas of the stomach and esophagus (United States). *Cancer Causes*
3 *Control* 2001;12:721-732.
- 4 49. Guo M, Liu S, Herman JG, et al. Gefitinib-sensitizing mutation in esophageal carcinoma cell line
5 Kyse450. *Cancer Biol Ther* 2006;5:152-155.
- 6 50. Guo M, Liu S, Lu F. Gefitinib-sensitizing mutations in esophageal carcinoma. *N Engl J Med*
7 2006;354:2193-2194.
- 8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1. Clinical factors and NKD2 methylation in 154 cases of esophageal cancer

Clinical factor	No.	NKD2 methylation status		<i>p</i> * value
		Methylated n=82(53.2%)	Unmethylated n=72 (46.8%)	
<i>Age (year)</i>				
<50	10	5	5	<i>p</i> =0.9085
≥50	144	77	67	
<i>Gender</i>				
Male	116	65	51	<i>p</i> =0.3058
Female	38	17	21	
<i>Tumor Size (cm)</i>				
<5	99	50	49	<i>p</i> =0.4555
≥5	55	32	23	
<i>Differentiation</i>				
Well	9	6	3	<i>p</i> =0.3792
Moderate	97	54	43	
Poor	48	22	26	
<i>TNM Stage</i>				
I + II	104	47	57	<i>p</i> =0.0066<0.01
III+IV	50	35	15	
<i>Lymph node</i>				
<i>Metastasis</i>				
N0	90	39	51	<i>p</i> =0.0058<0.01
N1	64	43	21	

* *p* values are obtained from chi-square test and the Fisher's exact test, significant difference, *p*< 0.05

Figure legends

Table 1. Clinical factors and NKD2 methylation in 154 cases of esophageal cancer

Figure 1. The expression and methylation status of NKD2 in esophageal cancer cells and normal esophageal mucosa

A. Semi-quantitative RT-PCR shows NKD2 expression levels in esophageal cancer cell lines. KYSE450, KYSE30, KYSE150, KESE70, TE8, KYSE410, TE1, KYSE140 and COLO680 are esophageal cancer cell lines. 5-AZA: 5-aza-2'-deoxycytidine; GAPDH: internal control of RT-PCR; H₂O: double distilled water. (-): absence of 5-AZA; (+): presence of 5-AZA.

B. MSP results of NKD2 in esophageal cancer cell lines. U: unmethylated alleles; M: methylated alleles; IVD: *in vitro* methylated DNA, serves as methylation control; NL: normal peripheral lymphocytes DNA, serves as unmethylation control; H₂O: double distilled water.

C. BSSQ results of NKD2. KYSE150, KYSE410, TE1, KYSE450: esophageal cancer cells, NE: normal esophageal mucosa. Double-headed arrow: MSP PCR product spanned 103 bp in NKD2. Bisulfite sequencing focused on a 287 bp region of the CpG island

(-287 bp to +38 bp) across the NKD2 transcription start site. Filled circles: methylated CpG sites, open circles: unmethylated CpG sites. TSS: transcription start site.

Figure 2. Methylation status and expression of NKD2 in primary esophageal cancer samples

A. MSP results of NKD2 in normal esophageal mucosa. NE: normal esophageal mucosa.

B. Representative results of MSP for NKD2 in primary esophageal cancer samples. EC: primary esophageal cancer samples.

C. Representative IHC results showing NKD2 expression in esophageal cancer and matched adjacent tissue samples (upper: $\times 100$; lower: $\times 400$).

D. NKD2 expression scores are shown as box plots, horizontal lines represent the median score; the bottom and top of the boxes represent the 25th and 75th percentiles, respectively; vertical bars represent the range of data. The expression level of NKD2 was significantly different between adjacent tissue and esophageal cancer samples. $***p < 0.001$.

E. The bar diagram shows the expression and DNA methylation status of NKD2 in different cancer samples. Reduced expression of NKD2 was significantly associated with promoter region methylation .

**** $p<0.01$.**

Figure 3. NKD2 inhibits esophageal cancer cell proliferation.

A. Growth curves represent the cell viability analyzed by the MTT assay in NKD2 re-expressed and unexpressed KYSE150 and TE1 cells. The experiment was performed in triplicate. $*p<0.05$, $***p<0.001$.

B. Colony formation results show that colony number was reduced by re-expression of NKD2 in KYSE150 and TE1 cells. Each experiment was repeated three times. The average number of tumor clones is represented by bar diagram. $***p<0.001$.

C. Cell phase distribution in NKD2 unexpressed and re-expressed KYSE150 and TE1 cells. The ratio is presented by bar diagram. Each experiment was repeated three times. $**p<0.01$, $***p<0.001$.

Figure 4. Restoration of NKD2 expression inhibits cell migration and invasion

A. Cell migration in NKD2 unexpressed and re-expressed KYSE150 and TE1 cells. The ratio is presented by bar diagram. Each experiment was repeated three times. $***p<0.001$.

B. Cell invasion in NKD2 unexpressed and re-expressed KYSE150 and TE1 cells. The ratio is presented by bar diagram. Each experiment was repeated three times. $***p<0.001$.

1 C. The expression levels of NKD2, MMP-2, MMP-7 and MMP-9 were
2
3 detected by western blot in NKD2 unexpressed and re-expressed
4
5 KYSE150 and TE1 cells. Knockdown of NKD2 by siRNA was
6
7 performed to validate the results in NKD2 highly expressed
8
9 KYSE450 cells.
10
11
12
13
14

15 **Figure 5. NKD2 inhibits canonical Wnt signaling in human**
16
17 **esophageal cancer cells.**
18
19
20

21 A. The expression levels of β -catenin, cyclin D1 and c-myc were reduced
22
23 and the level of phosphorylated β -catenin (p- β -catenin) increased after
24
25 re-expression of NKD2 in KYSE150 and TE1 cells.
26
27
28
29

30 B. The level of p- β -catenin was reduced and the expression of β -catenin,
31
32 c-myc and cyclin D1 were increased after knockdown of NKD2 by
33
34 siRNA in KYSE450 cells
35
36
37
38

39 **Figure 6. NKD2 suppresses esophageal cancer cell growth in**
40
41 **xenograft mice**
42
43
44

45 A. Representative burdened nude mice in NKD2 re-expressed and
46
47 unexpressed KYSE150 cells.
48
49
50

51 B. Subcutaneous tumor growth curves for xenograft mice in NKD2
52
53 unexpressed and re-expressed groups at different times. *** $p < 0.001$.
54
55
56
57

58 C. Tumor weight in nude mice at the 24th day after inoculation of NKD2
59
60
61
62
63
64
65

unexpressed and re-expressed KYSE150 cells. Bars: mean of 6 mice.

*** $p<0.001$.

D. Representative photographs of IHC analysis of NKD2 and p- β -catenin in xenografts. Staining of NKD2 and p- β -catenin was found in NKD2 re-expressed KYSE150 cell xenografts. Magnification: 400 \times .

Figure 1




Figure 2






Figure 3

A



B



C




Figure 4





Figure 5

A



B




Figure 6

