

Problem Solving Guide: Complex 1D kinematics w/ constant accel. Intervals

The Model

Analytical problems involving one dimensional motion are problems where the details of the motion are described by equations. Assumptions are:

- Objects are treated as point particles, with only a single, definite position.
- The motions of the object or objects have constant acceleration over given intervals. Acceleration may change, however, from interval to interval. If this occurs, we assume such a change is instantaneous.
- The motion is strictly along one straight line.

Problem Solving Steps

- 1) Organize and plan
 - a) Use diagrams, graphs, or other tools to visualize the situation. Label key instants in time clearly.
 - b) Define all variables clearly, using subscripts to denote the time and the object referred to. Note down known variables.
 - c) Identify what strategies will solve the problem.
 - d) Identify relationships between variables at key times or positions.
- 2) Solve
 - a) Apply the constant accel. equations to intervals of constant accel. in order to solve for unknowns. This often involves repeated application of these equations.
- 3) Reflect
 - a) Do your answers make sense?
 - b) Are the units correct?
 - c) Did you use any new techniques?
 - d) Any other insights?

Constant Acceleration Motion Equations

$$v_B = v_A + a_{AB}\Delta t_{AB}$$

$$v_B^2 = v_A^2 + 2a_{AB}\Delta x_{AB}$$

$$x_B = x_A + v_A\Delta t_{AB} + \frac{a_{AB}}{2}\Delta t_{AB}^2$$

$$x_B = x_A + \left(\frac{v_A + v_B}{2}\right)\Delta t_{AB}$$