


Implicit Differentiation

A graph can be expressed as an explicit function of x or an implicit function of x :

Explicitly in terms of x	Implicitly in terms of x
y is written explicitly in terms of x $y = \sqrt{x^2 + 1}$ $y = x^2 + 2x + 1$ \vdots	$y = \frac{1}{x}$ can be written implicitly in terms of x as $xy = 1$ The function $y = f(x)$ is implied by the equation. $xy = 1$ $x^2 + y^2 = 25$ \vdots

It is oftentimes difficult to define y in terms of x explicitly. To find the derivative of such a relation, we must use **implicit differentiation**.



In implicit differentiation, we differentiate with respect to x , however we treat each y as $f(x)$, and use chain rule on each $y \leftrightarrow f(x)$.

A few things to remember when executing implicit differentiation:

- ✓ Remember that $y = f(x)$. You have to apply chain rule when differentiating y .

$f(x)$	y	$\cos(f(x))$	$\cos(y)$	$[f(x)]^5$	y^5
\downarrow	\downarrow	\downarrow	\downarrow	\downarrow	\downarrow
$f'(x)$	y'	$-\sin(f(x)) \cdot f'(x)$	$-\sin(y) \cdot y'$	$5[f(x)]^4 \cdot f'(x)$	$5y^4 \cdot y'$

- Differentiate a component involving y just like x , however, you multiply the result of that component by y'
- ✓ If an “ x ” is multiplying a “ y ” then product rule must be used.
 $xy = 3y^2 \cos(x)$
 \downarrow
 $1 \cdot y + x \cdot y' = 6y \cdot y' \cos(x) + 3y^2 [-\sin(x)]$
- ✓ When solving for y' a.k.a. $\frac{dy}{dx}$, or y'' a.k.a. $\frac{d^2y}{dx^2}$ you must solve in terms of x and y only.
 - When finding y'' , if $y' = [\dots]$, then replace every y' with $[\dots]$ in the expression for y''

When solving for y'

- I. Differentiate, and after differentiating distribute wherever possible.
- II. Move all terms with y' to one side, and all terms without y' to the other.
- III. Factor out the common factor of y'
- IV. Divide both sides by the factor of y'

Find y' in terms of x and y only: $\cos(xy) = \sin(y) + 1$

Then find $y'|_{(0,0)}$.

Find $\frac{d^2y}{dx^2}$ in terms of x and y only: $x^2 + xy + y^2 = 1$