

What Is the Second Difference Matrix?

Nicholas J. Higham*

January 31, 2022

The second difference matrix is the tridiagonal matrix T_n with diagonal elements 2 and sub- and superdiagonal elements -1 :

$$T_n = \begin{bmatrix} 2 & -1 & & & \\ -1 & 2 & -1 & & \\ & -1 & 2 & \ddots & \\ & & \ddots & \ddots & -1 \\ & & & -1 & 2 \end{bmatrix} \in \mathbb{R}^{n \times n}.$$

It arises when a second derivative is approximated by the central second difference $f''(x) \approx (f(x+h) - 2f(x) + f(x-h))/h^2$. (Accordingly, the second difference matrix is sometimes defined as $-T_n$.) In MATLAB, T_n can be generated by `gallery('tridiag',n)`, which is returned as a sparse matrix.

This is Gil Strang's favorite matrix. The photo, from his home page, shows a birthday cake representation of the matrix.



The second difference matrix is symmetric positive definite. The easiest way to see this is to define the full rank rectangular matrix

$$L_n = \begin{bmatrix} 1 & & & & \\ -1 & 1 & & & \\ & -1 & \ddots & & \\ & & \ddots & 1 & \\ & & & -1 & 1 \end{bmatrix} \in \mathbb{R}^{(n+1) \times n}$$

*Department of Mathematics, University of Manchester, Manchester, M13 9PL, UK (nick.higham@manchester.ac.uk).

and note that $T_n = L_n^T L_n$. The factorization corresponds to representing a central difference as the product of a forward difference and a backward difference. Other properties of the second difference matrix are that it is diagonally dominant, a Toeplitz matrix, and an M -matrix.

Cholesky Factorization

In an LU factorization $A = LU$ the pivots u_{ii} are $2, 3/2, 4/3, \dots, (n+1)/n$. Hence the pivots form a decreasing sequence tending to 1 as $n \rightarrow \infty$. The diagonal of the Cholesky factor contains the square roots of the pivots. This means that in the Cholesky factorization $A = R^*R$ with R upper bidiagonal, $r_{nn} \rightarrow 1$ and $r_{n,n-1} \rightarrow -1$ as $n \rightarrow \infty$.

Determinant, Inverse, Condition Number

Since the determinant is the product of the pivots, $\det(T_n) = n+1$.

The inverse of T_n is full, with (i, j) element $i(n-j+1)/(n+1)$ for $j \geq i$. For example,

$$T_5^{-1} = \frac{1}{6} \begin{bmatrix} 5 & 4 & 3 & 2 & 1 \\ 4 & 8 & 6 & 4 & 2 \\ 3 & 6 & 9 & 6 & 3 \\ 2 & 4 & 6 & 8 & 4 \\ 1 & 2 & 3 & 4 & 5 \end{bmatrix}.$$

The 2-norm condition number satisfies $\kappa_2(T_n) \sim 4n^2/\pi^2$ (as follows from the formula (1) below for the eigenvalues).

Eigenvalues and Eigenvectors

The eigenvalues of T_n are

$$\mu_k = 2 - 2\cos(k\phi) = 4\sin^2\left(\frac{k\phi}{2}\right), \quad k = 1 : n, \quad (1)$$

where $\phi = \pi/(n+1)$, with corresponding eigenvector

$$v_k = [\sin(k\phi), \sin(2k\phi), \dots, \sin(nk\phi)]^T.$$

The matrix Q with

$$q_{ij} = \left(\frac{2}{n+1}\right)^{1/2} \sin\left(\frac{ij\pi}{n+1}\right)$$

is therefore an eigenvector matrix for T_n : $Q^* A Q = \text{diag}(\mu_k)$.

Variations

Various modifications of the second difference matrix arise and similar results can be derived. For example, consider the matrix obtained by changing the (n, n) element to 1:

$$\tilde{T}_n = \begin{bmatrix} 2 & -1 & & & \\ -1 & 2 & -1 & & \\ & -1 & 2 & \ddots & \\ & & \ddots & \ddots & -1 \\ & & & -1 & 1 \end{bmatrix} \in \mathbb{R}^{n \times n}.$$

It can be shown that \tilde{T}_n^{-1} has (i, j) element $\min(i, j)$ and eigenvalues $4 \cos(j\pi/(2n+1))^2$, $j = 1 : n$.

If we perturb the $(1, 1)$ of \tilde{T}_n to 1, we obtain a singular matrix, but suppose we perturb the $(1, 1)$ element to 3:

$$\hat{T}_n = \begin{bmatrix} 3 & -1 & & & \\ -1 & 2 & -1 & & \\ & -1 & 2 & \ddots & \\ & & \ddots & \ddots & -1 \\ & & & -1 & 1 \end{bmatrix} \in \mathbb{R}^{n \times n}.$$

The inverse is $\hat{T}_n^{-1} = G/2$, where G with (i, j) element $2 \min(i, j) - 1$ is a matrix of Givens, and the eigenvalues are $4 \cos((2j-1)\pi/(4n))^2$, $j = 1 : n$.

Notes

The factorization $T_n = L_n^T L_n$ is noted by Strang (2012).

For a derivation of the eigenvalues and eigenvectors see Todd (1977, p. 155 ff.). For the eigenvalues of \tilde{T}_n see Fortiana and Cuadras (1997). Givens's matrix is described by Newman and Todd (1958) and Todd (1977).

References

This is a minimal set of references, which contain further useful references within.

- J. Fortiana and C. N. Cuadras, A Family of Matrices, the Discretized Brownian Bridge, and Distance-Based Regression, *Linear Algebra Appl.* 264, 173–188, 1997.
- Morris Newman and John Todd, The Evaluation of Matrix Inversion Programs, *J. Soc. Indust. Appl. Math.* 6(4), 466–476, 1958.
- Gilbert Strang, *Essays in Linear Algebra*, Wellesley-Cambridge Press, Wellesley, MA, USA, 2012. Chapter A.3: “My Favorite Matrix”.
- John Todd, *Basic Numerical Mathematics*, Vol. 2: Numerical Algebra, Birkhäuser, Basel, and Academic Press, New York, 1977.

Related Blog Posts

- What Is a Diagonally Dominant Matrix? (2021)
- What Is a Sparse Matrix? (2020)
- What Is a Symmetric Positive Definite Matrix? (2020)
- What Is a Tridiagonal Matrix? (2022)
- What Is an M-Matrix? (2021)

This article is part of the “What Is” series, available from <https://nhigham.com/index-of-what-is-articles/> and in PDF form from the GitHub repository <https://github.com/higham/what-is>.