# Chapter 1.5 Elementary Matrices

IRISH C. SIDAYA

**Recall:** Three elementary row operations for matrices

- Interchange two rows.
- 2 Multiply a row by a nonzero constant.
- Add a multiple of a row to another row.

Recall: Three elementary row operations for matrices

- Interchange two rows.
- 2 Multiply a row by a nonzero constant.
- 3 Add a multiple of a row to another row.

### Definition

An  $n \times n$  matrix is called an **elementary matrix** when it can be obtained from the identity matrix  $I_n$  by a single elementary row operation.

**Recall:** Three elementary row operations for matrices

- Interchange two rows.
- Multiply a row by a nonzero constant.
- 3 Add a multiple of a row to another row.

### Definition

An  $n \times n$  matrix is called an **elementary matrix** when it can be obtained from the identity matrix  $I_n$  by a single elementary row operation.

**Example:** Which of the following matrices are elementary? For those that are, describe the corresponding elementary row operation.

**a.** 
$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

**a.** 
$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$
 **b.** 
$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$
 **c.** 
$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$
 **d.** 
$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

$$\mathbf{d.} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

e. 
$$\begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}$$

**e.** 
$$\begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}$$
 **f.**  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -1 \end{bmatrix}$ 

### **SOLUTION**

- **a.** This matrix is elementary. To obtain it from  $I_3$ , multiply the second row of  $I_3$  by 3.
- **b.** This matrix is *not* elementary because it is not square.
- **c.** This matrix is *not* elementary because to obtain it from I<sub>3</sub>, you must multiply the third row of  $I_3$  by 0 (row multiplication must be by a nonzero constant).
- **d.** This matrix is elementary. To obtain it from  $I_3$ , interchange the second and third rows of  $I_3$ .
- e. This matrix is elementary. To obtain it from  $I_2$ , multiply the first row of  $I_2$  by 2 and add the result to the second row.
- **f.** This matrix is *not* elementary because it requires two elementary row operations to obtain from  $I_3$ .

Sidava, Irish C. 3 / 13

### Theorem

### Representing Elementary Row Operations

Let E be the elementary matrix obtained by performing an elementary row operation on  $I_m$ . If that same elementary row operation is performed on an  $m \times n$  matrix A, then the resulting matrix is given by the product EA.

### Theorem

### Representing Elementary Row Operations

Let E be the elementary matrix obtained by performing an elementary row operation on  $I_m$ . If that same elementary row operation is performed on an  $m \times n$  matrix A, then the resulting matrix is given by the product EA.

### Example:

In the matrix product below, E is the elementary matrix in which the first two rows of  $I_3$  are interchanged.

$$\begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 2 & 1 \\ 1 & -3 & 6 \\ 3 & 2 & -1 \end{bmatrix} = \begin{bmatrix} 1 & -3 & 6 \\ 0 & 2 & 1 \\ 3 & 2 & -1 \end{bmatrix}$$

### Theorem

### Representing Elementary Row Operations

Let E be the elementary matrix obtained by performing an elementary row operation on  $I_m$ . If that same elementary row operation is performed on an  $m \times n$  matrix A, then the resulting matrix is given by the product EA.

### Example:

In the matrix product below, E is the elementary matrix in which the first two rows of  $I_3$  are interchanged.

$$\begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 2 & 1 \\ 1 & -3 & 6 \\ 3 & 2 & -1 \end{bmatrix} = \begin{bmatrix} 1 & -3 & 6 \\ 0 & 2 & 1 \\ 3 & 2 & -1 \end{bmatrix}$$

Note that the first two rows of A are interchanged by multiplying on the left by E.

Sidava, Irish C. 4/1

### Definition

### Row Equivalence

Let A and B be  $m \times n$  matrices. Matrix B is **row-equivalent** to A when there exists a finite number of elementary matrices  $E_1, E_2, \ldots, E_k$  such that

$$B = E_k E_{k-1} \cdot \cdot \cdot E_2 E_1 A.$$

### Definition

Row Equivalence

Let A and B be  $m \times n$  matrices. Matrix B is **row-equivalent** to A when there exists a finite number of elementary matrices  $E_1, E_2, \ldots, E_k$  such that

$$B = E_k E_{k-1} \cdot \cdot \cdot E_2 E_1 A.$$

### Example:

Find a sequence of elementary matrices that can be used to write the matrix A in row-echelon form.

$$A = \begin{bmatrix} 0 & 1 & 3 & 5 \\ 1 & -3 & 0 & 2 \\ 2 & -6 & 2 & 0 \end{bmatrix}$$

### Theorem

Elementary Matrices Are Invertible

If E is an elementary matrix, then  $E^{-1}$  exists and is an elementary matrix.

### Theorem

Elementary Matrices Are Invertible

If E is an elementary matrix, then  $E^{-1}$  exists and is an elementary matrix.

**Note:** The inverse of an elementary matrix E is the elementary matrix  $E^{-1}$  that converts back to  $I_n$ .

### Theorem

Elementary Matrices Are Invertible

If E is an elementary matrix, then  $E^{-1}$  exists and is an elementary matrix.

**Note:** The inverse of an elementary matrix E is the elementary matrix  $E^{-1}$  that converts back to  $I_n$ .

**Examples:** Find the inverses of the following elementary matrices

$$E_1 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

### Theorem

Elementary Matrices Are Invertible

If E is an elementary matrix, then  $E^{-1}$  exists and is an elementary matrix.

**Note:** The inverse of an elementary matrix E is the elementary matrix  $E^{-1}$  that converts back to  $I_n$ .

**Examples:** Find the inverses of the following elementary matrices

$$E_1 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix}$$

**◆□▶◆□▶◆臺▶◆臺▶ 臺 か**900

### Theorem

Elementary Matrices Are Invertible

If E is an elementary matrix, then  $E^{-1}$  exists and is an elementary matrix.

**Note:** The inverse of an elementary matrix E is the elementary matrix  $E^{-1}$  that converts back to  $I_n$ .

Examples: Find the inverses of the following elementary matrices

$$E_1 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix} \quad E_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \frac{1}{2} \end{bmatrix}$$

### Theorem

A Property of Invertible Matrices

A square matrix *A* is invertible if and only if it can be written as the product of elementary matrices.

### Theorem

A Property of Invertible Matrices

A square matrix A is invertible if and only if it can be written as the product of elementary matrices.

### Example:

Find a sequence of elementary matrices whose product is the nonsingular matrix

$$A = \begin{bmatrix} -1 & -2 \\ 3 & 8 \end{bmatrix}.$$

### Theorem

### Equivalent Conditions

If A is an  $n \times n$  matrix, then the following statements are equivalent.

- **1.** A is invertible.
- **2.**  $A\mathbf{x} = \mathbf{b}$  has a unique solution for every  $n \times 1$  column matrix  $\mathbf{b}$ .
- 3.  $A\mathbf{x} = O$  has only the trivial solution.
- **4.** A is row-equivalent to  $I_n$ .
- **5.** A can be written as the product of elementary matrices.

### Definitions

• A square matrix is **lower triangular** if all the entries above its main diagonal are zero.

### Definitions

- A square matrix is **lower triangular** if all the entries above its main diagonal are zero.
- A square matrix is upper triangular if all the entries below its main diagonal are zero.

Sidaya, Irish C. 9 / 13

### **Definitions**

- A square matrix is **lower triangular** if all the entries above its main diagonal are zero.
- 2 A square matrix is **upper triangular** if all the entries below its main diagonal are zero.

$$\begin{bmatrix} a_{11} & 0 & 0 \\ a_{21} & a_{22} & 0 \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

 $3 \times 3$  lower triangular matrix  $3 \times 3$  upper triangular matrix

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{bmatrix}$$

### Definitions

- A square matrix is **lower triangular** if all the entries above its main diagonal are zero.
- 2 A square matrix is **upper triangular** if all the entries below its main diagonal are zero.

$$\begin{bmatrix} a_{11} & 0 & 0 \\ a_{21} & a_{22} & 0 \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

 $3 \times 3$  lower triangular matrix  $3 \times 3$  upper triangular matrix

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{bmatrix}$$

### Definition

If the  $n \times n$  matrix A can be written as the product of a lower triangular matrix L and an upper triangular matrix U, then A = LU is an LU-factorization of A.

### Examples:

$$\mathbf{a.} \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & -2 \end{bmatrix} = LU$$

is an LU-factorization of the matrix

$$A = \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}$$

### Examples:

**a.** 
$$\begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & -2 \end{bmatrix} = LU$$

is an LU-factorization of the matrix

$$A = \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}$$

**b.** 
$$A = \begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 2 & -10 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & -4 & 1 \end{bmatrix} \begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 0 & 0 & 14 \end{bmatrix} = LU$$

is an LU-factorization of the matrix A.

### Example:

Find an *LU*-factorization of the matrix 
$$A = \begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 2 & -10 & 2 \end{bmatrix}$$
.

### Example:

Find an *LU*-factorization of the matrix 
$$A = \begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 2 & -10 & 2 \end{bmatrix}$$
.

### SOLUTION

Begin by row reducing A to upper triangular form while keeping track of the elementary matrices used for each row operation.

### Example:

Find an *LU*-factorization of the matrix 
$$A = \begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 2 & -10 & 2 \end{bmatrix}$$
.

### SOLUTION

Begin by row reducing A to upper triangular form while keeping track of the elementary matrices used for each row operation.

Matrix

### Example:

Find an *LU*-factorization of the matrix 
$$A = \begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 2 & -10 & 2 \end{bmatrix}$$
.

### SOLUTION

Begin by row reducing A to upper triangular form while keeping track of the elementary matrices used for each row operation.

Matrix Elementary Row Operation

### Example:

Find an *LU*-factorization of the matrix 
$$A = \begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 2 & -10 & 2 \end{bmatrix}$$
.

### SOLUTION

Begin by row reducing A to upper triangular form while keeping track of the elementary matrices used for each row operation.

Matrix

Elementary Row Operation

### Example:

Find an *LU*-factorization of the matrix 
$$A = \begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 2 & -10 & 2 \end{bmatrix}$$
.

### SOLUTION

Begin by row reducing A to upper triangular form while keeping track of the elementary matrices used for each row operation.

Matrix Elementary Row Operation

$$\begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 0 & -4 & 2 \end{bmatrix}$$

### Example:

Find an *LU*-factorization of the matrix 
$$A = \begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 2 & -10 & 2 \end{bmatrix}$$
.

### SOLUTION

Begin by row reducing A to upper triangular form while keeping track of the elementary matrices used for each row operation.

Matrix Elementary Row Operation Elementary Matrix

$$\begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 0 & -4 & 2 \end{bmatrix} \qquad R_3 + (-2)R_1 \to R_3$$

$$R_3 + (-2)R_1 \rightarrow R_3$$

### Example:

Find an *LU*-factorization of the matrix 
$$A = \begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 2 & -10 & 2 \end{bmatrix}$$
.

### SOLUTION

Begin by row reducing A to upper triangular form while keeping track of the elementary matrices used for each row operation.

Matrix Elementary Row Operation

$$\begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 0 & -4 & 2 \end{bmatrix} \qquad R_3 + (-2)R_1 \to R_3$$

$$E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix}$$

### Example:

Find an *LU*-factorization of the matrix 
$$A = \begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 2 & -10 & 2 \end{bmatrix}$$
.

### SOLUTION

Begin by row reducing A to upper triangular form while keeping track of the elementary matrices used for each row operation.

### Matrix

### Elementary Row Operation

$$\begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 0 & -4 & 2 \end{bmatrix} \qquad R_3 + (-2)R_1 \to R_3$$

$$R_3 + (-2)R_1 \to R$$

$$E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 0 & 0 & 14 \end{bmatrix}$$

### Example:

Find an *LU*-factorization of the matrix 
$$A = \begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 2 & -10 & 2 \end{bmatrix}$$
.

### SOLUTION

Begin by row reducing A to upper triangular form while keeping track of the elementary matrices used for each row operation.

# Matrix $\begin{vmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 0 & -4 & 2 \end{vmatrix}$ $R_3 + (-2)R_1 \to R_3$

$$R_3 + (-2)R_1 \rightarrow R_3$$

$$R_3 + (-2)R_1 \rightarrow R_3$$

$$\begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 0 & 0 & 14 \end{bmatrix}$$

$$R_3 + (4)R_2 \rightarrow R_3$$

$$E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix}$$

### Example:

Find an *LU*-factorization of the matrix 
$$A = \begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 2 & -10 & 2 \end{bmatrix}$$
.

### SOLUTION

Begin by row reducing A to upper triangular form while keeping track of the elementary matrices used for each row operation.

# Matrix Elementary Row Operation $\begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 0 & -4 & 2 \end{bmatrix}$ $R_3 + (-2)R_1 \rightarrow R_3$ $\begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 0 & 0 & 14 \end{bmatrix}$ $R_3 + (4)R_2 \rightarrow R_3$

$$E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix}$$

$$E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 4 & 1 \end{bmatrix}$$

### Example:

Find an *LU*-factorization of the matrix 
$$A = \begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 2 & -10 & 2 \end{bmatrix}$$
.

### SOLUTION

Begin by row reducing A to upper triangular form while keeping track of the elementary matrices used for each row operation.

Matrix Elementary Row Operation Elementary Matrix  $\begin{vmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 0 & -4 & 2 \end{vmatrix}$   $R_3 + (-2)R_1 \to R_3$  $E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix}$  $\begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ \vdots & \vdots & 14 \end{bmatrix}$  $E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 4 & 1 \end{bmatrix}$ 

Observe that the above matrix is upper triangular. We denote it by U so that we have  $E_2E_1A=U$ . ◆□ → ◆□ → ◆重 → ◆■ ・ ◆ ● ● ・ ◆ ● ・ ◆ ● ● ・ ◆ ● ・ ◆ ● ・ ◆ ● ・ ◆ ● ・ ◆ ● ・ ◆ ● ● ・ ◆ ● ・ ◆ ● ・ ◆ ● ・ ◆ ● ・ ◆ ●

### Example:

Find an *LU*-factorization of the matrix 
$$A = \begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 2 & -10 & 2 \end{bmatrix}$$
.

### **SOLUTION**

Begin by row reducing *A* to upper triangular form while keeping track of the elementary matrices used for each row operation.

# Matrix Elementary Row Operation Elementary Matrix $\begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 0 & -4 & 2 \end{bmatrix} \qquad R_3 + (-2)R_1 \rightarrow R_3 \qquad E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix}$ $\begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 3 \\ 0 & 0 & 14 \end{bmatrix} \qquad E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 4 & 1 \end{bmatrix}$

Observe that the above matrix is upper triangular. We denote it by U so that we have  $E_2E_1A=U$ . This implies that  $A=E_1^{-1}E_2^{-1}U$ .

Moreover, notice that the product of lower triangular matrices

$$E_1^{-1}E_2^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -4 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & -4 & 1 \end{bmatrix}$$

is a lower triangular matrix, denote it by L. Hence the factorization A = LU is complete.

Moreover, notice that the product of lower triangular matrices

$$E_1^{-1}E_2^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -4 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & -4 & 1 \end{bmatrix}$$

is a lower triangular matrix, denote it by L. Hence the factorization A = LU is complete.

### Remark:

If A row reduces to an upper triangular matrix U using only the row operation of adding a multiple of one row to another row below it, then A has an LU-factorization.

$$E_k \cdot \cdot \cdot E_2 E_1 A = U$$
  
 $A = E_1^{-1} E_2^{-1} \cdot \cdot \cdot E_k^{-1} U = LU$ 

Here L is the product of the inverses of the elementary matrices used in the row reduction.

# Solving a Linear System Using LU-Factorization

### Remark:

Once you have obtained an LU-factorization of a matrix A, you can then solve the system of n linear equations in n variables  $A\mathbf{x} = \mathbf{b}$  very efficiently in two steps.

- **1.** Write  $\mathbf{y} = U\mathbf{x}$  and solve  $L\mathbf{y} = \mathbf{b}$  for  $\mathbf{y}$ .
- 2. Solve  $U\mathbf{x} = \mathbf{y}$  for  $\mathbf{x}$ .

# Solving a Linear System Using LU-Factorization

### Remark:

Once you have obtained an LU-factorization of a matrix A, you can then solve the system of n linear equations in n variables  $A\mathbf{x} = \mathbf{b}$  very efficiently in two steps.

- 1. Write  $\mathbf{y} = U\mathbf{x}$  and solve  $L\mathbf{y} = \mathbf{b}$  for  $\mathbf{y}$ .
- 2. Solve  $U\mathbf{x} = \mathbf{y}$  for  $\mathbf{x}$ .

### Example:

Solve the linear system.

$$x_1 - 3x_2 = -5$$
  
 $x_2 + 3x_3 = -1$   
 $2x_1 - 10x_2 + 2x_3 = -20$