

MSCS 446 Numerical Analysis I
Written Assignment 8
Adhere to the Homework Guidelines
 Dr. Keith Wojciechowski

1. (N) Compute

$$\int_0^{2\pi} \frac{\cos(2x)}{e^x} dx \quad \text{the exact answer is } \frac{1}{5}(1 - e^{-2\pi})$$

using $n = 120$ equispaced points for the

- (a) composite trapezoidal rule
- (b) composite Simpson's rule (remember that you need 120 subintervals!)
- (c) Gaussian Three-Point Rule (this rule is NOT Gaussian quadrature, it is the rule given below)

Which method is most accurate in terms of absolute error?

The **Gaussian Three-Point Rule** is defined

$$\int_a^b f(x) dx \approx h \sum_{i=1}^{n/2} \left[\frac{5}{9} f \left(x_{2i-1} - h\sqrt{\frac{3}{5}} \right) + \frac{8}{9} f(x_{2i-1}) + \frac{5}{9} f \left(x_{2i-1} + h\sqrt{\frac{3}{5}} \right) \right]$$

Notice that this rule only uses the odd-indexed nodes (half as many nodes as the other two methods). Compute it using a dot product.

- 2. (A) In class we derived Simpson's $\frac{1}{3}$ Rule for 3 points. Complete that calculation from start to finish.
- 3. (A) Determine the degree of precision of the quadrature rule.

$$\int_{-1}^1 f(x) dx \approx \frac{1}{4} [3f(-2/3) + 2f(0) + 3f(2/3)]$$

- 4. (A) How large must n be if the composite trapezoidal rule is being used to estimate $\int_0^\pi \sin(x) dx$ with error $\leq 10^{-12}$? Will the estimate be too big or too small (justify using Calculus)? Repeat this exercise using the composite Simpson's rule.
- 5. (A) Construct a rule of the form (determine the α , β , and γ)

$$\int_{-1}^1 f(x) dx \approx \alpha f \left(-\frac{1}{2} \right) + \beta f(0) + \gamma f \left(\frac{1}{2} \right)$$

that has degree of precision 2.