

# 11.16.2.2.7

EE24BTECH11058 - P.Shiny Diavajna

## Question:

A die is thrown

A: The event that a number less than 7 appears.

B: The event that a number greater than 7 appears.

Find  $P(A + B)$

## Solution:

For 2 Boolean variables  $A$  and  $B$ , the axioms of Boolean Algebra are defined as:

$$A + A' = 1 \quad (0.1)$$

$$A + A = A \quad (0.2)$$

$$AB = BA \quad (0.3)$$

$$A + B = B + A \quad (0.4)$$

$$AA' = 0 \quad (0.5)$$

$$\Pr(1) = 1 \quad (0.6)$$

$$\Pr(A + B) = \Pr(A) + \Pr(B), \text{ if } \Pr(AB) = 0 \quad (0.7)$$

Using these axioms, we will try to prove that

$$\Pr(A + B) = \Pr(A) + \Pr(B) - \Pr(AB) \quad (0.8)$$

We will start by representing  $A$  and  $B$  as:

$$A = AB + AB' \quad (0.9)$$

$$B = AB + A'B \quad (0.10)$$

$$\Pr(A) = \Pr(AB) + \Pr(AB') \quad (0.11)$$

$$\Pr(B) = \Pr(AB) + \Pr(A'B) \quad (0.12)$$

On adding (0.9) and (0.10),

$$A + B = AB + AB + AB' + A'B \quad (0.13)$$

$$A + B = AB + AB' + A'B \quad (0.14)$$

$$\Pr(A + B) = \Pr(AB + AB' + A'B) \quad (0.15)$$

$$\Pr(A + B) = \Pr(AB) + \Pr(AB') + \Pr(A'B) \quad (0.16)$$

$$\Pr(A + B) = \Pr(AB) + \Pr(A) - \Pr(AB) + \Pr(B) - \Pr(AB) \quad (0.17)$$

$$\Rightarrow \Pr(A + B) = \Pr(A) + \Pr(B) - \Pr(AB) \quad (0.18)$$

$$P(A) = 1 \quad (0.19)$$

$$P(B) = 0 \quad (0.20)$$

$$P(AB) = 0 \quad (0.21)$$

Using the values of  $\Pr(A)$ ,  $\Pr(B)$  and  $\Pr(AB)$ ,

$$\Pr(A + B) = 1 + 0 - 0 \quad (0.22)$$

$$\Pr(A + B) = 1 \quad (0.23)$$

Therefore, the value of  $\Pr(A + B)$  is 1.

### **Simulated Solution:**

Let  $X_1$  be an indicator random variable of the event  $A$ .

$X_1$  is defined as:

$$X_1 = \begin{cases} 1, & A \\ 0, & A' \end{cases} \quad (0.24)$$

Let  $X_2$  be the indicator random variable of the event  $B$ .

$X_2$  is defined as:

$$X_2 = \begin{cases} 1, & B \\ 0, & B' \end{cases} \quad (0.25)$$

Let  $X_3$  be the indicator random variable of the event  $AB$ .

$X_3$  is defined as:

$$X_3 = \begin{cases} 1, & AB \\ 0, & (AB)' \end{cases} \quad (0.26)$$

The PMF of the random variable  $X_1$  is:

$$p_{X_1}(n) = \begin{cases} p_1, & n = 1 \\ 1 - p_1, & n = 0 \end{cases} \quad (0.27)$$

The PMF of the random variable  $X_2$  is:

$$p_{X_2}(n) = \begin{cases} p_2, & n = 1 \\ 1 - p_2, & n = 0 \end{cases} \quad (0.28)$$

The PMF of the random variable  $X_3$  is:

$$p_{X_3}(n) = \begin{cases} p_3, & n = 1 \\ 1 - p_3, & n = 0 \end{cases} \quad (0.29)$$

where,

$$p_1 = 1 \quad (0.30)$$

$$p_2 = 0 \quad (0.31)$$

$$p_3 = 0 \quad (0.32)$$

$$(0.33)$$

Let  $Y$  be the random variable which is defined as follows:

$$Y = X_1 + X_2 - X_3 \quad (0.34)$$

But we know that  $X_3$  can never be 0 when  $X_1$  and  $X_2$  are 1 and vice versa. So,  $Y$  is another Indicator Random variable whose PMF is defined as:

$$p_Y(n) = \begin{cases} p, & n = 1 \\ 1 - p, & n = 0 \end{cases} \quad (0.35)$$

From (34),

$$E(Y) = E(X_1 + X_2 - X_3) \quad (0.36)$$

$$E(Y) = E(X_1) + E(X_2) - E(X_3) \quad (0.37)$$

$$1.(p) + 0.(1 - p) = 1.(p_1) + 0.(1 - p_1) + 1.(p_2) + 0.(1 - p_2) - 1.(p_3) - 0.(1 - p_3) \quad (0.38)$$

$$p = p_1 + p_2 - p_3 \quad (0.39)$$

Through our definition, we know that,

$$\Pr(A) = p_1 \quad (0.40)$$

$$\Pr(B) = p_2 \quad (0.41)$$

$$\Pr(AB) = p_3 \quad (0.42)$$

Therefore, by comparison of the axiom

$$\Pr(A + B) = \Pr(A) + \Pr(B) - \Pr(AB) \quad (0.43)$$

and the equation (0.39),

$$p = \Pr(A + B) \quad (0.44)$$


$$\Pr(A + B) = 1 + 0 - 0 \quad (0.45)$$

$$\implies \Pr(A + B) = 1 \quad (0.46)$$

Below is the plot for the simulation of the probabilities, where the grey stems represent the theoretical probabilities and the coloured stems represent the simulated ones. Through observation in the last stem, we have proved through the code that

$$\Pr(A + B) = \Pr(A) + \Pr(B) - \Pr(AB) \quad (0.47)$$

**Plot:**



Acknowledgement: <https://github.com/Akshara778/EE1003/tree/main/assignment-9>