

Matrix, Linear Algebra, Differential Equation
MAT 2207

Jannat Tohfa Chowdhury

April 2024

Matrix

Definition of Matrix:

A system of any $m \times n$ numbers arranged in a rectangular arrangement of m rows and n columns is called a matrix of order $m \times n$ or an $m \times n$ matrix.

Ex:

$$\begin{bmatrix} 1 & -2 & 4 \\ 3 & 1 & 7 \end{bmatrix} \quad \text{is a } 2 \times 3 \text{ matrix.}$$

in general form:

$$A = \begin{bmatrix} \sigma_{11} & \sigma_{12} & \cdots & \sigma_{1n} \\ \sigma_{21} & \sigma_{22} & \cdots & \sigma_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ \sigma_{m1} & \sigma_{m2} & \cdots & \sigma_{mm} \end{bmatrix} = (\sigma_{ij})_{m \times n}$$

Singular and Non-singular Matrix:

Let A be any square matrix. If $\det(A) = 0$, then A is called a singular matrix, and if $\det(A) \neq 0$, then A is called a non-singular matrix.

Ex: $A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \quad B = \begin{bmatrix} 1 & 5 \\ 2 & 12 \end{bmatrix}$

$$\text{Then } |A| = \det(A) = \det \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} = 4 - 4 = 0$$

So, A is a singular matrix

$$\text{Again, } |B| = \det(B) = \det \begin{bmatrix} 1 & 5 \\ 2 & 12 \end{bmatrix} = 12 - 10 = 2 \neq 0$$

Hence, B is a non-singular matrix.

Inverse Matrix:

Let A and B be two $n \times n$ square matrices such that $AB = BA = I_n = I$, then B is said to be the inverse of A , and we write $B = A^{-1}$. Also, $A = B^{-1}$.

Ex: Let $A = \begin{bmatrix} 4 & 3 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & -3 \\ -1 & 4 \end{bmatrix}$.

$$\begin{aligned} \therefore AB &= \begin{bmatrix} 4 & 3 \\ 1 & 1 \end{bmatrix} \times \begin{bmatrix} 1 & -3 \\ -1 & 4 \end{bmatrix} \\ &= \begin{bmatrix} 4-3 & -12+12 \\ 1-1 & -3+4 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I_2 \end{aligned}$$

$$\begin{aligned} \text{and } BA &= \begin{bmatrix} 1 & -3 \\ -1 & 4 \end{bmatrix} \times \begin{bmatrix} 4 & 3 \\ 1 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 4-3 & 3-3 \\ -4+4 & -3+4 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I_2 \end{aligned}$$

$$\therefore AB = BA = I_2 = I$$

Therefore, we can write $A = B^{-1}$ and $B = A^{-1}$.

[N.B.: The inverse of a matrix exists only when the matrix is non-singular, i.e., $|A| \neq 0$.]

*** Multiplication of two matrices is possible only when the number of columns in the first matrix is equal to the number of rows in the second matrix.

Echelon Matrix:

Let $A = (a_{ij})_{m \times n}$ be any matrix. Then A is said to be an echelon matrix or is said to be in echelon form if:

1. all the non-zero rows (if any) precede the zero rows,
2. the number of zero entries preceding the first non-zero entry in each row increases by row.

Ex:

$\begin{bmatrix} 1 & -1 & 2 \\ 0 & 3 & 2 \\ 0 & 0 & 5 \end{bmatrix}$ is an echelon matrix, but $\begin{bmatrix} 1 & 2 & 3 \\ 0 & -1 & 3 \\ 2 & 5 & 4 \end{bmatrix}$ is not an echelon matrix.

Rank of a Matrix:

Rank of a matrix is the largest non-zero row in the matrix of row echelon form.

Ex: $A = \begin{bmatrix} 1 & 4 & 5 \\ 0 & 3 & 7 \\ 0 & 0 & 6 \end{bmatrix}$ $B = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 6 & 7 \\ 0 & 0 & 0 \end{bmatrix}$

Rank of matrix A = 3, Rank of matrix B = 2

Find the rank of the following matrices:

$$A = \begin{bmatrix} 1 & -1 & -1 & 2 \\ 2 & 1 & 3 & -1 \\ 3 & 2 & 1 & 2 \\ 4 & 1 & 2 & 3 \end{bmatrix} \quad B = \begin{bmatrix} 1 & 3 & 1 & -2 & -3 \\ 1 & 4 & 3 & -1 & -4 \\ 2 & 3 & -4 & -7 & -3 \\ 3 & 8 & 1 & -7 & -8 \end{bmatrix}$$

$$C = \begin{bmatrix} 1 & -1 & 2 & 1 \\ 3 & 0 & 2 & 2 \\ 2 & 1 & -1 & 1 \\ 1 & 0 & 1 & 1 \end{bmatrix} \quad D = \begin{bmatrix} 1 & -1 & 1 & 3 \\ 2 & -1 & 0 & 2 \\ 1 & 0 & 1 & 1 \\ 4 & 2 & 0 & 1 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & -1 & 1 & 2 \\ 2 & 1 & 3 & -1 \\ 3 & 2 & 1 & 2 \\ 4 & 1 & 2 & 3 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & -1 & 1 & 2 \\ 0 & 3 & 5 & -5 \\ 0 & 5 & 4 & -4 \\ 0 & 5 & 6 & -5 \end{bmatrix} \quad \begin{bmatrix} r_2' = r_2 - 2r_1 \\ r_3' = r_3 - 3r_1 \\ r_4' = r_4 - 4r_1 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & -1 & 1 & 2 \\ 0 & 3 & 5 & -5 \\ 0 & 0 & -2 & 1 \\ 0 & 0 & 2 & -1 \end{bmatrix} \quad \begin{bmatrix} r_3'' = r_3' - r_4' \\ r_4'' = r_4' - r_3' \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & -1 & 1 & 2 \\ 0 & 3 & 5 & -5 \\ 0 & 0 & -2 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \begin{bmatrix} r_4''' = r_4'' + r_3'' \end{bmatrix}$$

This is the echelon form of matrix A; and the Rank of A is 3.

$$B = \begin{bmatrix} 1 & 3 & 1 & -2 & -3 \\ 1 & 4 & 3 & -1 & -4 \\ 2 & 3 & -4 & -7 & -3 \\ 3 & 8 & 1 & -7 & -8 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & 3 & 1 & -2 & -3 \\ 0 & 1 & 2 & 1 & -1 \\ 0 & -5 & -10 & -5 & 5 \\ 0 & -4 & -8 & -4 & 4 \end{bmatrix} \quad \begin{bmatrix} r_2' = r_2 - r_1 \\ r_3' = r_3 - 2r_2 \\ r_4' = r_4 - 3r_2 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & 3 & 1 & -2 & -3 \\ 0 & 1 & 2 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad \begin{bmatrix} r_3'' = r_3' - 5r_2' \\ r_4'' = r_4' + 4r_2' \end{bmatrix}$$

This is the echelon form of matrix B; and the Rank of B is 2.

$$C = \begin{bmatrix} 1 & -1 & 2 & 1 \\ 3 & 0 & 2 & 2 \\ 2 & 1 & -1 & 1 \\ 1 & 0 & 1 & 1 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & -1 & 2 & 1 \\ 0 & 3 & -4 & -1 \\ 0 & 3 & -5 & -1 \\ 0 & 0 & -1 & -1 \end{bmatrix} \quad \begin{bmatrix} r_2' = r_2 - 3r_1 \\ r_3' = r_3 - 2r_1 \\ r_4' = r_2 - 3r_4 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & -1 & 2 & 1 \\ 0 & 3 & -4 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & -1 & -1 \end{bmatrix} \quad \begin{bmatrix} r_3'' = r_3' - r_2' \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & -1 & 2 & 1 \\ 0 & 3 & -4 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix} \quad \begin{bmatrix} r_4''' = r_4'' - r_3'' \end{bmatrix}$$

This is the echelon form of matrix C; and the Rank of C is 4.

$$D = \begin{bmatrix} 1 & -1 & 1 & 3 \\ 2 & -1 & 0 & 2 \\ 1 & 0 & 1 & 1 \\ 4 & 2 & 0 & 1 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & -1 & 1 & 3 \\ 0 & 1 & -2 & -4 \\ 0 & 1 & 0 & -2 \\ 0 & 4 & 0 & -3 \end{bmatrix} \quad \begin{bmatrix} r_2' = r_2 - 2r_1 \\ r_3' = r_3 - r_1 \\ r_4' = r_4 - 2r_2 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & -1 & 1 & 3 \\ 0 & 1 & -2 & -4 \\ 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 5 \end{bmatrix} \quad \begin{bmatrix} r_3'' = r_3' - r_2' \\ r_4'' = r_4' - 4r_2' \end{bmatrix}$$

This is the echelon form of matrix D; and the Rank of D is 4.

Inverse Matrix Calculation:

Find the inverse of the matrix by using the formula $[A:I]$

example; find the inverse matrix of $A = \begin{bmatrix} 1 & -1 & 2 & 1 \\ 3 & 0 & 2 & 2 \\ 2 & 1 & -1 & 1 \\ 1 & 0 & 1 & 1 \end{bmatrix}$

Solution:

$$\begin{array}{l}
 A = \left[\begin{array}{cccc|cccc} 1 & -1 & 2 & 1 & 1 & 0 & 0 & 0 \\ 3 & 0 & 2 & 2 & 0 & 1 & 0 & 0 \\ 2 & 1 & -1 & 1 & 0 & 0 & 1 & 0 \\ 1 & 0 & 1 & 1 & 0 & 0 & 0 & 1 \end{array} \right] \\
 \sim \left[\begin{array}{cccc|cccc} 1 & -1 & 2 & 1 & 1 & 0 & 0 & 0 \\ 0 & 3 & -4 & -1 & -3 & 1 & 0 & 0 \\ 0 & 3 & -5 & -1 & -2 & 0 & 1 & 0 \\ 0 & 1 & -1 & 0 & -1 & 0 & 0 & 1 \end{array} \right] \begin{array}{l} r_2' = r_2 - 3r_1 \\ r_3' = r_3 - 2r_1 \\ r_4' = r_4 - r_2 \end{array} \\
 \sim \left[\begin{array}{cccc|cccc} 1 & -1 & 2 & 1 & 1 & 0 & 0 & 0 \\ 0 & 3 & -4 & -1 & -3 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 & 1 & -1 & 1 & 0 \\ 0 & 0 & 2 & 1 & -1 & 0 & -1 & 3 \end{array} \right] \begin{array}{l} r_3'' = r_3' - r_2' \\ r_4'' = 3r_4' - r_3' \end{array} \\
 \sim \left[\begin{array}{cccc|cccc} 1 & -1 & 2 & 1 & 1 & 0 & 0 & 0 \\ 0 & 3 & -4 & -1 & -3 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 & 1 & -1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & -2 & 1 & 3 \end{array} \right] \begin{array}{l} r_4''' = r_4'' + 2r_3'' \end{array} \\
 \sim \left[\begin{array}{cccc|cccc} 3 & 0 & 2 & 2 & 0 & 1 & 0 & 0 \\ 0 & 3 & -4 & -1 & -3 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 & 1 & -1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & -2 & 1 & 3 \end{array} \right] \begin{array}{l} r_1' = 3r_1 + r_2 \end{array} \\
 \sim \left[\begin{array}{cccc|cccc} 3 & 0 & 0 & 2 & 2 & -1 & 2 & 0 \\ 0 & 3 & 0 & -1 & -7 & 5 & -4 & 0 \\ 0 & 0 & -1 & 0 & 1 & -1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & -2 & 1 & 3 \end{array} \right] \begin{array}{l} r_1'' = r_1' + 2r_3'' \\ r_2'' = r_2'' - 4r_3'' \end{array} \\
 \sim \left[\begin{array}{cccc|cccc} 3 & 0 & 0 & 0 & 0 & 3 & 0 & -6 \\ 0 & 3 & 0 & 0 & -6 & 3 & -3 & 3 \\ 0 & 0 & -1 & 0 & 1 & -1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & -2 & 1 & 3 \end{array} \right] \begin{array}{l} r_1''' = r_1'' - 2r_4''' \\ r_2''' = r_2''' + r_4''' \end{array} \\
 \sim \left[\begin{array}{cccc|cccc} 1 & 0 & 0 & 0 & 0 & 1 & 0 & -2 \\ 0 & 1 & 0 & 0 & -2 & 1 & -1 & 1 \\ 0 & 0 & 1 & 0 & -1 & 1 & -1 & 0 \\ 0 & 0 & 0 & 1 & 1 & -2 & 1 & 3 \end{array} \right] \begin{array}{l} r_1''' = r_1''' \times 1/3 \\ r_2''' = r_2''' \times 1/3 \\ r_3''' = r_3''' \times -1 \end{array}
 \end{array}$$

Home Work:

find the inverse matrix of $B = \begin{bmatrix} 1 & 2 & -3 \\ 2 & 1 & 0 \\ 4 & -2 & 5 \end{bmatrix}; C = \begin{bmatrix} 1 & 3 & 1 & 1 \\ 2 & 5 & 2 & 2 \\ 1 & 3 & 8 & 9 \\ 1 & 3 & 2 & 2 \end{bmatrix}$