



**Gramin Krishi Mausam Sewa**  
**Agromet Advisory Bulletin for FIROZABAD District**  
**Chander Shekhar Azad University of Agriculture & Technology**  
 Kanpur, Uttar Pradesh



## **Agromet Advisory Bulletin**

**Date : 2025-02-04**

**Weather Forecast of District FIROZABAD (Uttar Pradesh) Issued On : 2025-02-04 (Valid Till 08:30 IST of the next 5 days)**

Parameter	2025-02-05	2025-02-06	2025-02-07	2025-02-08	2025-02-09
Rainfall(mm)	2.0	0.0	0.0	0.0	0.0
Tmax(°C)	27.0	27.0	25.0	25.0	26.0
Tmin(°C)	14.0	13.0	10.0	10.0	10.0
RH-I(%)	93	91	75	71	65
RH-II(%)	54	55	46	44	41
Wind Speed(kmph)	6	8	14	13	8
Wind Direction(Degree)	337	330	312	299	304
Cloud Cover(Octa)	2	0	1	2	4

### **Weather Summary/Alert:**

According to the weather forecast received from the Indian Meteorological Department, due to light to moderate cloud cover in the next five days, there is a possibility of light rain dated on 05 February 2025. There will be light mist in the upper surface of the atmosphere and light fogs is likely to appear in the morning and night. The maximum temperature is between 25.0- 27.0°C, which is likely to be 2-3°C higher normal and the minimum temperature is between 10.0-13.0°C, which is likely to be 1-2°C higher normal. The maximum and minimum range of relative humidity is between 65-93 and 41- 55%. Wind direction is north-west and wind speed is between 8.0-14.0 kmph with gusts expected to be 4-5 kmph high speed than normal.

### **General Advisory:**

Farmers are advised to do light irrigation in crops like wheat, mustard etc. as per requirement and prepare the field, prepare fertilizers and seeds for sowing maize and summer vegetables crops start sowing. For insecticides, pesticides and weedicide chemicals, separate or use only by washing the equipment with clean water and do not spray or spray insecticides, pesticides and weedicide chemicals standing in the opposite direction of the air. Spraying should be done in the evening, if possible, after spraying, hands should be thoroughly washed with soap or hand wash before eating and after washing the clothes.

### **SMS Advisory:**

Farmers are advised to do sowing work by arranging fertilizers and seeds for sowing of maize and summer vegetables crops.

### **Crop Specific Advisory:**

Crop (Stage)	Crop Specific Advisory
WHEAT	Third irrigation in wheat crop 60-65 days after sowing, do light irrigation at the stage of knot formation and do topdressing of one-third quantity of nitrogen after second irrigation when oat comes. If both narrow and broad

Crop (Stage)	Crop Specific Advisory
	leaf weeds appear in wheat crop sown very late, then Sulfosulfuran 75% WP at 33 gm/ha or Matribuzin 70% WP at the rate of 250 gm/ha can be used to control it. Make a solution in 600 liters of water and spray.
MUSTARD	Apply second irrigation in mustard crop before flowering or at seed filling stage. There is a due to continuous cloudy sky, mustard crop is likely to show outbreak of Gundi bug, Painted class and leaf mulch in mustard crop, so Chlorpyrifos rate of 20% EC 1.0 liter / ha or Monocrotophos 37% SL @ 500 ml / ha for its prevention spray from 500 - 600 liters of water in clear sky.
FIELD PEA	Due to excess moisture in the environment, symptoms like white powder appear on the leaves, stems and legumes of the pea crop, (Buchani's disease) for its prevention, spraying of tridomorph 80% EC @ 500 ml per hectare in 500 to 600 liters of water and spraying it at 12-15 days interval when the sky is clear.
CHICK PEA	Stop the work of nipping in the gram crop . In irrigated gram, do light irrigations before flowering or at the stage of grain filling. There is a possibility of outbreak of cutworm (caterpillar insect) in gram crop, to prevent this, chlorpyrifos 50% EC + Spemethrin 5% EC at a rate of 2.0 liters / ha is sprayed in 500-600 liters of water.
MAIZE	Do to Zaid maize is composite variety - Kanchan, Gaurav, Shveta, Azad Uttam and Hybrid variety - Prakash, Decalb - 7074, PHB - 8144, Deccan 115, MMH - 133 arrange of fertilizer and seed for sowing of any one of the varieties. For sowing of maize composite variety, sow 20-25 kg seed / hectare and Shankar variety at the rate of 18-20 kg seed / ha.

#### Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
POTATO	Do not do any kind of agricultural operations in the potato crop sown on time. Increasing humidity / cloudy environment and falling temperature causes rapid outbreak of scorching disease in potato crop, hence Mancozeb or Ridomil 2.5 g / liter of water or copper oxychloride of 3.0 g / liter of water to prevent it make slurry and spray it at 12-15 days interval.
ONION	Prepare the field for sowing vegetables such as ladyfinger, luffa, cucumber, bitter gourd, gourd and pumpkin during summer season. Transplant the prepared onions and irrigate them immediately after planting. For weed control in onion crop, spraying of Pendimethalin 30% EC.3.5 liters of medicine / ha in 800 liters of water immediately after transplanting or just before irrigation. In tomato, chili crop, in case of late blight and outbreak of bacterial built disease, spray a mixture of 30 grams of copper oxychloride and 1 gram of streptocillin in 10 liters of water.
MANGO	In mango orchards, there is a possibility of moth and mango hopper pest infestation, so spraying Chlorpyrifos 20 EC 2.0 ml or Phenitrothian 50 EC 3.0 ml / liter water is sprayed to prevent it. To prevent the plum fruits from falling, spray the Super 6 hormone 1.0 ml per 4.5 liter water solution in clear sky. Planting soil on plants while weeding the banana plantation, prune the leaves and support the plants with bamboo or wood when the flowers and fruits emerge. Clean the mango trees and cut the twigs that are dry and diseased.

#### Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	In view of the current weather, farmers are advised to swing the animals in the morning and evening to protect them from the cold, their animals in the open during the night and Draw and put curtains of jut sack on windows and doors at night and remove put curtains in the sun during the day. Give sufficient quantity of grains to the animals with green and dry fodder. Animals must be given fresh and fresh water 3-4 times a day. Keep the animals in a clean place.

**Poultry Specific Advisory:**

Poultry	Poultry Specific Advisory
HEN	Farmers are advised to give provide light to the chickens 15 to 16 hours a day. The hen supplements the food, mix vitamins and energy food material in their feed as well as calcium content to hens. Arrange enough heat to keep the chickens from freezing.