



Gramin Krishi Mausam Sewa
Agromet Advisory Bulletin for UNNAO District
Chander Shekhar Azad University of Agriculture & Technology
Kanpur, Uttar Pradesh



Agromet Advisory Bulletin

Date : 2025-05-27

Weather Forecast of District UNNAO (Uttar Pradesh) Issued On : 2025-05-27 (Valid Till 08:30 IST of the next 5 days)

Parameter	2025-05-28	2025-05-29	2025-05-30	2025-05-31	2025-06-01
Rainfall(mm)	0.0	0.0	2.0	0.0	10.0
Tmax(°C)	39.0	38.0	40.0	39.0	39.0
Tmin(°C)	29.0	29.0	29.0	28.0	27.0
RH-I(%)	65	65	63	68	78
RH-II(%)	47	47	41	45	49
Wind Speed(kmph)	16	15	10	12	14
Wind Direction(Degree)	103	99	96	92	110
Cloud Cover(Octa)	2	2	5	6	5
Warning	No Warning	No Warning	No Warning	No Warning	No Warning

Forecast Summary:

According to the weather forecast received from the Indian Meteorological Department, due to light to moderate cloud cover in the next five day there is a possibility of light to moderate rain on 30 May & 01 June, 2025. The maximum temperature is between 38.0- 40.0°C, which is likely to be near to its normal and the minimum temperature is between 27.0-29.0°C, which is likely to be 1-2°C higher normal. The maximum and minimum range of relative humidity is between 65-78 and 41-47%. Wind direction is south-east and wind speed is between 10.0-16.0 kmph with gusts expected to be 4-5 kmph high speed than normal.

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the Indian Meteorological Department, there is a no warning in next five days.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

In view of light to moderate rain with thunderstorm/lightning and strong surface winds, farmers are advised not to tie animals in open space/under trees and also postpone their travel or activity. If you are in a group during a thunderstorm, separate from each other. This will reduce the number of injuries if the lightning strikes the ground.

General Advisory:

In view of the possibility of light to moderate rain with thunderstorms lightning and strong surface winds, farmers are advised to irrigate the standing maize crop only when the winds are calm. Spray

insecticides, fungicides and weedicides only when the winds are calm/the sky is clear and do not spray insecticides, pesticides and weedicides while standing in the opposite direction of the wind. If possible, hands should be washed thoroughly with soap or hand wash after spraying, before eating and after washing clothes.

SMS Advisory:

In view of the possibility of rain, thunderstorms, strong winds, farmers are advised to postpone irrigation work in Zaid crops .

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
MAIZE (Millking)	Maize crop is going on at the stage of cob formation/grain filling which is sensitive to moisture deficiency. Therefore, maintain proper moisture by giving light irrigation as per requirements. There is a possibility of outbreak of stem borer / shoot fly pest in maize crop, so for its prevention, solution of Emamectin Benzoate 5% SG at the rate of 200 g/ha or Dimethoate 30% EC 1.0 l/ha in 600-700 liters of water make a spray.
BLACK GRAM (Pod initiation)	Black gram/green gram crop is progressing from flower to pod formation stage which is sensitive to moisture deficiency. Therefore, maintain proper moisture by applying light irrigation as per requirement. There is a possibility of outbreak of pod borer in black gram crop, so to prevent it, apply Emamectin Benzoate 5% SG 220 g / hectare or Dimethoate 30% EC. Spray at the rate of 1.0 liters/hectare in 600-700 liters of water when the sky is clear.
MAIZE (Sowing)	Prepare the field for sowing Kharif maize only after ploughing, and also make arrangement for fertilizers and seeds for sowing any one of the improved recommended complex varieties- Kanchan, Gaurav, Sweta, Azad Uttam and hybrid varieties- Prakash, Y-1402, Bio-9681, Pro-316, D-9144, Dikalb- 7074, PHB- 8144, Deccan 115, MMH-133, etc. For sowing complex varieties of maize, sow at the rate of 20-25 kg seed/hectare and Shankar variety at the rate of 18-20 kg seed/hectare.
RICE (Nursery)	Prepare the fields for planting paddy nursery, as well as the advanced varieties Narendra-359, Narendra paddy-2026, Narendra paddy-2064, Narendra paddy-2065, Narendra paddy-3112, Sarju-52, Sita etc. and hybrid varieties- Pro .Agro-6444 (2001)Arise , Prof.Agro-6201 (2001)Arise, PHB-71, Pioneer-27 P 31, 27 P 37, 28 P 67 , RR X-113, Kaveri - 468, KRH-2, US-312, RH. -312, rh. -1531etc and fragrant paddy varieties- Type-3, Pusa Basmati-1, Malviya Sugandh-105, Malviya Sugandha-3-4, Narendra Sugandh and Narendra Usar Paddy-1, Narendra Usar Paddy-2, Narendra Usar Paddy-2008, Arrange any one variety of seeds and fertilizers from CLR-10 etc. Light irrigation should be done in the field 15 days before planting the nursery of paddy seeds so that the weeds coming out in the field are destroyed while preparing the field.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
BRINJAL (Physiological Maturity)	Harvest the finished crops like Pumpkin, Gourd, Taroi, Bitter gourd, Cucumber, Cucumber, Watermelon and Melon etc. and send them to the market. For the prevention of fruit borer / leaf borer pest in vegetable crops, make a solution of neem oil in 1.5-2.0 ml / liter of water and do 3-4 sprays at an interval of 8 -10 days. Do weeding and hoeing in the standing crops of vegetables and do irrigation work at an interval of 8-10 days in the evening.
MANGO	Dig pits for planting new gardens. Do irrigation work in orchards of mango, guava, lemon plum, grapes, papaya and litchi etc. Irrigate the orchards to prevent fall of mango fruit or apply 4.5 SL of 20 ml alpha naphthalene

Horticulture (Stage)	Horticulture Specific Advisory
	acetic acid make a solution per liter of water and spray it. To prevent cuckoo disorder or internal rot disease in mango pods, make borax 0.8% solution in 0.8 g / liter water and spray 2 at 10 days interval.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	Considering the current weather, farmers are advised to tie the animals in the open during the night. Tie the animals in a shade time place during the day. Do not tie the animals under the tree because this week the chances of falling trees twigs are high due to the speed of the winds. FMD for the prevention of foot-mouth disease to animal's vaccine and VQ for prevention of Lagadia fever. Give sufficient quantity of grains to the animals with green and dry fodder. Animals must be given clean and fresh water 3-4 times a day. Do not tie pregnant animals to the sloping place. The best time to give anthelmintic drugs for the prevention of insects in the stomach of animals.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Farmers are advised to add dietary supplements; vitamins and energy feed ingredients to the hens as well as add calcium content to the hen. Give medicine for the prevention of worms in the stomach of hen (Divmirg). To protect the chickens from heat, arrange curtains, fans and ventilation in the hen house. To save the chickens from the heat, sprinkle water on the curtains engaged in the chicken house so that the coolness remains.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation)	Others (Soil / Land Preparation) Specific Advisory
AGRI FIELD	Deep plow the fields empty of wheat and other crops with disc plow so that the eggs and larvae of harmful insects hidden inside the ground are destroyed in the sun.

Likely impacts of weather warnings (General)

According to the weather forecast received from the Indian Meteorological Department, there is a warning on thunderstorms & lightning, squall, strong surface winds at a local level.

Impact based advisories (General)

In view of light to moderate rain with thunderstorm/lightning and strong surface winds, farmers are advised not to tie animals in open space/under trees and also postpone their travel or activity. If you are in a group during a thunderstorm, separate from each other. This will reduce the number of injuries if the lightning strikes the ground.
--

Farmers are advised to download Unified "Mausam" and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: <https://play.google.com/store/apps/details/>

Meghdoot MobileApp link: <https://play.google.com/store/apps/details>

Damini MobileApp link : <https://play.google.com/store/apps/details>