



**Gramin Krishi Mausam Sewa**  
**Agromet Advisory Bulletin for MATHURA District**  
**Chander Shekhar Azad University of Agriculture & Technology**  
Kanpur, Uttar Pradesh



## **Agromet Advisory Bulletin**

Date : 2026-01-27

Weather Forecast of District MATHURA (Uttar Pradesh) Issued On : 2026-01-27 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-01-28	2026-01-29	2026-01-30	2026-01-31	2026-02-01
Rainfall(mm)	15.0	0.0	0.0	0.0	0.0
Tmax(°C)	16.0	18.0	20.0	21.0	22.0
Tmin(°C)	9.0	10.0	8.0	10.0	11.0
RH-I(%)	99	97	96	87	86
RH-II(%)	70	71	62	56	56
Wind Speed(kmph)	7	3	9	9	7
Wind Direction(Degree)	103	284	331	336	23
Cloud Cover(Octa)	6	2	4	4	2
Warning	Thunderstorms & Lightning, Squall etc; Hailstorm; Strong Surface Winds	Fog	No Warning	No Warning	No Warning

### Forecast Summary:

According to the weather forecast from the Indian Meteorological Department, due to moderate to dense cloud cover in the remaining days, there is a possibility of light to moderate rain with local level at strong winds, hailstorm, thunderstorm & lightning dated on 28 January, 2026. The maximum temperature is between 16.0- 22.0°C, which is likely to be 1-2°C higher normal and the minimum temperature is between 9.0-11.0°C, which is likely to be 2-3°C higher normal. The maximum and minimum range of relative humidity is between 86-99 and 56-71%. The wind direction will be from the south-east, north-west and the wind speed will be between 3.0-9.0 km/hour, which is expected to be 3-4 km/hour higher than normal.

### Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the Indian Meteorological Department, a warning has been issued for light rain with Thunderstorms & Lightning, Strong Winds dated on 28 January, 2026.

### Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Farmers are advised to postpone irrigation and herbicide application to standing Rabi crops in view of the possibility of rain. To protect crops such as mustard, chickpeas, kidney beans, and peas from cold waves/frost, spray a solution of 0.1% sulfuric acid (1 liter in 1000 liters of water) when the sky is clear.

### General Advisory:

Farmers are advised to postpone irrigation and herbicide/pesticide spraying on standing Rabi crops in view of the possibility of rain. To protect crops like mustard, chickpeas, kidney beans, and peas from the cold wave, spray them with a solution of 0.1% sulfuric acid (1 liter in 1000 liters of water). People are advised to wear warm clothes, stay indoors as much as possible, and minimize travel to avoid exposure to cold winds. Use fog lights while driving. In dense fog, follow the road markings for safe driving. Also, cover livestock with jute sacks at night and do not tie them in open areas.

### SMS Advisory:

Farmers are advised to postpone irrigation and herbicide/pesticide spraying on standing Rabi crops in view of the possibility of rain.

### Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
WHEAT (Tillering)	Given the possibility of rain, farmers are advised to postpone irrigation and weedicide application in standing wheat crops. If there is no rain, apply the second irrigation 40-45 days after sowing at the tillering stage and the third irrigation 60-65 days after sowing at the jointing stage. Apply light irrigation and nitrogen fertilizer. After the second irrigation, when the crop starts to show signs of growth, apply one-third of the total fertilizer dose. If both narrow-leaved and broad-leaved weeds are present in the wheat crop, after the first irrigation, when there is sufficient moisture, apply sulfosulfuron 75% WP at the rate of 33 grams per hectare or metribuzin 70% WP at the rate of 250 grams per hectare, dissolved in 500-600 liters of water.
MUSTARD (Pod Development)	Given the possibility of rain, postpone irrigation of standing mustard crops. Apply second irrigation in mustard crop before flowering or at seed filling stage. There is a due to continuous cloudy sky, mustard crop is likely to show outbreak of Gundi bug, Painted class and leaf mulch in mustard crop, so Chlorpyrifos rate of 20% EC 1.0 liter / ha or Monocrotophos 37% SL @ 500 ml / ha for its prevention spray from 500 - 600 liters of water in clear sky.
FIELD PEA (Pod formation)	Given the possibility of rain, postpone irrigation, pesticide application, and disease control measures for standing pea crops. For the prevention of ducal disease spreading like white powder on leaves, stems and pods due to excess moisture in pea crop, use soluble sulfur 80 % 2 kg or Tridemorph 80% EC at the rate of 50 ml/ha Spray the solution in 500-600 liters of water.
CHICK PEA (Flowering)	Given the possibility of rain, postpone irrigation, pesticide application, and disease control measures for standing chickpea crops. Stop the work of nipping in the gram crop. In irrigated gram, do light irrigations before flowering or at the stage of grain filling. There is a possibility of outbreak of cutworm (caterpillar insect) in chick pea crop, to prevent this, chlorpyrifos 50% EC + Spemethrin 5% EC at a rate of 2.0 liters / ha is sprayed in 500-600 liters of water.

### Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
POTATO (Vegetative )	Given the possibility of rain, postpone irrigation, pesticide application, and disease control measures for standing potato crops. There is a possibility of outbreak of late blight disease in potato crop, so for its prevention, spray Mancozeb (2.0 g/Litre water) or 0.2% Dithane M-45 (1.5 Ml/Litre water) solution do in clear weather.If the weather is damp and cloudy for several days, then 3-4 spraying should be done at an interval of 10-12 days as per the requirement.

Horticulture (Stage)	Horticulture Specific Advisory
ONION (Transplanting)	Farmers are advised to postpone irrigation and herbicide/pesticide spraying on onion crop. In case of late blight and bacterial wilt disease outbreak in tomato crop, spray mixture of 30 gm copper oxychloride and 1 gm streptomycin in 10 liters of water. If the infestation of fruit borer / leaf borer insect is visible in vegetable crops, then for its control, make a solution of neem oil in 1.5 ml / liter of water and do 3-4 sprays at an interval of 8-10 days. Damping off (humid rot) disease is likely to be seen in onion nursery, for its prevention, spray Thiram 2.5 gm or Macozeb 2.5 gm/Litre of water by making a solution. Recommend onion species - Kalyanpur Red Gol, Pusa Ratanar, Agrifound Light Red, X Kylever, Burgundy, KP, Orient, Rosie etc. for nursery of any one variety also, transplant the prepared seedlings.
MANGO (Transplanting)	Farmers are advised to postpone irrigation and herbicide/pesticide spraying on mango crop. To prevent mango blossoms from blight, spray a 0.2 percent solution of Mancozeb + Carbendazim (2.0 grams per liter of water) or a solution of Trifloxystrobin 25% + Tebuconazole 50% (0.25 grams per liter of water). For better flowering in mango trees, spray an NPK mixture dissolved in water at a rate of 5 kg per 2000 liters of water; this is sufficient for 50 trees. If mango plants show signs of flower and flower cluster midge infestation, spray with Dimethoate (30 percent active ingredient) at a rate of 2.0 ml per liter of water for control.

#### Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	In view of the possibility of change in weather, farmers are advised not to tie their animals in the open during night and put jute sack curtains on windows and doors at night and remove the curtains during the day in the sun. Vaccinate animals to prevent foot and mouth disease. Wash wounds of animals affected by this disease with potassium permanganate. Do not tie pregnant buffaloes/cows on slopes. Feed pregnant buffaloes/cows nutritious fodder and grains, and make sure to feed newborn calves knees for three days. Provide animals with adequate amounts of grain, along with green and dry fodder. To prevent the infection of Kilni/Lice, spray lime in the cattle sheds so that the eggs and larvae of Kilni/Lice are destroyed. To prevent all reproductive diseases, livestock farmers are advised to feed green fodder, straw, animal feed and mineral salts. Animals must be provided with clean, fresh water two to three times a day. Farmers/livestock owners can avail the benefits of this scheme by contacting the toll-free helpline number 1962 to provide veterinary services at their doorstep.

#### Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Farmers are advised to provide light to the chickens for 15 to 16 hours a day. Add dietary supplements, vitamins and energy feed ingredients to the hens as well as the calcium content of the chickens. Provide adequate heat to keep the chicks from freezing.

#### Likely impacts of weather warnings (General)

According to the weather forecast received from the Indian Meteorological Department, a warning has been issued for light rain with Thunderstorms & Lightning, Strong Winds dated on 28-29 January, 2026.

#### Impact based advisories (General)

Farmers are advised to postpone irrigation and herbicide application to standing Rabi crops in view of the possibility of rain. To protect crops such as mustard, chickpeas, kidney beans, and peas from cold waves/frost, spray a solution of 0.1% sulfuric acid (1 liter in 1000 liters of water) when the sky is clear.

**Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.**

**Mausam MobileApp link: <https://play.google.com/store/apps/details/>**

**Meghdoot MobileApp link: <https://play.google.com/store/apps/details>**

**Damini MobileApp link : <https://play.google.com/store/apps/details>**