



Gramin Krishi Mausam Sewa
Agromet Advisory Bulletin for KANPUR-NAGAR District
Chander Shekhar Azad University of Agriculture &
Technology
Kanpur, Uttar Pradesh



Agromet Advisory Bulletin

Date : 2025-06-20

Weather Forecast of District KANPUR-NAGAR (Uttar Pradesh) Issued On : 2025-06-20 (Valid Till 08:30 IST of the next 5 days)

Parameter	2025-06-21	2025-06-22	2025-06-23	2025-06-24	2025-06-25
Rainfall(mm)	25.0	12.0	25.0	15.0	6.0
Tmax(°C)	33.0	34.0	34.0	32.0	33.0
Tmin(°C)	26.0	27.0	26.0	25.0	25.0
RH-I(%)	88	85	90	91	90
RH-II(%)	67	62	63	66	67
Wind Speed(kmph)	5	4	5	3	6
Wind Direction(Degree)	132	285	278	220	245
Cloud Cover(Octa)	7	6	7	7	5
Warning	Thunderstorms & Lightning, Squall etc; Strong Surface Winds	Heavy Rain; Thunderstorms & Lightning, Squall etc; Strong Surface Winds	Heavy Rain; Thunderstorms & Lightning, Squall etc; Strong Surface Winds	Thunderstorms & Lightning, Squall etc	No Warning

Forecast Summary:

According to the weather forecast received from the Indian Meteorological Department, due to moderate to dense cloud cover in the next five day there is a possibility of light to moderate rain with thunderstorms & lightning, strong winds at local level between dated on 21-25 June, 2025. The maximum temperature is between 32.0- 34.0°C, which is likely to be 2-3°C lower normal and the minimum temperature is between 25.0-27.0°C, which is likely to be near its normal. The maximum and minimum range of relative humidity is between 88-91 and 62-67%. Wind direction is south-east, north-west, south-west and wind speed is between 5.0-7.0 kmph with gusts expected to be 1-2 kmph high speed than normal.

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the Indian Meteorological Department, there is a warning on moderate to heavy rain, thunderstorms & lightning, strong winds at a local level between dated on 21-25 June 2025.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

In view of the possibility of moderate to heavy rainfall in the coming week, farmers are advised to make proper arrangements for the drainage of excess rainwater. Harvest the ripe Zaid crops and keep the grains safe. Taking advantage of the rains that occurred last week, prepare the field for transplanting the ready paddy saplings and plant the saplings and sow Kharif crops only when the sky is clear and there is proper moisture. Do not tie the animals in open spaces during the rainy season.

General Advisory:

If the harvested crop in the field has got wet due to last week's rain, then dry the crop well in the sun and complete the threshing work soon. In view of the possibility of moderate to heavy rain in the coming week, farmers are advised to make proper arrangement for drainage of excess rain water. Keep the harvested crop at a safe place in the field or collect the crop in the field and cover it well with a polythene sheet. Strengthen the bunds of the fields for transplanting paddy saplings, so that rainwater remains retained in the fields. Postpone sowing of remaining Kharif crops except paddy crop.

SMS Advisory:

In view of the possibility of rain, transplant the prepared paddy saplings and postpone the sowing of Kharif crops like maize, sesame, pigeon pea etc.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
RICE (Transplanting)	Make proper arrangements for the drainage of excess rainwater from the paddy nursery. Strengthen the bunds of the fields for transplanting paddy saplings. So that rainwater remains in the fields. Plant nurseries of early maturing paddy varieties. If transplanting paddy saplings in normal fields, transplant them in 20 to 25 days and in barren land, transplant saplings of 25 to 30 days. Khaira disease in the nursery occurs due to zinc deficiency in which leaves turn yellow and later brown spots appear. To control this, when symptoms appear, spray 5 kg zinc sulphate with 2 percent urea solution or 2.5 kg slaked lime by making a solution in about 1000 liters of water per hectare.
MAIZE (Sowing)	In view of the possibility of rain, postpone sowing work. After preparing the field for sowing Kharif maize, arrange for fertilizers and seeds for sowing any one of the improved recommended complex varieties - Kanchan, Gaurav, Shweta, Azad Uttam and hybrid varieties - HMM-3904, PMH-3, NK-61, Prakash, Y-1402, Bio-9681, Pro-316, D-9144, Dikab- 7074, PHB- 8144, Deccan 115, MMH-133 etc. For sowing complex variety of maize, sow at the rate of 20-25 kg seed/hectare and Shankar variety at the rate of 18-20 kg seed/hectare.
SESAME (GINGELLY/TIL) (Sowing)	In view of the possibility of rain, postpone sowing work. Sow improved varieties of sesame such as Gujarat Til-6, R.T.-346, R.T.-351, Tarun, Pragati, Shekhar Type-78, Type-13, Type-4 and Type-12, M.T.-2013-3 and U.A.T. Til-1.
SUGARCANE (Vegetative)	In view of the possibility of rain, postpone irrigation work. Weeding should be done to control weeds in standing sugarcane crop. All these agricultural works should be completed before the onset of monsoon rains. For controlling top borer pest, destroy the affected young plant by cutting it from the ground level. If severe infestation is seen, then for its control, spray 150-200 ml of Chlorantraniliprole 18.5 S.C. Dissolve the pesticide in 400-500 liters of water and spray it per acre and irrigate.
MAIZE (Maturity and Harvesting)	Considering the possibility of rain, keep the harvested maize crop in a safe place or collect the harvested crop in the field and cover it with polythene sheet. After rain, dry the crop thoroughly in the sun and do threshing. Farmers are advised to harvest the mature maize corn and after separating the grain from the broken corn, dry it well in the sun and store it.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
CAULIFLOWER (Nursery Bed Preparation)	This is the suitable time for nursery of vegetables sown in Kharif like brinjal, chili, early cauliflower and sowing of ladyfinger, cowpea, pumpkin, bottle gourd, ridge gourd, bitter gourd, cucumber etc. For controlling stem/fruit/leaf borer insects in vegetable crops, mix 1.5-2.0 ml of Neem oil in/liter of water and spray 3-4 times at an interval of 8-10 days. For controlling green hopper and white fly insects in cucurbitaceous crops, spray 1.0 ml of Imidacloprid 30.5 percent by dissolving the chemical in one litre of water and to protect from fruit fly, use Q Lure pheromone trap at the rate of 8-10 traps per hectare.
MANGO (Transplanting)	For filling the pits dug in the month of May for planting fruit orchards, mix rotten manure of cow dung in the upper half of the soil and fill it up to 15 to 20 cm above the ground level. To protect guava from fruit flies, hang 8-10 traps per hectare using methyl eugenol and Q lure traps tied to branches at a height of 6 to 8 feet. Spray 5% neem extract dissolved in one litre of water at an interval of 10 to 15 days. Change the lure at an interval of 20-25 days.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	Farmers are advised not to tie animals in open spaces/under trees during rains. Tie animals in the open during nights. Tie animals in shady places during daytime. Vaccinate animals with FMD vaccine to prevent foot and mouth disease and VQ vaccine to prevent dengue fever. Give sufficient amount of grains to animals along with green and dry fodder. Make sure to give clean and fresh water to animals 3-4 times a day. Do not tie pregnant animals at sloping places. This is the right time to give deworming medicine to animals to prevent stomach worms.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Farmers are advised to add dietary supplements; vitamins and energy feed ingredients to the hens as well as add calcium content to the hen. Give medicine for the prevention of worms in the stomach of hen (Divmirg). To protect the chickens from heat, arrange curtains, fans and ventilation in the hen house. To save the chickens from the heat, sprinkle water on the curtains engaged in the chicken house so that the coolness remains.

Likely impacts of weather warnings (General)

According to the weather forecast received from the Indian Meteorological Department, there is a warning on moderate to heavy rain, thunderstorms & lightning, strong winds at a local level between dated on 21-25 June 2025.

Impact based advisories (General)

In view of the possibility of moderate to heavy rainfall in the coming week, farmers are advised to make proper arrangements for the drainage of excess rainwater. Harvest the ripe Zaid crops and keep the grains safe. Taking advantage of the rains that occurred last week, prepare the field for transplanting the ready paddy saplings and plant the saplings and sow Kharif crops only when the sky is clear and there is proper moisture. Do not tie the animals in open spaces during the rainy season.

Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: <https://play.google.com/store/apps/details/>

Meghdoot MobileApp link: <https://play.google.com/store/apps/details>

Damini MobileApp link : <https://play.google.com/store/apps/details>