

### Gramin Krishi Mausam Sewa

# Agromet Advisory Bulletin for SITAPUR District Chander Shekhar Azad University of Agriculture & Technology



Kanpur, Uttar Pradesh

# **Agromet Advisory Bulletin**

Date: 2025-04-08

Weather Forecast of District SITAPUR (Uttar Pradesh) Issued On: 2025-04-08 (Valid Till 08:30 IST of the next 5 days)

Parameter	2025-04- 09	2025-04- 10	2025-04-11	2025-04-12	2025-04-13
Rainfall(mm)	0.0	0.0	1.0	8.0	1.0
Tmax(°C)	39.0	38.0	37.0	37.0	37.0
Tmin(°C)	25.0	25.0	24.0	23.0	23.0
RH-I(%)	71	75	79	81	80
RH-II(%)	32	35	36	34	35
Wind Speed(kmph)	15	15	17	10	8
Wind Direction(Degree)	104	101	91	111	107
Cloud Cover(Octa)	1	3	3	3	4
Warning	No Warning	No Warning	Thunderstorms & Lightning, Squall etc	Thunderstorms & Lightning, Squall etc	Thunderstorms & Lightning, Squall etc

#### **Forecast Summary:**

According to the weather forecast received from the Indian Meteorological Department, Due to light to moderate cloud cover in the next five days there is a possibility of light rain last three days. The maximum temperature is between 37.0-39.0°C, which is likely to be 4-5°C higher normal and the minimum temperature is between 24.0-25.0°C, which is likely to be 3-4°C higher normal. The maximum and minimum range of relative humidity is between 71-81 and 32-36%. Wind direction is south-east, north-east and wind speed is between 8.0-17.0 kmph with gusts expected to be 6-7 kmph high speed than normal.

#### Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the Indian Meteorological Department, there is a warning of light rain with thunderstorms locally in last three days due to light to moderate clouds.

### Likely impacts of weather warnings on Agriculture and associated Agromet advisories

In view of the weather conditions, farmers are advised to harvest and thresh the standing crops in the fields in the morning/evening to protect them from the heat wave. Do not do any work on the fields between 11 am and 4 pm and do light irrigation in the standing crops in the evening at an interval of 10-12 days. Do not leave the animals outside for grazing between 11 am and 4 pm. Bathe the animals in the morning and evening.

# **General Advisory:**

Farmers are advised to maintain proper moisture in standing vegetables and crops by doing light irrigation as per requirement. While working in the field, tie a towel on the head and keep a water bottle with you. Keep the threshed crop grains in a safe place. To prevent infection of thrips insect and purple spot disease in onion, it is advised to keep a regular watch on the crops. Use separate equipment for insecticides, fungicides and weedicide chemicals or wash the equipment with clean water and do not spray insecticides, fungicides and weedicide chemicals while standing in the opposite direction of the wind. If possible, spray in the evening, after spraying, wash hands thoroughly with soap or hand wash before eating or drinking and wash clothes and take a bath.

# **SMS Advisory:**

Farmers are advised to maintain proper moisture in standing crops by doing light irrigation as per requirement. To avoid heat stroke, do not go out of the house unnecessarily during the afternoon.

# **Crop Specific Advisory:**

Crop (Stage)	Crop Specific Advisory	
WHEAT	In view of the possibility of rain, the harvested crop should be threshed quickly to protect the grains. Harvest the wheat opera house and then thresh it. After harvesting the wheat crop, tie the bundles together so that the bundles do not fly away due to strong winds and threshing should be done in the evening and at night when the wind remains calm.	
MAIZE	Keep the maize crop free from weeds and maintain proper moisture by irrigating lightly at an interval of 12-15 days as per requirement. For prevention of stem borer pest in maize crop, spray Emamectin Benzoate 5% SG 200 gm per acre or Chlorantraniliprole 18.5% SC 60 ml per acre by dissolving it in 500-700 liters of water.	
BLACK GRAM	Keep black gram crops free from weeds and maintain proper moisture by giving light irrigation at intervals of 12 -15 days as per requirement. Thrips/green leaf pest is likely to be seen in black gram crop, so for its prevention, spray Oxidemetan-methyl 25% EC or Dimethoate 30% EC at the rate of 1.0 liter/ha by making a solution in 600-700 liters of water.	
GREEN GRAM	Recommended species of Zaid Moong - Narendra Moong-1, Samrat (PDM-139), Sweta etc., do the work of sowing any one species at proper moisture. For sowing of zaid moong crop, complete the sowing work by April 10 at the rate of 20-25 kg seed/hectare.	

### **Horticulture Specific Advisory:**

Horticulture (Stage)	Horticulture Specific Advisory
ONION	In the summer season, irrigation should be done in the planted/sown vegetables at an interval of 8-10 days in the evening. Thrips pest is likely to appear in the onion crop, so for its prevention, use Methyl-O-Dimethan 25% EC.at the rate of 1.5 liters / ha spray the solution in 600-700 liters of water.
MANGO	Irrigate the orchards to prevent fall of mango fruit or apply 4.5 SLof of 20 ml alpha naphthalene acetic acid make a solution per liter of water and spray it. There is a possibility of showing the outbreak of midge pest in mango blossom, so for its prevention, spray Fenitrothian in 1.0 ml / liter of water and spray it.

# **Live Stock Specific Advisory:**

Live Stock	Live Stock Specific Advisory
BUFFALO	It is the appropriate time to give anthelmintic medicine to the animals for the prevention of worms in the stomach. FMD for the prevention of foot-mouth disease to animals vaccine and VQ for prevention of Lagadia fever. Get vaccinated with the vaccine. In view of the prevailing weather, farmers are advised tie the animals in a shaded place or in the shade of a tree during the day. Feed animals with green fodder and easily digestible mineral mixture and salt. The animals must be given clean and cold water 3-4 times a day. Do not leave the animals to graze from 11 am to 4 pm. bathe the animals in the morning and evening.

## **Poultry Specific Advisory:**

Poultry	Poultry Specific Advisory		
HEN	Poultry Specific Advisory  Farmers are advised to add dietary supplements; vitamins and energy feed ingredients to the hens as well as add calcium content to the hen. Give medicine for the prevention of worms in the stomach of hen (Divmirg). To protect the chickens from heat, arrange curtains, fans and ventilation in the hen house. To reduce the temperature of the hen house, cover the roof of the shed with straw.		

### Likely impacts of weather warnings (General)

According to the weather forecast received from the Indian Meteorological Department, there is a warning of light rain with thunderstorms locally in last three days due to light to moderate clouds.

### **Impact based advisories (General)**

In view of the weather conditions, farmers are advised to harvest and thresh the standing crops in the fields in the morning/evening to protect them from the heat wave. Do not do any work on the fields between 11 am and 4 pm and do light irrigation in the standing crops in the evening at an interval of 10-12 days. Do not leave the animals outside for grazing between 11 am and 4 pm. Bathe the animals in the morning and evening.

Farmers are advised to download Unified "Mausam" and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: https://play.google.com/store/apps/details/

Meghdoot MobileApp link: https://play.google.com/store/apps/details

Damini MobileApp link: https://play.google.com/store/apps/details