



Gramin Krishi Mausam Sewa
Agromet Advisory Bulletin for UNNAO District
Chander Shekhar Azad University of Agriculture &
Technology
Kanpur, Uttar Pradesh



Agromet Advisory Bulletin

Date : 2026-07-10

Weather Forecast of District UNNAO (Uttar Pradesh) Issued On : 2026-07-10 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-07-11	2026-07-12	2026-07-13	2026-07-14	2026-07-15
Rainfall(mm)	54.0	17.0	2.0	3.0	0.0
Tmax(°C)	32.0	34.0	37.0	36.0	36.0
Tmin(°C)	26.0	27.0	29.0	27.0	28.0
RH-I(%)	86	79	75	77	81
RH-II(%)	59	54	45	48	62
Wind Speed(kmph)	8	16	21	3	9
Wind Direction(Degree)	211	262	252	194	100
Cloud Cover(Octa)	6	7	6	6	7
Warning	Heavy Rain; Thunderstorms & Lightning, Squall etc	Heavy Rain	No Warning	No Warning	No Warning

Forecast Summary:

According to the weather forecast received from the Indian Meteorological Department, due to moderate to dense cloud cover in the next five days there is a possibility of moderate to heavy rain with local level high wind speed, thunderstorm– lightning between dated on 11-14 July, 2026. The maximum temperature is between 32.0- 37.0°C, which is likely to be near its normal and the minimum temperature is between 26.0-29.0°C, which is likely to be 2-3°C higher normal. The maximum and minimum range of relative humidity is between 75-86 and 48- 62%. Wind direction is south-west, south-east and wind speed is between 8.0-21.0 kmph with gusts expected to be 7-8 kmph high speed than normal.

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the India Meteorological Department (IMD), there is a warning of thunderstorms, strong winds, and moderate to heavy rain in upcoming days . Humid weather is also expected to continue.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Carry out sowing, transplanting, and other agricultural activities when there is no rain and the sky is clear. Apply fertilizers, insecticides, and fungicides only when the weather clears up. Regularly monitor crops for pests and diseases after the rain. Keep livestock in a dry, safe area and provide them with clean drinking water. Do not tether animals under trees or near electricity poles, as falling branches or electric shocks from the poles could prove fatal to the animals.

General Advisory:

Farmers are advised to transplant ready paddy seedlings and, in the absence of rain, proceed with the sowing of Kharif crops such as maize, groundnut, sorghum (jowar), sesame, and pigeon pea (arhar). If possible, harvest mature crops before the rain starts and store the harvested produce in a dry, safe place. Regularly monitor crops for pests and diseases after the rain. Do not tether animals under trees or near electricity poles. Bathe the animals in the morning and evening, keep them in a dry, safe area, and provide clean drinking water. Use clean water for insecticides, fungicides, and herbicides, and do not spray them while standing against the wind. If possible, wash your hands thoroughly with soap after spraying, before eating, and after washing clothes.

SMS Advisory:

Farmers are advised to transplant ready paddy seedlings and, in the absence of rain, proceed with the sowing of crops such as groundnut, sorghum, sesame, pigeon pea, black gram and green gram.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
RICE (Transplanting)	Make proper arrangements for the drainage of excess rainwater from the paddy nursery. In case of excessive rainfall, postpone the transplanting work. Prepare the field for transplanting paddy saplings and transplant 22-25 days old paddy saplings. Strengthen the bunds of the fields for transplanting paddy saplings. So that rainwater remains in the fields. For control of broad and narrow leaf weeds in paddy crop, anilophos 30% EC is applied within 2-3 days of transplanting. Or prittilechlor at the rate of 1.25 liters / ha and make a solution in 500 -600 liters of water and spray 2-3 inches of water. If Khaira disease is visible in paddy, then spraying 20-25 kg zinc sulphate and 2.5 kg lime in 800 liters of water is sprayed for its control.
MAIZE (Germination)	Make proper arrangement for drainage of excess rain water. In case of no rain. In the maize crop sown from time to time, do first weeding - hoeing after 15-20 days. sow one of the recommended hybrid varieties of Kharif maize- D-7074, D-9144, M-2525, Ganga, Azad Shekhar Makkah-1, Azad Shekhar Makkah-2, Azad Uttam and Azad Kamal etc.
SORGHUM (JOWAR/GREAT MILLET) (Sowing)	In view of the possibility of rain, farmers should postpone sowing work. sow any one of the recommended hybrid varieties of sorghum - Varsha, CSV-13, CSV-15, SPV-1388 (Bundela), Vijayeta and hybrid varieties - CSH-16, CSH-9, CSH-14, CSH-18, CSH-13 and CSH-23 etc. when the sky is clear.
GROUNDNUT (Sowing)	In view of the possibility of rain, farmers should postpone sowing. Sow recommended hybrid varieties of kharif groundnut - Chitra, Kaushal, Prakash, Amber, Utkarsh, Divya, Kaushal - bunched variety - Chandra, Ta-28, 64 M-13 etc. And for sowing peanuts, sow at the rate of 70-80 kg / hectare when the sky is clear.
SESAME (GINGELLY/TIL) (Sowing)	In view of the possibility of rain, farmers should postpone sowing. Sow any one of the recommended varieties of sesame crop - Type-4, 12, 13, Type-78, Shekhar, Pragati, Tarun, RT-351 and RT-346 etc. when the sky is clear.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
CAULIFLOWER (Nursery/ seedling)	Make proper arrangements for excessive rain water drainage. In case of no rain, do sowing/planting/weeding-hoeing work. Raise nurseries of crops such as brinjal, chili, tomato and early cauliflower etc. to be planted in Kharif, as well as transplant prepared seedlings. For prevention of fruit borer/leaf borer

Horticulture (Stage)	Horticulture Specific Advisory
	pest in vegetable crops, prepare a solution of 1.5-2.0 ml/liter of water with neem oil and spray it 3-4 times at an interval of 8-10 days. Weeding-hoeing should be done in standing vegetable crops when sky is clear.
MANGO (Harvesting)	For filling the pits dug in the month of May for planting fruit orchards, mix rotten manure of cow dung in the upper half of the soil and fill it up to 15 to 20 cm above the ground level. To protect mango and guava from fruit flies, hang 8-10 traps per hectare using methyl eugenol and Q lure traps tied to branches at a height of 6 to 8 feet. Spray 5% neem extract dissolved in one litre of water at an interval of 10 to 15 days. Change the lure at an interval of 20-25 days.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	Farmers are advised to give sufficient quantity of grains with green and dry fodder to the animals. Take special care of calving animals and protect lactating animals from milk fever, ketosis etc. after calving. Protect animals and newborns from internal and external parasites. Use disinfectant medicine in the cattle shed. The best time to give antithetical drugs for the prevention of insects in the stomach of animals. Get the animals vaccinated with throat and lenthia fever. The animals must give clean and fresh water 3-4 times a day. Give the animals a bath two to three times a day. Tie the animals in the shade at the time of day and do not tie the animals under the tree. Because this week due to the high wind speed, there is a high probability of falling of trees / tree twigs.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Farmers are advised to add supplements, vitamins and energy food items to the food of chickens and also give calcium to the chickens. Give medicine to prevent worms in the stomach of chickens (deworming).

Likely impacts of weather warnings (General)

Heavy rainfall may lead to flooding and waterlogging in low-lying areas. Road, bridge, and other transport networks could be disrupted. There is a risk of serious injury or death due to lightning strikes, including a threat to livestock grazing in open areas; power supply and communication services may be affected; there is a risk of mosquito-borne diseases such as malaria, dengue, and chikungunya; and eye and respiratory issues may increase due to airborne dust and debris.
--

Impact based advisories (General)

Regularly follow weather department warnings and avoid unnecessary agricultural activities or travel during thunderstorms, lightning, gusty winds, or heavy rainfall. Apply fertilizers and plant protection chemicals only after the rain has stopped, and ensure the prompt drainage of excess water from fields. Keep livestock in dry, elevated areas; do not seek shelter under trees, and stay away from electric poles and water bodies. Provide clean drinking water and dry fodder, Protect green fodder from getting wet to prevent fungal growth. Keep animal shelters clean, and ensure immediate treatment for sick animals. Consume clean and boiled water, prevent water stagnation, use mosquito nets, and maintain personal hygiene.

Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: <https://play.google.com/store/apps/details/>

Meghdoot MobileApp link: <https://play.google.com/store/apps/details>

Damini MobileApp link : <https://play.google.com/store/apps/details>