



Gramin Krishi Mausam Sewa
Agromet Advisory Bulletin for KASGANJ District
Chander Shekhar Azad University of Agriculture & Technology
Kanpur, Uttar Pradesh



Agromet Advisory Bulletin

Date : 2026-07-17

Weather Forecast of District KASGANJ (Uttar Pradesh) Issued On : 2026-07-17 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-07-18	2026-07-19	2026-07-20	2026-07-21	2026-07-22
Rainfall(mm)	1.0	9.0	14.0	26.0	26.0
Tmax(°C)	38.0	36.0	34.0	32.0	32.0
Tmin(°C)	29.0	29.0	28.0	26.0	26.0
RH-I(%)	73	76	82	88	88
RH-II(%)	43	51	58	62	63
Wind Speed(kmph)	4	3	1	12	12
Wind Direction(Degree)	243	325	154	119	104
Cloud Cover(Octa)	7	7	7	7	8
Warning	Thunderstorms & Lightning, Squall etc; Hot Day	No Warning	Thunderstorms & Lightning, Squall etc	Heavy Rain; Thunderstorms & Lightning, Squall etc	Heavy Rain; Thunderstorms & Lightning, Squall etc

Forecast Summary:

According to the weather forecast received from the India Meteorological Department, due to moderate to dense cloud cover in the next five days there is a possibility of moderate to heavy rain with local level thunderstorm– lightning between dated on 18 -22 July, 2026. The maximum temperature is between 32.0- 38.0°C, which is likely to be 1-2°C above normal and the minimum temperature is between 26.0-29.0°C, which is likely to be 2-3°C above normal. The maximum and minimum range of relative humidity is between 73-88 and 43- 63%. Wind direction is south-east, south-west and wind speed is between 1.0-12.0 kmph with gusts expected to be 2-3 kmph high speed than normal

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the India Meteorological Department (IMD), there is a warning of thunderstorms, strong winds, moderate to heavy rain. Humid weather is also expected to continue.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Carry out sowing, transplanting, and other agricultural activities when there is no rain and the sky is clear. Apply fertilizers, insecticides, and fungicides only when the weather clears up. Regularly monitor crops for pests and diseases after the rain. Keep livestock in a dry, safe area and provide them with clean drinking water. Do not tether animals under trees or near electricity poles, as falling branches or electric shocks from the poles could prove fatal to the animals.

General Advisory:

Farmers are advised to complete the transplanting of ready paddy seedlings as soon as possible and, in the absence of rain, to proceed with the sowing of crops such as groundnut, sorghum, sesame, black gram (urd), green gram (moong), and pigeon pea (arhar). Regularly monitor crops for pests and diseases following rainfall. Do not tether livestock under trees or near electric poles. Bathe the animals in the morning and evening, keep them in a dry and safe location, and provide them with clean drinking water. Use clean water for preparing pesticides, fungicides, and herbicides, and avoid spraying these chemicals while standing against the wind direction. If possible, wash hands thoroughly with soap after spraying, before eating, and after washing clothes.

SMS Advisory:

Farmers are advised to complete the transplantation of ready paddy seedling soon in view of the possibility of rain and in case of no rain, they should sow crops like sesame, urd, green gram etc.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
RICE (Tillering)	Complete the transplanting of ready 20-25 days old paddy seedlings as soon as possible. For the control of broad and narrow leaf weeds in paddy crop, Enilophos 30% EC within 2-3 days of transplanting. Or sprinkle 2-3 inches of water by making a solution of Pritalchlor 1.50 liters/ha in 500-600 liters of water. After 15 - 20 days of transplanting of paddy, apply bispyribac sodium 10 SC at the rate of 0.20 liters/ha in 500-600 liters of water in proper moisture condition. If Khaira disease is visible in paddy, then to control it, make a solution of 20-25 kg zinc sulphate and 2.5 kg lime in 800 liters of water and spray.
MAIZE (Vegetative)	In case of no rain, weeding and urea top dressing of maize crop should be done only when sky is clear. There is a possibility of stem borer and leaf roller pest infestation in the maize crop, so for its prevention, spray Emamectin Benzoate 5% SG 200 grams/hectare or Dimethoate 30% EC 1.0 liter/hectare by mixing it in 600-700 liters of water in clear weather.
SORGHUM (JOWAR/GREAT MILLET) (Sowing)	In view of the possibility of rain, farmers should postpone sowing work. sow any one of the recommended hybrid varieties of sorghum - Varsha, CSV-13, CSV-15, SPV-1388 (Bundela), Vijayeta and hybrid varieties - CSH-16, CSH-9, CSH-14, CSH-18, CSH-13 and CSH-23 etc. when the sky is clear.
GROUNDNUT (Sowing)	In view of the possibility of rain, farmers should postpone sowing. Sow recommended hybrid varieties of kharif groundnut - Chitra, Kaushal, Prakash, Amber, Utkarsh, Divya, Kaushal - bunched variety - Chandra, Ta-28, 64 M-13 etc. And for sowing peanuts, sow at the rate of 70-80 kg / hectare when the sky is clear.
SESAME (GINGELLY/TIL) (Sowing)	In view of the possibility of rain, farmers should postpone sowing. Sow any one of the recommended varieties of sesame crop - Type-4, 12, 13, Type-78, Shekhar, Pragati, Tarun, RT-351 and RT-346 etc. when the sky is clear.
BLACK GRAM (Sowing)	Recommended species of urd to be sown in Kharif - Narendra urd-1, Azad urd-2, Azad urd-3, Shekhar-1, Shekhar-2, Shekhar-3, Pant U-30 etc. For this, by arranging fertilizers and seeds at the rate of 12 -15 kg seed / hectare, sowing in case of no rain.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
BRINJAL (Transplanting)	Make proper arrangements for excessive rain water drainage. In case of no rain, do sowing/planting/weeding-hoeing work. Raise nurseries of crops such as brinjal, chili, tomato and early cauliflower etc. to be planted in Kharif, as well as transplant prepared seedlings. For prevention of fruit borer/leaf borer pest in vegetable crops, prepare a solution of 1.5-2.0 ml/liter of water with neem oil and spray it 3-4 times at an interval of 8-10 days. Weeding-hoeing should be done in standing vegetable crops when sky is clear.
MANGO (Planting)	Perform the planting of new plants in the pits already filled with mangoes. For prevention of mango leaf moth, apply 2% carbaryl 1.5% dust on the clear sky. After weeding the banana/papaya garden, put 25-30 cm high soil around the stem and do the work of standing in banana plants.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	Farmers are advised to give sufficient quantity of grains with green and dry fodder to the animals. Take special care of calving animals and protect lactating animals from milk fever, ketosis etc. after calving. Protect animals and newborns from internal and external parasites. Use disinfectant medicine in the cattle shed. The best time to give antithetical drugs for the prevention of insects in the stomach of animals. Get the animals vaccinated with throat and lenthia fever. The animals must give clean and fresh water 3-4 times a day. Give the animals a bath two to three times a day. Tie the animals in the shade at the time of day and do not tie the animals under the tree, because this week due to the high wind speed, there is a high probability of falling of trees / tree twigs.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Farmers are advised to add dietary supplements; vitamins and energy feed ingredients to the hens as well as add calcium content to the hen. Give medicine for the prevention of worms in the stomach of hen (Divmigr).

Likely impacts of weather warnings (General)

Sultry heat (high humidity and temperature) increases the humidity in the air, making it difficult for sweat to dry easily, leading to discomfort. Humidity can increase the risk of dehydration and skin problems like prickly heat and fungal infections. Constant humidity can also lower energy levels, increase irritability, and cause insomnia, leading to mental stress. Physical discomfort and fatigue affect people's performance at work and in studies. The humidity associated with the rainy season promotes rapid mosquito and bacterial growth, increasing the risk of malaria, dengue, and viral infections.

Impact based advisories (General)

Drink water every hour, even if you don't feel thirsty. Mix lemon water, ORS, or a salt-sugar solution with water. Eat light meals and water-rich fruits like cucumber, radish, and oranges. Limit tea, coffee, alcohol, and spicy or fried foods. Wear light-colored, loose-fitting cotton clothing that allows for ventilation. To avoid heat rash and fungal infections, shower twice daily and use anti-fungal powder. Open the windows to allow air to circulate, reducing indoor humidity. If you're using a cooler, keep the window or door slightly open to allow moisture to escape, otherwise the humidity will increase.

Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: <https://play.google.com/store/apps/details/>

Meghdoot MobileApp link: <https://play.google.com/store/apps/details>

Damini MobileApp link : <https://play.google.com/store/apps/details>