Gramin Krishi Mausam Sewa

Agromet Advisory Bulletin for KANNAUJ District

Chander Shekhar Azad University of Agriculture & Technology

Kanpur, Uttar Pradesh

Agromet Advisory Bulletin

Date : 2026-10-30

Weather Forecast of District KANNAUJ (Uttar Pradesh) Issued On : 2026-10-30 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-10-31	2026-11-01	2026-11-02	2026-11-03	2026-11-04
Rainfall(mm)	10.0	0.0	0.0	0.0	0.0
Tmax(°C)	24.0	25.0	29.0	31.0	30.0
Tmin(°C)	21.0	20.0	20.0	19.0	19.0
RH-I(%)	91	89	89	80	75
RH-II(%)	73	70	53	38	39
Wind Speed(kmph)	16	11	5	1	1
Wind Direction(Degree)	71	33	282	252	18
Cloud Cover(Octa)	8	4	1	1	0
Warning	No Warning	No Warning	No Warning	No Warning	No Warning

Forecast Summary:

According to the weather forecast received from the Indian Meteorological Department, due to moderate to dense cloud cover in the next four day there is a possibility of light to moderate rain with thunderstorms & lightning at local level between dated on 31 Oct, 2026. The maximum temperature is between 24.0- 31.0°C, which is likely to be 1-2°C lower normal and the minimum temperature is between 19.0-21.0°C, which is likely to be 3-4°C higher normal. The maximum and minimum range of relative humidity is between 75-91 and 38-73%. Wind direction is north-east, north-west and wind speed is between 1.0-16.0 km/h with winds expected to be 3-4 Km/h normal.

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast of the India Meteorological Department, a warning of light to moderate rain with strong winds, thunderstorm and lightning has been issued.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Heavy rainfall and strong winds can cause standing or harvested paddy crops to fall over. This increases moisture content in the grains, potentially causing them to turn black and damaging the harvest. Rain immediately after sowing can form a crust on the soil surface, which may severely hinder the germination of mustard, potato, and early-sown wheat seeds. High humidity and persistent cloud cover can lead to fungal growth on harvested crops, as well as an increased risk of root rot and blast disease in vegetables and pulse crops. If the paddy crop is fully mature, postpone harvesting. Crops that have already been harvested should be moved from the fields to a safe, elevated location or covered thoroughly with tarpaulins. For fields prepared for mustard, early-sown wheat, and potatoes, wait until after the rain once the soil is dry enough to be worked before proceeding with sowing. In fields used for vegetables, peas, and pulses, ensure that a system of ridges and furrows (or drainage

General Advisory:

In view of the potential weather changes or light rain expected in the coming days, please move harvested crops from the fields to safe, elevated locations, or cover them thoroughly with tarpaulins. Farmers sowing mustard, potatoes, or early-sown wheat should ensure appropriate soil moisture and proceed with sowing only once the weather has completely cleared. Avoid heavy irrigation, as well as the application of pesticides or fertilizers, for the next 2–3 days to prevent the wastage of water and unnecessary expenditure. To protect milch animals from dry, cold north-westerly winds, cover their shelters (enclosures) at night using tarpaulins or jute sacks.

SMS Advisory:

Farmers are advised to ensure that the weather is favorable for harvesting/threshing of standing Kharif crops and sowing of Rabi crops like wheat, gram, peas, mustard, linseed, potatoes and vegetables

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
RICE (Maturity and Harvesting)	Harvest the paddy crops and after drying for 2 -3 days after harvesting, and after harvesting, dry the sun for 3-4 days and store the seeds.
GROUNDNUT (Maturity and Harvesting)	Dig the groundnut crops only when the veins above the peel become visible and the inner part becomes brown and the peanut grain turns pink. After digging peanuts, dry the pods thoroughly in the sun and store them.
WHEAT (Sowing)	According to the weather, the temperature is favorable for sowing of wheat in the irrigated condition, so the sowing of wheat should start from 10 November. Prepare field for sowing wheat and regional species PBW-723, HPBW-01, DBW-39, K.-9107, K.-1006, K.-402, K.-607, D. .BW-90, WB-02, NW-5054, HD-3043, U.P-2382, H.U. W-468, PBW-443, HD-2967 etc. Recommendation species for erosive land: - KRL-210, KRL-213, K.- 1317 etc. for the sowing of any species of manure and seeds should be arranged.
RAPESEED (Sowing)	There is a possibility of infestation of hairy caterpillar in the rapeseed crop, so for its prevention, Emamectin benzoate 5% SG at the rate of 200 g / ha is sprayed in 500-600 liters of water. Within 20-22 sowing of rapeseed crop, weeding should be done within 10-22 centimeters distance from plant to plant. In the rapeseed crop, the first irrigation should be done within 25-30 days of sowing and on arrival of oats, make top dressing of urea at the rate of 108 kg / ha.
MUSTARD (Sowing)	Saw fly in mustard crop - These pests cause more damage in 7 to 10 days of germination. To prevent this, methyl parathion 2 percent or malathion 5 percent or carbaryl 5 percent powder at the rate of 25 kg per hectare in the morning or evening. It on clear weather for sowing work at the rate of 3-4 kg seed/ha for sowing any one of the recommended varieties of rye/mustard- Pusa Mustard-34, Pusa Mustard-32, Varuna (V-WRR-2), C5-64 (CS2005-143), Pusa OOM-36 (PDZ-15), BMP-II (DIMR), Azad Mahak, Surekha (KMR-16-02), Rohini, Narendra Rai- 8501, Maya, Vaibhav etc.
FIELD PEA (Sowing)	For sowing of any of pea crop recommended Variety - rachana, Malaviya pea-15, Sapna etc., sow at the rate of 100-125 kg seed / hectare.
CHICK PEA (Sowing)	For sowing of any varieties of chickpea, crop recommendation species- Awarodhi, Pusa-256, Pusa-362, K.W.R-108, K-850, K.D.G.-1168 etc. Small grain seeds of 75-80 kg/ hectare for large grain seeds, sow at the rate of 90-100 kg of seed / hectare.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
POTATO (Sowing)	Main varieties of potato - Kufri Bahar, Kufri Anand, Kufri Badshah, Kufri Sinduri, Kufri Sutlej, Kufri Evergreen Lalima, Kufri Arun, Kufri and Kufri Pukhraj, Kufri Garim, Kufri Lima, Kufri Lalit, Kufri Sangam, Kufri Neelkanth, Kufri Ganga, Kufri Thar-3, Kufri Mohan, Kufri Neelkanth, Kufri Surya, Kufri Chipsona-1, Kufri Chipsona-3, Kufri Chipsona-4 And the weather is favorable for sowing of Kufri, Friesona etc. To prevent chicken pox in potato crop, spray 100 grams of streptocycline or boric acid in 10 liters of water and spray the tubers on the seeds and do not treat the germinated seeds.

Horticulture (Stage)	Horticulture Specific Advisory
VEGETABLE PEA (Sowing)	Sow vegetables like peas, fennel, celery, beetroot, garlic, coriander, spinach, soya, fenugreek, radish, carrot, turnip and leafy vegetables. Raise nursery of tomato, onion, chilli, cauliflower and cabbage. If there is a possibility of water stagnation in the field, then plant cauliflower, brinjal, tomato, chilli etc. on the ridges. If there is an infestation of fruit borer/leaf borer insect in vegetable crops, then for its control, make a solution of neem oil in 1.5 ml/litre water and spray it 3-4 times at an interval of 8-10 days.
MANGO (Vegetative)	Plant new plants in place of destroyed plants in new orchards of mango, guava, amla, lemon etc. After hoeing the banana/papaya gardens, add 25-30 cm high soil around the stem. Install wooden or boss stands in banana/papaya fruit trees. To prevent leaf cutting insect in mango, sprinkle 2% carbaryl 1.5% dust.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	Farmers are advised not to tie pregnant buffaloes/cows at sloping places. Feed nutritious fodder and grains to pregnant buffaloes/cows and feed colostrum to the newborn baby for three days. Keep animals in a clean place. Be sure to vaccinate animals to protect them from strangulation. In view of the changing weather, do not tie animals in open places at night and ensure to take care of the health and illness of the animals. Provide adequate amount of grains along with green and dry fodder to the animals. Make sure to feed clean and fresh water to animals 3-4 times a day.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Farmers are advised to Clean every day at the place where the chickens live. Arrange for clean water and balanced diet to the hens. This is not the right time to raise new chicks for eggs but it is a suitable time for broiler chickens. Add supplementary food, vitamins and energy food material to the chickens' diet and also give calcium material to the chickens. Chlorination of water must be done. Give deworming medicine to kill the worms in the stomach of the chickens. Keep the chicken house dry and keep turning the bedding.

Likely impacts of weather warnings (General)

Heavy rainfall impacts human health, damages infrastructure, and causes complete disruption to transportation. Lightning strikes, flash floods, and landslides pose a direct threat to human life and property. During the rainy days, damp conditions in animal shelters can lead to hoof infections, while the consumption of wet fodder can cause serious gastrointestinal ailments in livestock. Unseasonal heavy rain and hailstorms during the harvest period cause standing crops to flatten in the fields, where they subsequently rot. Sudden shifts in weather or high humidity levels accelerate the proliferation of pests and fungal infections in crops. The loss of a harvest results in the total forfeiture of the capital invested by farmers in seeds and fertilizers, trapping them in financial distress and debt. Following weather warnings, farmers face increased operational costs associated with managing drainage and relocating crops to safer areas. Furthermore, produce—such as grains and vegetables—that suffers a decline in quality due to adverse weather conditions fails to fetch fair prices in the markets.

Impact based advisories (General)

In light of alerts for heavy rain and lightning, stay indoors in sturdy, permanent structures during thunderstorms; avoid standing under trees, near weak walls, or beneath electricity poles. Expect waterlogging on roads and reduced visibility; therefore, avoid non-essential travel and do not drive through waterlogged areas. To protect livestock from strong winds and rain, keep them in well-ventilated, permanent shelters rather than tying them in open fields or under trees. Ensure that the floors of animal sheds remain dry and clean to prevent foot infections, and avoid feeding livestock wet or moldy fodder. As the harvesting of Kharif crops (such as paddy) is currently underway, suspend all harvesting activities until the weather clears. Ensure proper drainage in fields to prevent water

accumulation, which could cause crop roots to rot. Move any already-harvested crops to a safe, elevated location, or cover them securely with tarpaulins to prevent the grains from darkening. Given the c

Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: https://play.google.com/store/apps/details/

Meghdoot MobileApp link: https://play.google.com/store/apps/details

Damini MobileApp link : https://play.google.com/store/apps/details