



Gramin Krishi Mausam Sewa
Agromet Advisory Bulletin for KANPUR-DEHAT District
Chander Shekhar Azad University of Agriculture & Technology
 Kanpur, Uttar Pradesh



Agromet Advisory Bulletin

Date : 2026-10-01

Weather Forecast of District KANPUR-DEHAT (Uttar Pradesh) Issued On : 2026-10-01 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-10-02	2026-10-03	2026-10-04	2026-10-05	2026-10-06
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	34.0	36.0	36.0	35.0	34.0
Tmin(°C)	23.0	24.0	24.0	25.0	26.0
RH-I(%)	75	69	66	68	71
RH-II(%)	46	43	43	45	52
Wind Speed(kmph)	7	6	8	8	6
Wind Direction(Degree)	279	228	253	235	220
Cloud Cover(Octa)	0	0	0	0	2
Warning	No Warning	No Warning	No Warning	No Warning	No Warning

Forecast Summary:

According to the weather forecast received from the Indian Meteorological Department, due to clear sky weather in the next four day and light cloud cover in remaining day but there is a no possibility of rain. The maximum temperature is between 34.0- 36.0°C, which is likely to be 2-3°C higher normal and the minimum temperature is between 23.0-26.0°C, which is likely to be 1-2°C higher normal. The maximum and minimum range of relative humidity is between 66-75 and 43-52%. Wind direction is south-west and wind speed is between 6.0-8.0 km/h with winds expected to be 2-3 Km/h normal.

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the India Meteorological Department (IMD), there is a no warning.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Farmers are advised that weather conditions are favorable for the harvesting, threshing, and sowing of crops. Do not tether livestock in open areas or let them out to graze, and prevent them from entering waterlogged areas. Keep animals in a dry, safe place and provide them with clean drinking water; do not tie them under trees or near electric poles, as falling branches or electric shocks from the poles could prove fatal.

General Advisory:

Weather conditions are favorable for the harvesting, threshing, and sowing of mature crops. If harvested crops in the field have gotten wet, dry them thoroughly in the sun before proceeding with threshing. Prevent livestock from entering areas with flowing water or waterlogging, and do not allow them to drink stagnant rainwater. When stepping out of the house at night during the rainy season, carry a torch; to protect against snakes, scorpions, and similar creatures, seal gaps and holes in the walls and spray a small amount of kerosene or phenyl, as the odor repels snakes.

SMS Advisory:

There is no possibility of rain in the coming week; therefore, the weather is favorable for the harvesting, threshing, and sowing of mature crops.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
RICE (Millking)	The rice crop is in the stage of emergence of ears and flowering/milking, which is sensitive to moisture deficiency, so in the absence of rain, maintain proper moisture by doing light irrigation as per requirement. The current weather is favorable for pests, so on an average, 2 - 3 odorous pests / hill are visible in the paddy crop after flowering, then for its control, Imidacoloroprid 17.8% SL 350 ml / ha or Burfozin 25% SP 750 ml Make a solution in 500-600 liters of water at the rate of / hectare and spray it on clear weather. If the outbreak of stem borer insect is seen in the paddy crop, then Imidacloprid 17.8 EC for its control. SL at the rate of 25 g/ha make a solution in 500-600 liters of water and spray it in clear weather.
SESAME (GINGELLY/TIL) (Harvesting)	Weather conditions are favorable for the harvesting and threshing of mature crops. If harvested crops in the field have gotten wet, dry them thoroughly in the sun before proceeding with threshing. There is a possibility of showing an outbreak of leaf and pod larva pest in sesame crop, so for its prevention, Emaxectin Benzoate 5 % SG 200 g / hectare or Quinalphos 25 % EC at the rate of 1.25 liters / hectare make a solution in 500-600 liters water and spray on clear weather.
GROUNDNUT (Physiological Maturity)	There is a likelihood of white grub or termite infestation in the groundnut crop; therefore, to control this, apply Fipronil 0.3% GR by broadcasting at a rate of 20 kg/hectare.
BLACK GRAM (Physiological Maturity)	To prevent the pod borer larvae in standing black gram/green gram crops, which bore into the pods and eat the grains, spray 750 ml of Cypermethrin 10 EC per hectare by dissolving it in 500-600 liters of water in clear weather.
RAPESEED (Sowing)	For sowing any one of the recommended varieties of rapeseed - Ta-36, Ta-9 Bhavani, PT-303, PT-30 etc., arrangement of fertilizer and seeds at the rate of 3-4 kg seeds/hectare do the sowing work .
SUGARCANE (Grand growth)	If the length of sugarcane has become five feet, then do the first tying at a height of two and half feet. For the first tying, tie two bunches of sugarcane in a line together with the dry leaves below. Destroy the plants affected by red rot disease by removing them along with the roots and spray the plants with a solution of thiophanate methyl 0.2 percent. If symptoms of bacterial rot disease are seen in sugarcane crop, cut and destroy the infected plant and for its control, spray 0.01% Streptomycin when weather is favorable.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
POTATO (Sowing)	Start sowing of early varieties of potato like Kufri Chandra Mukhi and Kufri Ashoka. Take out the potato seeds from cold storage 7-10 days before sowing. If sprouts appear in the seeds stored in cold storage, then sort them and separate them. If the potato seeds have not been treated before storing them in the cold storage, then immediately after taking out the potatoes from the cold storage and sorting them, treat the potato tubers with 3 percent boric acid solution for 30 minutes and dry them in a shady place.
CHILLI (Transplanting)	Sow radish, carrot, beetroot, turnip and leafy vegetables on the ridges. Transplanting of saplings planted brinjal, chilli, tomato and early cauliflower. If there is a possibility of water stagnation in the field, then plant the early cauliflower, brinjal, tomato, chilli etc. on the ridges. If there is an outbreak of fruit borer / leaf borer in vegetable crops, make a solution of neem oil in 1.5 ml / liter of water to control it and spray it 3-4 times after clearing the sky at intervals of 6-10 days.
MANGO (Planting)	Try to complete the planting of fruit plants like mango, guava, amla, lemon etc. as soon as possible. Plant new plants in place of the destroyed plants in the new bag. Make a stand of 25-30 meters high soil around the stem by folding banana / papaya bags. For prevention of mango leaf moth, apply 2% carbaryl 1.5% dust on the clear sky.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	Tie animals in the open only when the sky is clear at night. Tie animals in a shady place during the day. Do not feed the stagnant water of the ponds or other places, due to which the animals are prone to liver flu and stomach worms. The best time to give antithetical drugs for the prevention of insects in the stomach of animals. Give sufficient amount of grains to animals along with green and dry fodder. Make sure to give clean and fresh water to animals 3-4 times a day. Do not tie pregnant animals at a sloping place. This is the right time to give deworming medicine to animals to prevent stomach worms.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Farmers are advised to Clean every day at the place where the chickens live. Arrange for clean water and balanced diet to the hens. This is not the right time to raise new chicks for eggs but it is a suitable time for broiler chickens. Add supplementary food, vitamins and energy food material to the chickens' diet and also give calcium material to the chickens. Chlorination of water must be done. Give deworming medicine to kill the worms in the stomach of the chickens. Keep the chicken house dry and keep turning the bedding. If there is excess moisture in the chicken house, turn on the fan.

Likely impacts of weather warnings (General)

Prolonged exposure to sunlight and high humidity significantly increase the risk of heat stress and infectious diseases in livestock. High humidity and heat reduce the appetite of buffaloes and cows, potentially leading to a sudden drop in milk production. Stagnant water following recent monsoon rains, combined with rising temperatures, creates an environment where bacteria thrive in fodder and water sources. This can lead to the spread of ailments such as intestinal worms, skin infections, and Foot-and-Mouth Disease among the animals. The combination of a sudden rise in humid heat and a gradual drop in minimum temperatures creates conditions characterized by intense heat during the day and a slight chill at night or in the early morning. This fluctuating weather affects human health; venturing out in strong sunlight can lead to dehydration. Furthermore, this season sees a rapid surge in the prevalence of mosquitoes and viruses responsible for diseases like Chikungunya, Dengue, Malaria, and viral flu.

Impact based advisories (General)

Do not tether animals in direct sunlight during the day (between 11 AM and 4 PM); keep them in a well-ventilated, shaded shed instead. Ensure drinking water troughs are kept clean and provide fresh, clean water at least 3-4 times a day. If vaccinations for Hemorrhagic Septicemia (HS) or Black Quarter (BQ) have not been administered, get the animals vaccinated. At night, use neem leaf smoke in the animal shed to protect against mosquitoes and flies. Carry a water bottle when stepping out. Consume ORS, lemonade, or buttermilk to prevent electrolyte depletion in the body. Do not allow rainwater from recent showers to stagnate around the house or on rooftops. Use mosquito nets or repellents while sleeping. Avoid drinking cold water from the refrigerator or sitting directly in air-conditioned (AC) environments immediately after coming in from the heat; these habits are primary causes of colds, coughs, and lung infections. This dry week is ideal for preparing Rabi crops and for the harvesting

Farmers are advised to download Unified "Mausam" and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: <https://play.google.com/store/apps/details/>

Meghdoot MobileApp link: <https://play.google.com/store/apps/details>

Damini MobileApp link : <https://play.google.com/store/apps/details>

