



Gramin Krishi Mausam Sewa
Agromet Advisory Bulletin for SITAPUR District
Chander Shekhar Azad University of Agriculture & Technology
 Kanpur, Uttar Pradesh



Agromet Advisory Bulletin

Date : 2026-05-05

Weather Forecast of District SITAPUR (Uttar Pradesh) Issued On : 2026-05-05 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-05-06	2026-05-07	2026-05-08	2026-05-09	2026-05-10
Rainfall(mm)	15.0	3.0	12.0	0.0	0.0
Tmax(°C)	29.0	31.0	32.0	33.0	35.0
Tmin(°C)	20.0	21.0	23.0	23.0	24.0
RH-I(%)	89	86	78	76	74
RH-II(%)	60	56	46	53	51
Wind Speed(kmph)	10	4	12	18	21
Wind Direction(Degree)	63	360	360	104	106
Cloud Cover(Octa)	6	5	6	1	3
Warning	Thunderstorms & Lightning, Squall etc; Strong Surface Winds	No Warning	Thunderstorms & Lightning, Squall etc	No Warning	No Warning

Forecast Summary:

According to the weather forecast from the India Meteorological Department, due to light to moderate cloud cover in the next five days, there is a possibility of light to moderate rain with local level at high winds speeds, thunderstorms & Lightning, hailstorm between dated on 06-08 May, 2026. The maximum temperature is between 29.0-35.0°C, which is likely to be 3-4°C lower normal and the minimum temperature is between 20.0-24.0°C, which is likely to be near its normal. The maximum and minimum range of relative humidity is between 74-89 and 46-60%. The wind direction will be from the north-east, south-east, north and the wind speed will be between 4.0-21.0km/hour, which is expected to be 6-7 km/hour higher than normal.

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the India Meteorological Department, a warning has been issued for the period between May 06-08 May , 2026, alerting to light to moderate rainfall, thunderstorms accompanied by lightning, strong surface winds, and hailstorms.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

In view of the weather conditions, farmers working in the fields are advised not to take shelter or allow themselves or their animals to take shelter under trees in case of heavy winds/storms or rain. Suspend irrigation activities for standing crops such as summer maize, black gram (Urad), and green gram

(Moong). Furthermore, harvest mature crops such as wheat and pigeon pea (Arhar) and store them in a safe location.

General Advisory:

Farmers are advised to make proper arrangements for excess rain water drainage. Harvested crops should be collected and covered with polythene sheets. The grains should be dried thoroughly in the sun and stored safely. Farmers are advised to maintain proper moisture in standing vegetables and crops by doing light irrigation as per requirement. While working in the field, tie a towel on the head and keep a water bottle with you. Keep the threshed crop grains in a safe place. In view of the severe heat, give water to the animals 3-4 times a day and bathe them in the morning and evening. Use separate equipment for insecticides, fungicides and weedicide chemicals or wash the equipment with clean water and do not spray insecticides, fungicides and weedicide chemicals while standing in the opposite direction of the wind. If possible, spray in the evening, after spraying, wash hands thoroughly with soap or hand wash before eating or drinking and wash clothes and take a bath.

SMS Advisory:

Farmers are advised to postpone irrigation work. Do not seek shelter or allow yourself or your animals to take refuge under trees during heavy rain and strong winds/storms.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
MAIZE (Silking)	In view of the possibility of rain, postpone irrigation. For prevention of stem borer pest in maize crop, spray Emamectin Benzoate 5% SG 200 gm per acre or Chlorantraniliprole 18.5% SC 60 ml per acre by dissolving it in 500-700 liters of water.
BLACK GRAM (Vegetative)	In view of the possibility of rain, postpone irrigation. Keep Zaid black gram/green gram crops free from weeds. Thrips/green leaf pest is likely to be seen in black gram crop, so for its prevention, spray Oxidemetan-methyl 25% EC or Dimethoate 30% EC at the rate of 1.0 liter/ha by making a solution in 600-700 liters of water.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
ONION (Physiological Maturity)	The best time to harvest onions is one week after 50% of the leaves fall. Thrips pest is likely to appear in the onion crop, so for its prevention, use Methyl-O-Dimethan 25% EC at the rate of 1.5 liters / ha spray the solution in 600-700 liters of water. Irrigation should be done by weeding and hoeing the vegetables planted in the summer season, tomato, brinjal and chilli. After harvesting the ready crops of pumpkin, gourd, taroi, bitter gourd, cucumber, cucumber, watermelon, muskmelon etc. sown in summer, send them to the market.
MANGO (Fruit formation)	Do irrigation work in orchards of mango, guava, lemon, grapes, papaya and litchi etc. Irrigate the orchards to prevent fall of mango fruit or apply 4.5 SL of 20 ml alpha naphthalene acetic acid make a solution per liter of water and spray it. To prevent cuckoo disorder or internal rot disease in mango pods, make borax 0.8% solution in 0.8 g / liter water and spray 2 at 10 days interval.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	Considering the current weather, farmers are advised tie the animals in a shade time place during the day. Do not tie the animals under the tree because this week the chances of falling trees twigs are high due to the speed of the winds. Vaccination in animals for prevention of hoof-mouth disease. Give sufficient quantity of grains to the animals with green and dry fodder. Animals must be given clean and fresh water 3-4 times a day. Do not tie pregnant animals to the sloping place. The best time to give anthelmintic drugs for the prevention of insects in the stomach of animals.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Farmers are advised to put jute curtains in the poultry house and soak the curtains with cold water during the afternoon, make arrangements for fans and ventilation. add dietary supplements; vitamins and energy feed ingredients to the hens as well as add calcium content to the hen. Give medicine for the prevention of worms in the stomach of hen (Divmirg). To protect the chickens from heat, arrange curtains, fans and ventilation in the hen house. To reduce the temperature of the hen house, cover the roof of the shed with straw.

Likely impacts of weather warnings (General)

According to the weather forecast received from the India Meteorological Department, a warning has been issued for the period between May 06 -08May , 2026, alerting to light to moderate rainfall, thunderstorms accompanied by lightning, strong surface winds, and hailstorms.

Impact based advisories (General)

In view of the weather conditions, farmers working in the fields are advised not to take shelter or allow themselves or their animals to take shelter under trees in case of heavy winds/storms or rain. Suspend irrigation activities for standing crops such as summer maize, black gram (Urad), and green gram (Moong). Furthermore, harvest mature crops such as wheat and pigeon pea (Arhar) and store them in a safe location.

Farmers are advised to download Unified "Mausam" and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: <https://play.google.com/store/apps/details/>

Meghdoot MobileApp link: <https://play.google.com/store/apps/details>

Damini MobileApp link : <https://play.google.com/store/apps/details>