



Gramin Krishi Mausam Sewa
Agromet Advisory Bulletin for MATHURA District
Chander Shekhar Azad University of Agriculture &
Technology
Kanpur, Uttar Pradesh



Agromet Advisory Bulletin

Date : 2026-08-04

Weather Forecast of District MATHURA (Uttar Pradesh) Issued On : 2026-08-04 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-08-05	2026-08-06	2026-08-07	2026-08-08	2026-08-09
Rainfall(mm)	11.0	11.0	18.0	7.0	10.0
Tmax(°C)	29.0	28.0	28.0	32.0	32.0
Tmin(°C)	24.0	24.0	23.0	23.0	24.0
RH-I(%)	88	87	89	88	87
RH-II(%)	69	72	68	55	53
Wind Speed(kmph)	10	8	9	7	7
Wind Direction(Degree)	248	238	244	360	221
Cloud Cover(Octa)	8	8	8	5	6
Warning	Thunderstorms & Lightning, Squall etc	No Warning	Heavy Rain	No Warning	No Warning

Forecast Summary:

According to the weather forecast received from the Indian Meteorological Department, due to moderate to dense cloud cover in the next five day there is a possibility of light to moderate rain with thunderstorms & lightning at local level between dated on 01 to 05 August, 2026. The maximum temperature is between 28.0- 32.0°C, which is likely to be 1-2°C below normal. and the minimum temperature is between 23.0-24.0°C, which is likely to be 1-2°C below normal. The maximum and minimum range of relative humidity is between 87-89 and 53-72%. Wind direction is south-west and wind speed is between 7.0-10.0 kmph with gusts expected to be 3-4 kmph high speed than normal.

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the India Meteorological Department (IMD), there is a warning of thunderstorms, strong winds light to moderate rain.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Carry out sowing and other agricultural activities when there is no rain and the sky is clear. Apply fertilizers, insecticides, and fungicides only when the weather clears up. Regularly monitor crops for pests and diseases after the rain. Keep livestock in a dry, safe area and provide them with clean drinking water. Do not tether animals under trees or near electricity poles, as falling branches or electric shocks from the poles could prove fatal to the animals.

General Advisory:

Farmers are advised to make proper arrangements for draining excess rainwater from standing Kharif crops, with the exception of the paddy crop. Operations such as sowing, and the application of pesticides, fungicides, fertilizers, and irrigation for crops like Urd and Moong should be postponed. Following rainfall, crops should be regularly monitored for pests and diseases. Do not tether livestock under trees or near electric poles. Bathe the animals in the morning and evening, keep them in a dry and safe location, and ensure the availability of clean drinking water. Use clean water for preparing pesticides, fungicides, and herbicides, and avoid spraying these chemicals while standing against the wind direction. If possible, wash hands thoroughly with soap after spraying, before eating, and after washing clothes.

SMS Advisory:

Except for the paddy crop, ensure proper drainage of excess rainwater from all other crops. Postpone activities such as sowing, application of pesticides/fungicides and fertilizers, and irrigation.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
RICE (Tillering)	Postpone the application of herbicides, pesticides, fungicides, and fertilizers, as well as irrigation activities, for standing paddy crops. For control of broad and narrow leaf weeds in paddy crop, apply Bispyribac Sodium 10%SC @ 0.20 liters per hectare after 15-20 days of transplanting spray it with 500-600 liters of water at appropriate humidity. If Khaira disease is seen in paddy, then spray 20-25 kg zinc sulphate and 2.5 kg lime solution in 800 liters of water for its control. Topdressing of urea @ 50 to 60 kg per hectare is done 25 to 30 days after transplanting of paddy. Keep in mind that weeds should be removed before top dressing and while topdressing, there should not be more than 2 to 3 cm of water in the field. If infestation of root worm pest is seen in paddy crop, spray Cartap hydrochloride 4% granular chemical at the rate of 20 to 25 kg per hectare in three to four cm stagnant water.
MAIZE (Vegetative)	Ensure proper drainage of excess rainwater from standing maize crops, and postpone the application of pesticides, fungicides, and fertilizers, as well as irrigation. In maize crop, second weeding-hoeing after 35-40 days. In maize crop, first top dressing should be done 30 to 35 days after sowing and second top dressing after 40 to 45 days after sowing, top dressing of urea at the rate of 60 to 70 kg per hectare should be clear sky. In maize crop, there is a possibility of outbreak of stem borer / shoot fly pest, so for its prevention, Emamectin benzoate 5% SG 200 g / ha or dimethoate 30% EC 1.0 liter / ha at 600-700 liters of water solution Sprinkle on clear weather.
SESAME (GINGELLY/TIL) (Vegetative)	Ensure proper drainage of excess rainwater from standing sesame crops and suspend weeding, hoeing, the application of pesticides, fungicides, and fertilizers, as well as irrigation. The first weeding of sesame crop should be done 15-20 days after sowing when the sky is clear.
GROUNDNUT (Vegetative)	Ensure proper drainage of excess rainwater from standing groundnut crops, and postpone weeding, hoeing, the application of pesticides and fertilizers, and irrigation. After weeding the groundnut crops 20 to 25 days after sowing, add 100 kg gypsum per hectare and then do light hoeing and cover the plants with soil.
BLACK GRAM (Sowing)	Ensure proper drainage of excess rainwater from standing black gram crops, and postpone sowing operations. Recommendation of the black gram species sown in Kharif - Narendra Urd-1, Azad Urd-2, Azad Urd-3, Shekhar-1, Shekhar-2, Shekhar-3, Pant U-30 etc. for sowing of any species Sowing should be started by arranging manure and seeds at the rate of 12 -15 kg seeds / hectare.

Crop (Stage)	Crop Specific Advisory
GREEN GRAM (Sowing)	Ensure proper drainage of excess rainwater from standing green gram crops, and postpone sowing operations. The recommended varieties of green gram sown in Kharif - Narendra Moong-1, Pant Moong-1, Pant Moong-3, Pant Moong-4, Samrat, Malaviya Janchetna, Malaviya Janapriya, Malaviya Jagriti Shweta, Swati, Virat and IPM -2-14 etc For the sowing of any one variety. Make the sowing work at the rate of 12 -15 kg seed / hectare .
SUGARCANE (Grand growth)	Properly manage the drainage of excess rainwater from the field. For controlling white grit, use light traps or spray insecticide on the plants. To prevent the autumn sugarcane from falling, tying up must be done. For weed control in standing sugarcane crop, do weeding and hoeing. For controlling top borer insect, mix 150-200 ml of insecticide Chlorantraniliprole 18.5 SC in 400-500 liters of water and spray it at the rate of per acre.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
BRINJAL (Transplanting)	Ensure proper drainage for excess rainwater and postpone activities related to pesticides, fertilizers, transplanting, sowing, and irrigation. Raise nursery of crops such as brinjal, chilli, tomato and early cauliflower planted in kharif, along with planting of brinjal / chilli seedlings. If there is a possibility of water stagnation in the field, then plant the early cauliflower, brinjal, tomato, chilli etc. on the ridges. For prevention of fruit borer/leaf borer pest in vegetable crops, prepare a solution of 1.5-2.0 ml/liter of water with neem oil and spray it 3-4 times at an interval of 8-10 days. Weeding-hoeing should be done in standing vegetable crops when sky is clear.
MANGO (Planting)	Perform the planting of new plants in the pits already filled with mangoes. For prevention of mango leaf moth, apply 2% carbaryl 1.5% dust on the clear sky. After weeding the banana/papaya garden, put 25-30 cm high soil around the stem and do the work of standing in banana plants.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	Farmers are advised to give sufficient quantity of grains with green and dry fodder to the animals. Take special care of calving animals and protect lactating animals from milk fever, ketosis etc. after calving. Protect animals and newborns from internal and external parasites. Use disinfectant medicine in the cattle shed. The best time to give antithetical drugs for the prevention of insects in the stomach of animals. Get the animals vaccinated against Foot-and-Mouth Disease. The animals must give clean and fresh water 3-4 times a day. Give the animals a bath two to three times a day. Tie the animals in the shade at the time of day and do not tie the animals under the tree, because this week due to the high wind speed, there is a high probability of falling of trees / tree twigs. The department is operating a Mobile Veterinary Unit scheme to provide veterinary services at the doorsteps of farmers and livestock owners. Farmers and livestock owners can contact the toll-free helpline number 1962 to avail the benefits of this scheme.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Protect the chickens from dampness and waterlogging during the current weather conditions. Ensure proper ventilation in the poultry shed and provide clean drinking water along with a balanced diet. Maintain cleanliness around the shed to control mosquitoes and flies, and ensure timely vaccination for all chickens. Immediately isolate any chickens that appear sick or lethargic, and seek veterinary advice promptly if necessary.

Likely impacts of weather warnings (General)

Waterlogging in low-lying areas disrupting daily activities and local commuting. Increased risk of road accidents due to reduced visibility and slippery roads. Potential for loss of life, injuries, and property damage caused by thunderstorms, lightning strikes, and strong winds. Risk of casualties and damage to property from falling trees, branches, hoardings, and electric poles; temporary disruption to power, communication, and other essential services. Impact on residential areas, markets, schools, and public facilities due to localized flooding or waterlogging; increased risk of waterborne and mosquito-borne diseases caused by high humidity and waterlogging. Damage to crops, horticultural produce, livestock shelters, and agricultural infrastructure. Adverse effects on commercial activities, construction work, and the operations of outdoor workers.

Impact based advisories (General)

Regularly monitor the weather forecasts and warnings issued. During thunderstorms and lightning, avoid open fields, tall trees, electric poles, and water bodies. Travel only when necessary and avoid waterlogged roads, bridges, and culverts. Do not stand under weak structures, hoardings, or trees during strong winds. Inspect and secure rooftops, windows, and drainage systems around your home. Use electrical appliances with caution and keep flashlights and other essential emergency supplies ready in case of power outages. Take special care of children, elderly persons, pregnant women, and individuals with health conditions. Use clean and safe drinking water to prevent water-borne diseases. Keep livestock in safe shelters and ensure adequate fodder and clean drinking water for them.

Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightning.

Mausam MobileApp link: <https://play.google.com/store/apps/details/>

Meghdoot MobileApp link: <https://play.google.com/store/apps/details>

Damini MobileApp link : <https://play.google.com/store/apps/details>