Gramin Krishi Mausam Sewa

Agromet Advisory Bulletin for MATHURA District

Chander Shekhar Azad University of Agriculture & Technology

Kanpur, Uttar Pradesh

Agromet Advisory Bulletin

Date : 2026-10-19

Weather Forecast of District MATHURA (Uttar Pradesh) Issued On : 2026-10-19 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-10-20	2026-10-21	2026-10-22	2026-10-23	2026-10-24
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	33.0	34.0	34.0	35.0	35.0
Tmin(°C)	21.0	21.0	21.0	21.0	21.0
RH-I(%)	53	53	57	54	53
RH-II(%)	23	22	22	20	18
Wind Speed(kmph)	3	5	2	2	2
Wind Direction(Degree)	34	290	233	243	360
Cloud Cover(Octa)	0	0	0	0	0
Warning	NA	NA	NA	NA	NA

Forecast Summary:

According to the weather forecast received from the Indian Meteorological Department, due to clear sky weather in the next five days, there is a no possibility of rain . The maximum temperature is between 33.0- 35.0°C, which is likely to be 1-2°C higher normal and the minimum temperature is between 21.0-21.0°C, which is likely to be 3-4°C higher normal. The maximum and minimum range of relative humidity is between 53-57 and 18-23%. Wind direction is north-east, south-west and wind speed is between 2.0-5.0 km/h with winds expected to be near normal.

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the India Meteorological Department (IMD), there is a no warning in next five days

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Farmers are advised to carry out the harvesting/threshing of mature crops and the sowing of new crops. Do not tether livestock in open areas; keep them in a dry, safe place and provide them with clean drinking water. Avoid tying animals under trees or near electricity poles, as falling branches or electric shocks from the poles could prove fatal to the animals.

General Advisory:

Proceed with the harvesting, threshing, and sowing operations for mature Kharif crops such as paddy, maize, and sesame. Carry out the sowing of Rabi crops—such as gram, peas, lentils, mustard, and linseed only after treating the seeds. Harvest timely-sown paddy crops when 85% of the crop has turned golden in color.

SMS Advisory:

Farmers are advised to carry out the harvesting and threshing of mature Kharif crops and the sowing of Rabi crops.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
RICE (Physiological Maturity)	Harvest the crops when 85% golden color appears in the paddy ears and after harvesting, dry the lock in the sun and do the threshing on clear weather. The current weather is favorable for pests, so after flowering, on an average, 2 - 3 odorous pests / hill are visible in the paddy crop, then for its control, Imidacoloroprid 17. 8% SL 350 ml / ha or Burfozin 25% SP 750 ml Make a solution in 500 -600 liters of water at the rate of / hectare and spray it on clear weather.
GROUNDNUT (Maturity and Harvesting)	The groundnut crop should be excavated only when the veins on the top of the peel emerge and the inner part becomes brown in color and the groundnut becomes pink. After harvesting the groundnut, dry the beans well in the sun and store them.
RAPESEED (Vegetative)	Along with weeding and hoeing within 20-22 days of sowing of rapeseed crop, make the distance from plant to plant to 10 - 15 cm. There is a possibility of outbreak of sawfly and hairy lard insect in rapeseed crop, so for its prevention, Imamectin Benzoate 5% SG at the rate of 200 g/ha or quinalphos 25 EC at the rate of 1.25 liters/ha in 500-600 liters of water. Make a solution and spray it.
MUSTARD (Sowing)	The weather is favorable for sowing work at the rate of 3-4 kg seed/ha for sowing any one of the recommended varieties of rye/mustard- Pusa Mustard-34, Pusa Mustard-32, Varuna (V-WRR-2), C5-64 (CS2005-143), Pusa OOM-36 (PDZ-15), BMP-II (DIMR), Azad Mahak, Surekha (KMR-16-02), Rohini, Narendra Rai- 8501, Maya, Vaibhav etc.
CHICK PEA (Sowing)	Recommended varieties of gram crop - Block, Pusa-256, Pusa-362, KWR-108, K-850, KDG-1168 etc. Sow any one variety and seed 75-80 kg of small grains and for large grain varieties, do sowing at the rate of 90-100 kg seed/ha.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
POTATO (Sowing)	Main varieties of potato - Kufri Bahar, Kufri Anand, Kufri Badshah, Kufri Sinduri, Kufri Sutlej, Kufri Evergreen Lalima, Kufri Arun, Kufri and Kufri Pukhraj, Kufri Garim, Kufri Lima, Kufri Lalit, Kufri Sangam, Kufri Neelkanth, Kufri Ganga, Kufri Thar-3, Kufri Mohan, Kufri Neelkanth, Kufri Surya, Kufri Chipsona-1, Kufri Chipsona-3, Kufri Chipsona-4 And the weather is favorable for sowing of Kufri, Friesona etc. To prevent chicken pox in potato crop, spray 100 grams of streptocycline or boric acid in 10 liters of water and spray the tubers on the seeds and do not treat the germinated seeds.
VEGETABLE PEA (Sowing)	Sow vegetables like peas, fennel, celery, beetroot, garlic, coriander, spinach, soya, fenugreek, radish, carrot, turnip and leafy vegetables. Raise nursery of tomato, onion, chilli, cauliflower and cabbage. If there is a possibility of water stagnation in the field, then plant cauliflower, brinjal, tomato, chilli etc. on the ridges. If there is an infestation of fruit borer/leaf borer insect in vegetable crops, then for its control, make a solution of neem oil in 1.5 ml/litre water and spray it 3-4 times at an interval of 8-10 days when the sky is clear.
PAPAYA (Transplanting)	Plant new plants in place of the destroyed plants in the new bag. Make a stand of 25-30 meters high soil around the stem by folding banana / papaya bags. For prevention of mango leaf moth, apply 2% carburil 1.5% dust .

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	Farmers are advised to Pregnant buffaloes/cows should not be tied to the slope, feed nutritious fodder and feed to pregnant buffaloes/cows and feed

Live Stock	Live Stock Specific Advisory
	milk to the newborn child for three days. Keep the animals in a clean place and smoke the animals to prevent fly and mosquito. Get animals vaccinated for the prevention of cracking-mouth disease. Animals suffering from this disease should wash their wounds with potassium permanganate. Give sufficient quantity of grains with green and dry fodder to the animals. Do not feed the stagnant water of the ponds or other places, due to which the animals are prone to liver flu and stomach worms. The animals must give clean and fresh water 3-4 times a day.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Farmers are advised to Clean every day at the place where the chickens live. Arrange for clean water and balanced diet to the hens. This is not the right time to raise new chicks for eggs but it is a suitable time for broiler chickens. Add supplementary food, vitamins and energy food material to the chickens' diet and also give calcium material to the chickens. Chlorination of water must be done. Give deworming medicine to kill the worms in the stomach of the chickens. Keep the chicken house dry and keep turning the bedding.

Likely impacts of weather warnings (General)

Prolonged exposure to sunlight and high humidity significantly increase the risk of heat stress and infectious diseases in livestock. High humidity and heat reduce the appetite of buffaloes and cows, potentially leading to a sudden drop in milk production. Stagnant water following recent monsoon rains, combined with rising temperatures, creates an environment where bacteria thrive in fodder and water sources. This can lead to the spread of ailments such as intestinal worms, skin infections, and Foot-and-Mouth Disease among the animals. The combination of a sudden rise in humid heat and a gradual drop in minimum temperatures creates conditions characterized by intense heat during the day and a slight chill at night or in the early morning. This fluctuating weather affects human health; venturing out in strong sunlight can lead to dehydration. Furthermore, this season sees a rapid surge in the prevalence of mosquitoes and viruses responsible for diseases like Chikungunya, Dengue, Malaria, and viral flu.

Impact based advisories (General)

Do not tether animals in direct sunlight during the day (between 11 AM and 4 PM); keep them in a well-ventilated, shaded shed instead. Ensure drinking water troughs are kept clean and provide fresh, clean water at least 3–4 times a day. If vaccinations for Hemorrhagic Septicemia (HS) or Black Quarter (BQ) have not been administered, get the animals vaccinated. At night, use neem leaf smoke in the animal shed to protect against mosquitoes and flies.Carry a water bottle when stepping out. Consume ORS, lemonade, or buttermilk to prevent electrolyte depletion in the body. Do not allow rainwater from recent showers to stagnate around the house or on rooftops. Use mosquito nets or repellents while sleeping. Avoid drinking cold water from the refrigerator or sitting directly in air-conditioned (AC) environments immediately after coming in from the heat; these habits are primary causes of colds, coughs, and lung infections. This dry week is ideal for preparing Rabi crops and for the harvesting

Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: https://play.google.com/store/apps/details/

Meghdoot MobileApp link: https://play.google.com/store/apps/details

Damini MobileApp link : https://play.google.com/store/apps/details