Gramin Krishi Mausam Sewa

Agromet Advisory Bulletin for AGRA District

Chander Shekhar Azad University of Agriculture & Technology

Kanpur, Uttar Pradesh

Agromet Advisory Bulletin

Date : 2026-10-27

Weather Forecast of District AGRA (Uttar Pradesh) Issued On : 2026-10-27 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-10-28	2026-10-29	2026-10-30	2026-10-31	2026-11-01
Rainfall(mm)	0.0	0.0	0.0	7.0	0.0
Tmax(°C)	34.0	34.0	32.0	28.0	29.0
Tmin(°C)	19.0	20.0	22.0	21.0	20.0
RH-I(%)	52	59	82	86	88
RH-II(%)	18	24	46	64	58
Wind Speed(kmph)	6	3	7	9	7
Wind Direction(Degree)	4	354	79	88	37
Cloud Cover(Octa)	1	5	4	7	3
Warning	No Warning	No Warning	No Warning	No Warning	No Warning

Forecast Summary:

According to the weather forecast received from the Indian Meteorological Department, due to moderate to dense cloud cover in next five days , there is a possibility of light to moderate rain dated on 31 Oct, 2026. The maximum temperature is between 28.0- 34.0°C, which is likely to be 2-3°C higher normal and the minimum temperature is between 19.0-22.0°C, which is likely to be 3-4°C higher normal. The maximum and minimum range of relative humidity is between 52-88 and 18-64%. Wind direction is south-east, north-west, north-east and wind speed is between 3.0-9.0 km/h with winds expected to be 2-3 Km/h normal.

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the India Meteorological Department, there is a warning of light to moderate rain with Thunderstorms & Lightning, Squall etc; Strong Surface Winds.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Heavy rainfall and strong winds can cause standing or harvested paddy crops to fall over. This increases moisture content in the grains, potentially causing them to turn black and damaging the harvest. Rain immediately after sowing can form a crust on the soil surface, which may severely hinder the germination of mustard, potato, and early-sown wheat seeds. High humidity and persistent cloud cover can lead to fungal growth on harvested crops, as well as an increased risk of root rot and blast disease in vegetables and pulse crops. If the paddy crop is fully mature, postpone harvesting until October 30th. Crops that have already been harvested should be moved from the fields to a safe, elevated location or covered thoroughly with tarpaulins. For fields prepared for mustard, early-sown wheat, and potatoes, wait until after the rain—once the soil is dry enough to be worked—before proceeding with sowing. In fields used for vegetables, peas, and pulses, ensure that a system of ridges and fu

General Advisory:

In view of the potential weather changes or light rain expected in the coming days, please move harvested crops from the fields to safe, elevated locations, or cover them thoroughly with tarpaulins.

Farmers sowing mustard, potatoes, or early-sown wheat should ensure appropriate soil moisture and proceed with sowing only once the weather has completely cleared. Avoid heavy irrigation, as well as the application of pesticides or fertilizers, for the next 2–3 days to prevent the wastage of water and unnecessary expenditure. To protect milch animals from dry, cold north-westerly winds, cover their shelters (enclosures) at night using tarpaulins or jute sacks.

SMS Advisory:

In view of the potential weather changes or light rain expected in the coming days, please move harvested crops from the fields to safe, elevated locations, or cover them thoroughly with tarpaulins.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
RICE (Physiological Maturity)	In view of the possibility of rain, move harvested crops from the fields to a safe, elevated location or cover them thoroughly with a tarpaulin. Harvest the paddy crop when 85% of the panicles have turned golden yellow. After harvesting, dry the produce in the sun and carry out threshing during clear weather. The current weather conditions are favorable for pests; therefore, if an average of 2–3 rice ear-cutting bugs (gandhi bugs) per hill are observed after the flowering stage, take control measures by spraying a solution of either Imidacloprid 17.8% SL (350 ml/hectare) or Buprofezin 25% SP (750 ml/hectare) mixed in 500–600 liters of water. Perform this application during clear weather.
RAPESEED (Vegetative)	Ensure proper drainage. There is a possibility of an outbreak of hairy caterpillars in the *toriya* (rapeseed) crop; to prevent this, spray a solution of Emamectin Benzoate 5% SG (200 grams/hectare) mixed in 500–600 liters of water. Within 20–22 days of sowing, perform weeding and hoeing, and ensure a spacing of 10–15 cm between plants. Provide the first irrigation to the *toriya* crop within 25–30 days of sowing, and once the soil reaches the appropriate moisture level, apply a top dressing of urea at a rate of 108 kg/hectare.
MUSTARD (Sowing)	In the absence of rain, the mustard crop is susceptible to the mustard sawfly, which can cause significant damage 7 to 10 days after germination. To control this pest, apply 25 kg/hectare of either 2% methyl parathion, 5% malathion, or 5% carbaryl dust during the morning or evening hours. For sowing, select one of the recommended mustard/rapeseed varieties—such as Pusa Sarson-34, Pusa Sarson-32, C-5-64 (CS2005-143), Pusa OOM-36 (PDZ-15), BMP-II (DMR), Azad Mahak, Surekha (KMR-16-02), Varuna, Rohini, Narendra Rai-8501, Maya, or Vaibhav—and sow at a seed rate of 3–4 kg per hectare.
FIELD PEA (Sowing)	In the absence of rain, sow recommended pea varieties—such as Rachna, Malviya Matar-15, or Sapna—at a seed rate of 100–125 kg per hectare.
CHICK PEA (Sowing)	In the absence of rain, sow recommended chickpea varieties—such as Avrodhi, Pusa-256, Pusa-362, KWR-108, K-850, or KDG-1168—using a seed rate of 75–80 kg per hectare for small-seeded varieties, and 90–100 kg per hectare for large-seeded varieties.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
POTATO (Sowing)	Ensure proper drainage. In the absence of rain, proceed with the sowing of major potato varieties, such as Kufri Bahar, Kufri Anand, Kufri Badshah, Kufri Sindhuri, Kufri Satluj, Kufri Lalima, Kufri Arun, Kufri Sadabahar, Kufri Pukhraj, Kufri Garima, Kufri Lima, Kufri Lalit, Kufri Sangam, Kufri Neelkanth, Kufri Ganga, Kufri Thar-3, Kufri Mohan, Kufri Surya, Kufri Chipsona-1, Kufri Chipsona-3, Kufri Chipsona-4, and Kufri Frysona. To protect the potato crop from common scab (bacterial spot/lesions), treat the tubers by spraying them with a solution of 100 grams of Streptocycline or boric acid in 10 liters of water; do not treat tubers that have already sprouted.
VEGETABLE PEA (Sowing)	Ensure proper drainage. In the absence of rain, sow vegetable peas, fennel, ajwain (carom seeds), beetroot, garlic, coriander, spinach, dill, fenugreek, radish, carrot, turnip, and other leafy vegetables. Prepare nurseries for tomato, onion, chili, cauliflower, and cabbage. If there is a risk of water stagnation in

Horticulture (Stage)	Horticulture Specific Advisory
	the field, transplant crops like cauliflower, eggplant (brinjal), tomato, and chili on raised beds. If an infestation of fruit borers or leaf miners is observed in vegetable crops, control it by spraying a solution of 1.5 ml of neem oil per liter of water; apply 3–4 times at intervals of 8–10 days, ensuring the application is done when the sky is clear.
PAPAYA (Fruit formation)	n new orchards of papaya, mango, guava, amla (Indian gooseberry), lemon, etc., replace any lost or damaged plants by planting new ones. For banana and papaya orchards, hoe the soil and mound it up to a height of 25–30 cm around the base of the stems. Provide support for banana and papaya trees using wooden or bamboo stakes. To control leaf-cutting insects in mango trees, apply a 2% carbaryl dust (or 1.5% carbaryl dust formulation).

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	During heavy rain and storms, seek shelter in a sturdy, permanent building as soon as you hear thunder or see lightning; avoid standing under trees, electricity poles, transformers, or mobile towers. Never attempt to cross waterlogged roads, drains, or underpasses, as the current and depth of the water can be life-threatening. To prevent waterborne diseases (such as diarrhea or cholera), consume only boiled or filtered water. Keep livestock in a safe, elevated, dry, and secure area to prevent hoof rot or other infections. Never tether animals near electricity poles, wires, or metal structures/sheds; during lightning storms, do not leave them in open fields or under trees. Ensure you have a sufficient stock of dry fodder stored in advance, as consuming wet fodder can cause digestive issues, such as abdominal pain and bloating (gas), in livestock.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Farmers are advised to Clean every day at the place where the chickens live. Arrange for clean water and balanced diet to the hens. This is not the right time to raise new chicks for eggs but it is a suitable time for broiler chickens. Add supplementary food, vitamins and energy food material to the chickens' diet and also give calcium material to the chickens. Chlorination of water must be done. Give deworming medicine to kill the worms in the stomach of the chickens. Keep the chicken house dry and keep turning the bedding.

Likely impacts of weather warnings (General)

Due to a sudden drop in temperature, daytime temperatures may fall by 5–6°C. This change increases the risk of seasonal colds, coughs, and allergies; therefore, keep light, warm clothing or full-coverage garments ready. During rain, low-lying areas and underpasses may become waterlogged; check local route updates before traveling.In light of the rain, keep animal enclosures dry. Damp flooring can lead to hoof infections in livestock. Ensure an adequate stock of dry fodder is stored in a safe, dry place before the rain begins; consuming wet or moldy fodder can cause severe abdominal pain and bloating (gas) in animals. With a significant 12–15°C difference between day and night temperatures, the risk of catching a cold is highest during the early morning and late night hours. Protect the elderly and children from the cold morning air. Consume lukewarm water and eat fruits rich in Vitamin C to maintain a strong immune system. If the paddy crop is fully mature, it is advisable to postpone harvesting for 1–2 days given the expected rain. Crops that have already been harvested and are drying in the fields should be moved to a safe location immediately or covered securely with tarpaulins to prevent the grains from darkening or absorbing moisture. Store grain in elevated areas and keep tarpaulins handy to protect it from getting wet in the event of unexpected rain. For fields prepared for the sowing of mustard, early-sown wheat, or potatoes, it is best to wait until after the rain to sow. Sowing before the rain can cause a crust to form on the soil surface, which negatively affects seed germination. Ensure proper drainage in the fields to prevent water accumulation.

Impact based advisories (General)

During heavy rain and storms, seek shelter in a sturdy, permanent building as soon as you hear thunder or see lightning; avoid standing under trees, electricity poles, transformers, or mobile towers. Never attempt to cross waterlogged roads, drains, or underpasses, as the current and depth of the water can be life-threatening. To prevent waterborne diseases (such as diarrhea or cholera), consume only boiled or filtered water.

Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: https://play.google.com/store/apps/details/

Meghdoot MobileApp link: https://play.google.com/store/apps/details

Damini MobileApp link : https://play.google.com/store/apps/details