



Gramin Krishi Mausam Sewa
Agromet Advisory Bulletin for MATHURA District
Chander Shekhar Azad University of Agriculture &
Technology
Kanpur, Uttar Pradesh



Agromet Advisory Bulletin

Date : 2026-05-19

Weather Forecast of District MATHURA (Uttar Pradesh) Issued On : 2026-05-19 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-05-20	2026-05-21	2026-05-22	2026-05-23	2026-05-24
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	46.0	46.0	46.0	47.0	47.0
Tmin(°C)	29.0	29.0	29.0	29.0	30.0
RH-I(%)	46	46	46	46	46
RH-II(%)	26	26	26	26	25
Wind Speed(kmph)	8	13	17	12	17
Wind Direction(Degree)	300	289	268	287	271
Cloud Cover(Octa)	0	0	0	0	1
Warning	Heat Wave	Heat Wave	Heat Wave	Heat Wave	Heat Wave

Forecast Summary:

According to the weather forecast received from the India Meteorological Department, due to clear sky weather in the next five days there is any no possibility of rain but there is a heat wave on dated on 20-24 May, 2026. The maximum temperature is between 46.0- 47.0°C, which is likely to be 6-7°C higher normal and the minimum temperature is between 29.0-30.0°C, which is likely to be 5-6°C higher normal. The maximum and minimum range of relative humidity is between 46-46 and 25-26%. Wind direction is south-west, north-west and wind speed is between 8.0-17.0 kmph with gusts expected to be 5-6 kmph high speed than normal.

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the Indian Meteorological Department, there is a warning of intense heat due to clear skies in the next five days.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

In view of the prevailing weather conditions, farmers are advised to provide light irrigation to standing crops in their fields to maintain adequate moisture levels and protect them against heatwaves; irrigation activities should be carried out during the evening hours. The harvesting of mature crops should be undertaken during the early morning or late evening. Farmers should refrain from performing any field work between 11:00 AM and 4:00 PM, and livestock should not be allowed to graze outdoors during this period. Livestock should be bathed both in the morning and in the evening. Given the intense heat, farmers are advised to drink water at frequent intervals while working in the fields. Furthermore, one should avoid stepping out of the house unnecessarily during the afternoon hours; if venturing out for

essential tasks is unavoidable, one should do so only after covering the head with a cloth (such as a *gamcha*).

General Advisory:

Farmers are advised to maintain adequate soil moisture in standing vegetable crops and field crops by providing light irrigation as needed. While working in the fields, they should cover their heads with a cloth (Gamcha) and ensure they carry a water bottle with them. In view of the severe heat, livestock should be given water 3–4 times a day and bathed in the morning and evening. When using pesticides, fungicides, or herbicides, use separate sprayers for each type of chemical; alternatively, thoroughly wash the equipment with clean water before switching between different chemicals. Furthermore, strictly avoid spraying or dusting these chemicals while standing against the wind direction. Whenever possible, spraying should be carried out during the evening hours. After spraying—and before eating or drinking—one must thoroughly wash their hands with soap or hand wash, wash their clothes, and take a bath.

SMS Advisory:

Farmers are advised to maintain proper moisture in standing crops by doing light irrigation as per requirement. To avoid heat stroke, do not go out of the house unnecessarily during the afternoon.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
MAIZE (Millking)	Maize crop is going on at the stage of cumin extract /cob formation/grain filling which is sensitive to moisture deficiency. Therefore, maintain proper moisture by giving light irrigation as per requirements. There is a possibility of outbreak of stem borer / shoot fly pest in maize crop, so for its prevention, solution of Emamectin Benzoate 5% SG at the rate of 200 g/ha or Dimethoate 30% EC 1.0 l/ha in 600-700 liters of water make a spray.
BLACK GRAM (Pod initiation)	There is a possibility of outbreak of pod borer pest in black gram crop, so for its prevention, apply Emamectin Benzoate 5% SG 200 g/ha or Dimethoate 30% EC. Make a solution in 600-700 liters of water at the rate of 1.0 liter/ha and spray it when the sky is clear.
GREEN GRAM (Flowering)	There is a possibility of outbreak of pod borer pest in green gram crop, so for its prevention, apply Emamectin Benzoate 5% SG 200 g/ha or Dimethoate 30% EC. Make a solution in 600-700 liters of water at the rate of 1.0 liter/ha and spray it when the sky is clear.
MAIZE (Sowing)	Prepare the fields for the sowing of Kharif maize, as well as the advanced recommended complex species- Kanchan, Gaurav, Sweta, Azad, the best and hybrid varieties- Prakash, Y-1402, Bio-9681, Pro-316, D-9144, Dikalb- For sowing any one species from 7074, PHB-8144, Deccan 115, MMH-133 etc., arranging manure and seeds. For sowing of complex variety of maize, sow at the rate of 20 -25 kg seed / hectare and Shankar species at the rate of 18 -20 kg seed / hectare.
RICE (Nursery Bed Preparation)	For preparation of paddy fields, do deep plowing and fencing. Light irrigation of the field 15 days before planting paddy nursery so that the weeds coming out in the field are destroyed while preparing the field.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
BRINJAL (Physiological Maturity)	Harvest the finished crops like Pumpkin, Gourd, Taroi, Bitter gourd, Cucumber, Cucumber, Watermelon and Melon etc. and send them to the market. For the prevention of fruit borer / leaf borer pest in vegetable crops,

Horticulture (Stage)	Horticulture Specific Advisory
	make a solution of neem oil in 1.5-2.0 ml / liter of water and do 3-4 sprays at an interval of 8 -10 days. Do weeding and hoeing in the standing crops of vegetables and do irrigation work at an interval of 8-10 days in the evening.
MANGO (Physiological Maturity)	Dig pits for planting new gardens. Do irrigation work in orchards of mango, guava, lemon plum, grapes, papaya and litchi etc. Irrigate the orchards to prevent fall of mango fruit or apply 4.5 SL of 20 ml alpha naphthalene acetic acid make a solution per liter of water and spray it. To prevent cuckoo disorder or internal rot disease in mango pods, make borax 0.8% solution in 0.8 g / liter water and spray 2 at 10 days interval.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	Considering the current weather, farmers are advised to tie the animals in the open during the night. Tie the animals in a shade time place during the day. Do not tie the animals under the tree because this week the chances of falling trees twigs are high due to the speed of the winds. FMD for the prevention of foot-mouth disease to animal's vaccine and VQ for prevention of Lagadia fever. Give sufficient quantity of grains to the animals with green and dry fodder. Animals must be given clean and fresh water 3-4 times a day. Do not tie pregnant animals to the sloping place. The best time to give anthelmintic drugs for the prevention of insects in the stomach of animals.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Farmers are advised to add dietary supplements; vitamins and energy feed ingredients to the hens as well as add calcium content to the hen. Give medicine for the prevention of worms in the stomach of hen (Divmirg). To protect the chickens from heat, arrange curtains, fans and ventilation in the hen house. To save the chickens from the heat, sprinkle water on the curtains engaged in the chicken house so that the coolness remains.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation)	Others (Soil / Land Preparation) Specific Advisory
AGRI FIELD	Deep plow the fields empty of wheat and other crops with disc plow so that the eggs and larvae of harmful insects hidden inside the ground are destroyed in the sun.

Likely impacts of weather warnings (General)

According to the weather forecast received from the Indian Meteorological Department, there is a warning of intense heat due to clear skies in the next five days.
--

Impact based advisories (General)

In view of the prevailing weather conditions, farmers are advised to provide light irrigation to standing crops in their fields to maintain adequate moisture levels and protect them against heatwaves; irrigation activities should be carried out during the evening hours. The harvesting of mature crops should be undertaken during the early morning or late evening. Farmers should refrain from performing any field work between 11:00 AM and 4:00 PM, and livestock should not be allowed to graze outdoors during

this period. Livestock should be bathed both in the morning and in the evening. Given the intense heat, farmers are advised to drink water at frequent intervals while working in the fields. Furthermore, one should avoid stepping out of the house unnecessarily during the afternoon hours; if venturing out for essential tasks is unavoidable, one should do so only after covering the head with a cloth (such as a *gamcha*).

Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: <https://play.google.com/store/apps/details/>

Meghdoot MobileApp link: <https://play.google.com/store/apps/details>

Damini MobileApp link : <https://play.google.com/store/apps/details>