



**Gramin Krishi Mausam Sewa**  
**Agromet Advisory Bulletin for MAINPURI District**  
**Chander Shekhar Azad University of Agriculture & Technology**  
Kanpur, Uttar Pradesh



## **Agromet Advisory Bulletin**

Date : 2026-06-30

Weather Forecast of District MAINPURI (Uttar Pradesh) Issued On : 2026-06-30 (Valid Till 08:30 IST of the next 5 days)

| Parameter              | 2026-07-01 | 2026-07-02                                                  | 2026-07-03                                                  | 2026-07-04                            | 2026-07-05                            |
|------------------------|------------|-------------------------------------------------------------|-------------------------------------------------------------|---------------------------------------|---------------------------------------|
| Rainfall(mm)           | 14.0       | 39.0                                                        | 6.0                                                         | 7.0                                   | 4.0                                   |
| Tmax(°C)               | 33.0       | 33.0                                                        | 31.0                                                        | 31.0                                  | 32.0                                  |
| Tmin(°C)               | 25.0       | 25.0                                                        | 24.0                                                        | 24.0                                  | 24.0                                  |
| RH-I(%)                | 76         | 80                                                          | 87                                                          | 86                                    | 82                                    |
| RH-II(%)               | 52         | 52                                                          | 62                                                          | 62                                    | 56                                    |
| Wind Speed(kmph)       | 5          | 5                                                           | 6                                                           | 7                                     | 3                                     |
| Wind Direction(Degree) | 278        | 234                                                         | 360                                                         | 360                                   | 131                                   |
| Cloud Cover(Octa)      | 5          | 6                                                           | 8                                                           | 7                                     | 7                                     |
| Warning                | No Warning | Thunderstorms & Lightning, Squall etc; Strong Surface Winds | Thunderstorms & Lightning, Squall etc; Strong Surface Winds | Thunderstorms & Lightning, Squall etc | Thunderstorms & Lightning, Squall etc |

### Forecast Summary:

According to the weather forecast received from the India Meteorological Department, due to moderate to dense cloud cover in the next five days, there is a possibility of light to moderate with Thunderstorms & Lightning, Squall etc and Strong Surface Winds. The maximum temperature is between 31.0-33.0°C, which is likely to be 3-4°C lower its normal and the minimum temperature is between 24.0-25.0°C, which is likely to be 1-2°C lower normal. The maximum and minimum range of relative humidity is between 76-87 and 52-62%. Wind direction is south-east, south-west, north and wind speed is between 3.0-7.0 kmph with gusts expected to be 1-2 kmph high speed than normal.

### Weather Warnings (Valid Till 08:30 IST of the next day)

According to the India Meteorological Department, there is a warning of light to moderate rain with Thunderstorms & Lightning, Squall etc; Strong Surface Winds in next five days.

### Likely impacts of weather warnings on Agriculture and associated Agromet advisories

After harvesting and threshing the mature crops, structure the grain and if there is harvested crop, keep it in a safe place and cover it with a polythene sheet.

### General Advisory:

Farmers are advised to harvest and thresh mature maize and urad/moong crops, and to sow Kharif maize and vegetables, in the absence of rain. Do not tether livestock under trees. Bathe the animals in the morning and evening, keep them in a shaded area, and provide them with water 3–4 times a day. Do not spray insecticides, fungicides, or herbicides while standing against the wind direction. If possible, wash hands thoroughly with soap after spraying, before eating, and after washing clothes.

### SMS Advisory:

Farmers are advised to harvest and thresh mature maize and urad/moong crops, and to sow Kharif maize and vegetables, in the absence of rain.

### Crop Specific Advisory:

| Crop (Stage)                         | Crop Specific Advisory                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|--------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| MAIZE (Sowing)                       | Ensure proper drainage for timely sown maize crops to handle excessive rainfall. After preparing the fields, arrange for seeds and fertilizers to sow one of the recommended improved composite varieties (such as Kanchan, Gaurav, Shweta, Azad Uttam) or hybrid varieties (such as HMM-3904, PMH-3, NK-61, Prakash, Y-1402, Bio-9681, Pro-316, D-9144, Dekalb-7074, PHB-8144, Deccan 115, MMH-133, etc.). Sow at a rate of 20–25 kg/hectare for composite varieties and 18–20 kg/hectare for hybrid varieties, provided there is no rainfall at the time of sowing.                                                                                                                                                                                                                                                                                                                                                        |
| RICE (Transplanting)                 | Transplant the ready paddy seedlings. Strengthen the field embankments (bunds) to retain rainwater in the fields. Sow nurseries for early-maturing paddy varieties. Transplant 20–25 day-old seedlings in normal fields and 25–30 day-old seedlings in saline/alkaline (usar) lands. To control broad-leaved and narrow-leaved weeds in the paddy crop, apply Anilofos 30% EC or Pretilachlor at a rate of 1.25 liters/hectare—mixed in 500–600 liters of water—within 2–3 days of transplanting, maintaining a water depth of 2–3 inches. If symptoms of 'Khaira' disease (caused by zinc deficiency)—characterized by yellowing leaves followed by the appearance of brown spots—are observed in the nursery, control it by applying 5 kg of zinc sulphate mixed with a 2% urea solution, or 2.5 kg... In the absence of rain, spray a solution of slaked lime prepared in approximately 1000 liters of water per hectare. |
| SESAME (GINGELLY/TIL) (Sowing)       | Ensure proper drainage arrangements to handle excessive rainfall. Sow improved sesame varieties—such as Gujarat Til-6, RT-346, RT-351, Tarun, Pragati, Shekhar, Type-78, Type-13, Type-4, Type-12, MT-2013-3, and UAT Til-1—once rainfall occurs and soil moisture is adequate.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| MAIZE (Maturity and Harvesting)      | Farmers are advised to harvest mature maize cobs; after shelling the grains, dry them thoroughly in the sun as quickly as possible before storage. If the crop has already been harvested, store it in a safe place and cover it with a polythene sheet.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| BLACK GRAM (Maturity and Harvesting) | Harvest and thresh mature Urd (black gram) or Moong (green gram) crops when there is no rain; if the crop has already been harvested, store it in a safe place, cover it with a polythene sheet, and proceed with threshing promptly to preserve the grain.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |

### Horticulture Specific Advisory:

| Horticulture (Stage)      | Horticulture Specific Advisory                                                                                                                             |
|---------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------|
| CAULIFLOWER (Nursery Bed) | Ensure proper drainage for excess rainwater. Prepare nurseries for Kharif vegetables and transplant the seedlings onto raised beds. This is the right time |

| Horticulture (Stage)           | Horticulture Specific Advisory                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|--------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Preparation)                   | to sow nurseries for brinjal, chili, and early-maturing cauliflower, and to sow crops like okra, cowpea, pumpkin, bottle gourd, ridge gourd, bitter gourd, and cucumber. Harvest the mature Zaid season crops—such as pumpkin, bottle gourd, bitter gourd, cucumber, watermelon, and muskmelon—and send them to the market. To protect vegetable crops from fruit borers and leaf miners, prepare a solution by mixing 1.5–2.0 ml of neem oil per liter of water and spray it 3–4 times at intervals of 8–10 days. Carry out weeding and hoeing in standing vegetable crops and irrigate them in the evening at intervals of 8–10 days. To control green hopppers and whiteflies in cucurbitaceous crops, spray 1.0 ml of Imidacloprid 30.5%; to control stem and fruit borers, spray a solution of 1.0 ml of Flubendiamide 39.35% per liter of water. Install 'Cue-lure pheromone traps' at a rate of 8–10 traps per hectare to protect against fruit flies. |
| MANGO (Physiological Maturity) | To fill the pits dug in May for establishing fruit orchards, mix well-decomposed cow dung manure with the topsoil and fill the pit up to 15–20 cm above the ground level. To protect mango and guava crops from fruit flies, install 8–10 traps per hectare; use Methyl Eugenol and Cue-lure traps for this purpose and tie them to branches at a height of 6–8 feet. Dissolve 5% neem extract in one liter of water and spray at intervals of 10 to 15 days. Replace the lure at intervals of 20–25 days.                                                                                                                                                                                                                                                                                                                                                                                                                                                    |

#### Live Stock Specific Advisory:

| Live Stock | Live Stock Specific Advisory                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
|------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| BUFFALO    | Considering the current weather, farmers are advised to tie the animals in the open during the night. Tie the animals in a shade time place during the day. Do not tie the animals under the tree because this week the chances of falling trees twigs are high due to the speed of the winds. FMD for the prevention of foot-mouth disease to animal's vaccine and VQ for prevention of Lagadia fever. Give sufficient quantity of grains to the animals with green and dry fodder. Animals must be given clean and fresh water 3-4 times a day. Do not tie pregnant animals to the sloping place. The best time to give anthelmintic drugs for the prevention of insects in the stomach of animals. |

#### Poultry Specific Advisory:

| Poultry | Poultry Specific Advisory                                                                                                                                                                                                                                                                                                                                                                                                                        |
|---------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| HEN     | Farmers are advised to add dietary supplements; vitamins and energy feed ingredients to the hens as well as add calcium content to the hen. Give medicine for the prevention of worms in the stomach of hen (Divmirg). To protect the chickens from heat, arrange curtains, fans and ventilation in the hen house. To save the chickens from the heat, sprinkle water on the curtains engaged in the chicken house so that the coolness remains. |

#### Likely impacts of weather warnings (General)

|                                                                                                                                                                                    |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| According to the India Meteorological Department, there is a warning of light to moderate rain with Thunderstorms & Lightning, Squall etc; Strong Surface Winds in next five days. |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

#### Impact based advisories (General)

|                                                                                                                                                                   |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| After harvesting and threshing the mature crops, structure the grain and if there is harvested crop, keep it in a safe place and cover it with a polythene sheet. |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------|

**Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.**

**Mausam MobileApp link: <https://play.google.com/store/apps/details/>**

**Meghdoot MobileApp link: <https://play.google.com/store/apps/details>**

**Damini MobileApp link : <https://play.google.com/store/apps/details>**