



Gramin Krishi Mausam Sewa
Agromet Advisory Bulletin for LUCKNOW District
Chander Shekhar Azad University of Agriculture &
Technology
Kanpur, Uttar Pradesh



Agromet Advisory Bulletin

Date : 2026-06-25

Weather Forecast of District LUCKNOW (Uttar Pradesh) Issued On : 2026-06-25 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-06-26	2026-06-27	2026-06-28	2026-06-29	2026-06-30
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	38.0	42.0	42.0	42.0	39.0
Tmin(°C)	29.0	30.0	30.0	30.0	30.0
RH-I(%)	61	63	65	63	65
RH-II(%)	36	39	43	45	45
Wind Speed(kmph)	3	9	12	13	20
Wind Direction(Degree)	360	284	291	282	305
Cloud Cover(Octa)	1	2	3	2	4
Warning	Heat Wave	Heat Wave	Heat Wave	Heat Wave	No Warning

Forecast Summary:

According to the weather forecast received from the India Meteorological Department, light cloud cover in the next five days, there is no any possibility of rain. The maximum temperature is between 38.0-42.0°C, which is likely to be 2-3°C higher normal and the minimum temperature is between 29.0-30.0°C, which is likely to be 1-2°C higher normal. The maximum and minimum range of relative humidity is between 61-65 and 36-45%. Wind direction is north-west and wind speed is between 3.0-20.0 kmph with gusts expected to be 7-8 kmph high speed than normal.

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the India Meteorological Department, the sky is likely to remain partly cloudy over the next five days with no chance of rain, though conditions of humid heat may persist. Additionally, a heatwave warning has been issued.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Harvest and thresh the mature Zaid crops and store the grain; the weather is favorable for sowing Kharif maize, preparing paddy nurseries, and transplanting seedlings.

General Advisory:

Farmers are advised that the weather is favorable for harvesting and threshing mature maize and black gram (urd)/green gram (moong) crops, as well as for sowing Kharif maize and vegetables. Do not tether livestock under trees. Bathe the animals in the morning and evening, keep them in a shaded area, and

provide them with water 3–4 times a day. Do not spray insecticides, fungicides, or herbicides while standing against the wind direction. If possible, wash hands thoroughly with soap after spraying, before eating, and after washing clothes.

SMS Advisory:

Farmers are advised that the weather is favorable for the harvesting and threshing of mature maize and urd/moong crops, as well as for the sowing of Kharif maize and vegetables.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
MAIZE (Sowing)	After preparing the field for sowing Kharif maize, arrange for fertilizers and seeds for sowing any one of the improved recommended complex varieties - Kanchan, Gaurav, Shweta, Azad Uttam and hybrid varieties - HMH-3904, PMH-3, NK-61, Prakash, Y-1402, Bio-9681, Pro-316, D-9144, Dikab- 7074, PHB- 8144, Deccan 115, MMH-133 etc. For sowing complex variety of maize, sow at the rate of 20-25 kg seed/hectare and Shankar variety at the rate of 18-20 kg seed/hectare.
RICE (Transplanting)	Strengthen the field embankments to facilitate paddy transplanting and ensure rainwater is retained in the field. Prepare nurseries for early-maturing paddy varieties. Transplant seedlings when they are 20–25 days old for normal fields, and 25–30 days old for saline or alkaline (usar) lands. To control broad-leaved and narrow-leaved weeds in the paddy crop, apply Anilofos 30% EC or Pretilachlor (at a rate of 1.25 liters/hectare) mixed in 500–600 liters of water within 2–3 days of transplanting, while maintaining a water depth of 2–3 inches. If symptoms of 'Khaira' disease—caused by zinc deficiency and characterized by yellowing leaves followed by the appearance of brown spots—are observed in the nursery, control it by applying 5 kg of zinc minerals mixed with a 2% urea solution, or by dissolving 2.5 kg of slaked lime in approximately 1000 liters of water per hectare (to be applied in the absence of natural rainfall).
SESAME (GINGELLY/TIL) (Sowing)	Sow improved varieties of sesame such as Gujarat Til-6, R.T.-346, R.T.-351, Tarun, Pragati, Shekhar Type-78, Type-13, Type-4 and Type-12, M.T.-2013-3 and U.A.T. Til-1.
MAIZE (Maturity and Harvesting)	Farmers are advised to harvest the mature maize corn and after separating the grain from the broken corn, dry it well in the sun and store it.
BLACK GRAM (Maturity and Harvesting)	The weather is favorable for the harvesting and threshing of mature.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
CAULIFLOWER (Nursery Bed Preparation)	Raise nursery/ sow seedlings of vegetables to be sown in Kharif. This is the appropriate time for nursery of brinjal, chili, early cauliflower and sowing of ladyfinger, cowpea, pumpkin, bottle gourd, ridge gourd, bitter gourd, cucumber etc. Harvest the ready crops of Zaid pumpkin, bottle gourd, bitter gourd, cucumber, watermelon, muskmelon etc. and send them to market. For prevention of fruit borer/leaf borer pest in vegetable crops, prepare solution of neem oil 1.5-2.0 ml/liter in water and spray 3-4 times at an interval of 8-10 days. Do weeding in standing vegetable crops and do irrigation work in the evening at an interval of 8-10 days. For controlling green hopper and white fly in cucurbitaceous crops, spray 1.0 ml of Imidacloprid 30.5% chemical and for controlling stem and fruit borer pests, spray 1.0 ml of Flubendiamide 39.35% chemical by dissolving it in one litre

Horticulture (Stage)	Horticulture Specific Advisory
	of water. To protect against fruit fly, apply Q Lure Pheromone Trap at the rate of 8-10 traps per hectare.
MANGO (Physiological Maturity)	For filling the pits dug in the month of May for planting fruit orchards, mix rotten manure of cow dung in the upper half of the soil and fill it up to 15 to 20 cm above the ground level. To protect mango and guava from fruit flies, hang 8-10 traps per hectare using methyl eugenol and Q lure traps tied to branches at a height of 6 to 8 feet. Spray 5% neem extract dissolved in one litre of water at an interval of 10 to 15 days. Change the lure at an interval of 20-25 days.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	Considering the current weather, farmers are advised to tie the animals in the open during the night. Tie the animals in a shade time place during the day. Do not tie the animals under the tree because this week the chances of falling trees twigs are high due to the speed of the winds. FMD for the prevention of foot-mouth disease to animal's vaccine and VQ for prevention of Lagadia fever. Give sufficient quantity of grains to the animals with green and dry fodder. Animals must be given clean and fresh water 3-4 times a day. Do not tie pregnant animals to the sloping place. The best time to give anthelmintic drugs for the prevention of insects in the stomach of animals.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Farmers are advised to add dietary supplements; vitamins and energy feed ingredients to the hens as well as add calcium content to the hen. Give medicine for the prevention of worms in the stomach of hen (Divmirg). To protect the chickens from heat, arrange curtains, fans and ventilation in the hen house. To save the chickens from the heat, sprinkle water on the curtains engaged in the chicken house so that the coolness remains.

Likely impacts of weather warnings (General)

According to weather information from the India Meteorological Department, conditions such as localized strong winds, and sultry heat are likely to persist over the next five days. Additionally, there is a warning of heatwave conditions.

Impact based advisories (General)

Harvest and thresh the mature Zaid crops and store the grain; the weather is favorable for sowing Kharif maize, preparing paddy nurseries, and transplanting seedlings.

Farmers are advised to download Unified "Mausam" and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: <https://play.google.com/store/apps/details/>

Meghdoot MobileApp link: <https://play.google.com/store/apps/details>

Damini MobileApp link : <https://play.google.com/store/apps/details>