**Gramin Krishi Mausam Sewa**

**Agromet Advisory Bulletin for SITAPUR District**

**Chander Shekhar Azad University of Agriculture & Technology**

Kanpur, Uttar Pradesh

# Agromet Advisory Bulletin

**Date : 2026-10-23**

**Weather Forecast of District SITAPUR (Uttar Pradesh) Issued On : 2026-10-23 (Valid Till 08:30 IST of the next 5 days)**

| Parameter | 2026-10-24 | 2026-10-25 | 2026-10-26 | 2026-10-27 | 2026-10-28 |
|---|---|---|---|---|---|
| Rainfall(mm) | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Tmax(°C) | 31.0 | 31.0 | 32.0 | 32.0 | 31.0 |
| Tmin(°C) | 18.0 | 18.0 | 18.0 | 17.0 | 18.0 |
| RH-I(%) | 86 | 85 | 85 | 86 | 83 |
| RH-II(%) | 45 | 46 | 46 | 45 | 45 |
| Wind Speed(kmph) | 4 | 3 | 1 | 4 | 4 |
| Wind Direction(Degree) | 322 | 30 | 34 | 90 | 63 |
| Cloud Cover(Octa) | 0 | 0 | 0 | 0 | 1 |
| Warning | NA | NA | NA | NA | NA |

## Forecast Summary:

According to the weather forecast received from the Indian Meteorological Department, due to clear sky weather in the next five day there is a no possibility of rain . The maximum temperature is between 31.0- 32.0°C, which is likely to be near normal and the minimum temperature is between 17.0-18.0°C, which is likely to be 2-3°C higher normal. The maximum and minimum range of relative humidity is between 83-86 and 45-46%. Wind direction is south-west, east, north-east and wind speed is between 1.0-4.0 km/h with winds expected to be 1-2 Km/h higher normal.

## Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the India Meteorological Department, there is no warning in this next five days.

## Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Farmers are advised to irrigate standing rice crops as needed, and the weather is favorable for harvesting/threshing and sowing mature crops. Dry the crop thoroughly in the sun before threshing. Smoke cattle sheds to protect livestock from mosquitoes.

## General Advisory:

Complete the harvesting of mature crops immediately and store the grain in safe, dry locations to prevent damage from unseasonal moisture or fungal growth. Due to high daytime temperatures, the topsoil moisture is depleting rapidly, which affects the sowing of Rabi crops. Therefore, ensure that you provide a 'palewa' (light pre-sowing irrigation) to the field before planting mustard, chickpea, or potatoes; this will improve seed germination. Additionally, carry out sowing in the morning or evening rather than during the midday hours. Take advantage of the dry weather to finish sowing mustard and early-maturing potato varieties by the end of October.

## SMS Advisory:

| |
|---|
| Given the completely clear and dry weather, proceed with the harvesting of mature paddy crops, and sow Rabi crops only after providing a pre-sowing irrigation (palewa). |

## Crop Specific Advisory:

| Crop (Stage) | Crop Specific Advisory |
|---|---|
| RICE (Physiological Maturity) | Harvest the crops when 85% golden color appears in the paddy ears and after harvesting, dry the lock in the sun and do the threshing on clear weather. The current weather is favorable for pests, so after flowering, on an average, 2 - 3 odorous pests / hill are visible in the paddy crop, then for its control, Imidacoloroprid 17. 8% SL 350 ml / ha or Burfozin 25% SP 750 ml Make a solution in 500 -600 liters of water at the rate of / hectare and spray it on clear weather. |
| RAPESEED (Vegetative ) | There is a possibility of infestation of hairy caterpillar in the rapeseed crop, so for its prevention, Emamectin benzoate 5% SG at the rate of 200 g / ha is sprayed in 500-600 liters of water. Within 20-22 sowing of rapeseed crop, weeding should be done within 10-22 centimeters distance from plant to plant. In the rapeseed crop, the first irrigation should be done within 25-30 days of sowing and on arrival of oats, make top dressing of urea at the rate of 108 kg / ha. |
| MUSTARD (Sowing) | Saw fly in mustard crop - These pests cause more damage in 7 to 10 days of germination. To prevent this, methyl parathion 2 percent or malathion 5 percent or carbaryl 5 percent powder at the rate of 25 kg per hectare in the morning or evening. The weather is favorable for sowing work at the rate of 3-4 kg seed/ha for sowing any one of the recommended varieties of rye/mustard- Pusa Mustard-34, Pusa Mustard-32, Varuna (V-WRR-2), C5-64 (CS2005-143), Pusa OOM-36 (PDZ-15), BMP-II (DIMR), Azad Mahak, Surekha (KMR-16-02), Rohini, Narendra Rai- 8501, Maya, Vaibhav etc. |
| FIELD PEA (Sowing) | For sowing of any varieties of pea crop recommended Variety - rachana, Malaviya pea-15, Sapna etc., sow at the rate of 100-125 kg seed / hectare. . |
| CHICK PEA (Sowing) | For sowing of any varieties of chickpea, crop recommendation species- Awarodhi, Pusa-256, Pusa-362, K.W.R-108, K-850, K.D.G.-1168 etc. Small grain seeds of 75-80 kg/ hectare for large grain seeds, sow at the rate of 90-100 kg of seed / hectare |

## Horticulture Specific Advisory:

| Horticulture (Stage) | Horticulture Specific Advisory |
|---|---|
| POTATO (Sowing) | Main varieties of potato - Kufri Bahar, Kufri Anand, Kufri Badshah, Kufri Sinduri, Kufri Sutlej, Kufri Evergreen Lalima, Kufri Arun, Kufri and Kufri Pukhraj, Kufri Garim, Kufri Lima, Kufri Lalit, Kufri Sangam, Kufri Neelkanth, Kufri Ganga, Kufri Thar-3, Kufri Mohan, Kufri Neelkanth, Kufri Surya, Kufri Chipsona-1, Kufri Chipsona-3, Kufri Chipsona-4 And the weather is favorable for sowing of Kufri, Friesona etc. To prevent chicken pox in potato crop, spray 100 grams of streptocycline or boric acid in 10 liters of water and spray the tubers on the seeds and do not treat the germinated seeds. |
| VEGETABLE PEA (Sowing) | Sow vegetables like peas, fennel, celery, beetroot, garlic, coriander, spinach, soya, fenugreek, radish, carrot, turnip and leafy vegetables. Raise nursery of tomato, onion, chilli, cauliflower and cabbage. If there is a possibility of water stagnation in the field, then plant cauliflower, brinjal, tomato, chilli etc. on the ridges. If there is an infestation of fruit borer/leaf borer insect in vegetable crops, then for its control, make a solution of neem oil in 1.5 ml/litre water and spray it 3-4 times at an interval of 8-10 days when the sky is clear. |
| PAPAYA (Fruit formation) | Plant new plants in place of destroyed plants in new orchards of mango, guava, amla, lemon etc. After hoeing the banana/papaya gardens, add 25-30 cm high soil around the stem. Install wooden or boss stands in banana/papaya fruit trees. To prevent leaf cutting insect in mango, sprinkle 2% carbaryl 1.5% dust. |

## Live Stock Specific Advisory:

| Live Stock | Live Stock Specific Advisory |
|---|---|
| BUFFALO | Farmers are advised not to tie pregnant buffaloes/cows at sloping places. Feed nutritious fodder and grains to pregnant buffaloes/cows and feed colostrum to the newborn baby for three days. Keep animals in a clean place. Be sure to vaccinate animals to protect them from strangulation. In view of the changing weather, do not tie animals in open places at night and ensure to take care of the health and illness of the animals. Provide adequate amount of grains along with green and dry fodder to the animals. Make sure to feed clean and fresh water to animals 3-4 times a day. |

**Poultry Specific Advisory:**

| Poultry | Poultry Specific Advisory |
|---|---|
| HEN | Farmers are advised to Clean every day at the place where the chickens live. Arrange for clean water and balanced diet to the hens. This is not the right time to raise new chicks for eggs but it is a suitable time for broiler chickens. Add supplementary food, vitamins and energy food material to the chickens' diet and also give calcium material to the chickens. Chlorination of water must be done. Give deworming medicine to kill the worms in the stomach of the chickens. Keep the chicken house dry and keep turning the bedding. |

**Likely impacts of weather warnings (General)**

Prolonged exposure to sunlight and high humidity significantly increase the risk of heat stress and infectious diseases in livestock. High humidity and heat reduce the appetite of buffaloes and cows, potentially leading to a sudden drop in milk production. Stagnant water following recent monsoon rains, combined with rising temperatures, creates an environment where bacteria thrive in fodder and water sources. This can lead to the spread of ailments such as intestinal worms, skin infections, and Foot-and-Mouth Disease among the animals. The combination of a sudden rise in humid heat and a gradual drop in minimum temperatures creates conditions characterized by intense heat during the day and a slight chill at night or in the early morning. This fluctuating weather affects human health; venturing out in strong sunlight can lead to dehydration. Furthermore, this season sees a rapid surge in the prevalence of mosquitoes and viruses responsible for diseases like Chikungunya, Dengue, Malaria, and viral flu.

**Impact based advisories (General)**

Do not tether animals in direct sunlight during the day (between 11 AM and 4 PM); keep them in a well-ventilated, shaded shed instead. Ensure drinking water troughs are kept clean and provide fresh, clean water at least 3–4 times a day. If vaccinations for Hemorrhagic Septicemia (HS) or Black Quarter (BQ) have not been administered, get the animals vaccinated. At night, use neem leaf smoke in the animal shed to protect against mosquitoes and flies.Carry a water bottle when stepping out. Consume ORS, lemonade, or buttermilk to prevent electrolyte depletion in the body. Do not allow rainwater from recent showers to stagnate around the house or on rooftops. Use mosquito nets or repellents while sleeping. Avoid drinking cold water from the refrigerator or sitting directly in air-conditioned (AC) environments immediately after coming in from the heat; these habits are primary causes of colds, coughs, and lung infections. This dry week is ideal for preparing Rabi crops and for the harvesting

**Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.**

**Mausam MobileApp link: https://play.google.com/store/apps/details/**

**Meghdoot MobileApp link: https://play.google.com/store/apps/details**

**Damini MobileApp link : https://play.google.com/store/apps/details**