**Gramin Krishi Mausam Sewa**

**Agromet Advisory Bulletin for ETAWAH District**

**Chander Shekhar Azad University of Agriculture & Technology**

Kanpur, Uttar Pradesh

# Agromet Advisory Bulletin

**Date : 2026-04-17**

**Weather Forecast of District ETAWAH (Uttar Pradesh) Issued On : 2026-04-17 (Valid Till 08:30 IST of the next 5 days)**

| Parameter | 2026-04-18 | 2026-04-19 | 2026-04-20 | 2026-04-21 | 2026-04-22 |
|---|---|---|---|---|---|
| Rainfall(mm) | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Tmax(°C) | 40.0 | 41.0 | 41.0 | 41.0 | 41.0 |
| Tmin(°C) | 23.0 | 23.0 | 23.0 | 24.0 | 23.0 |
| RH-I(%) | 41 | 44 | 34 | 33 | 33 |
| RH-II(%) | 24 | 23 | 22 | 21 | 22 |
| Wind Speed(kmph) | 6 | 7 | 14 | 12 | 13 |
| Wind Direction(Degree) | 63 | 236 | 345 | 306 | 294 |
| Cloud Cover(Octa) | 3 | 1 | 1 | 1 | 1 |
| Warning | No Warning | No Warning | No Warning | No Warning | No Warning |

## Forecast Summary:

According to the weather forecast received from the India Meteorological Department, due to cloudy weather days in the next five days but there is a no possibility of rain . The maximum temperature is between 40.0- 41.0°C, which is likely to be 3-4°C higher normal and the minimum temperature is between 23.0-24.0°C, which is likely to be 1-2°C higher normal. The maximum and minimum range of relative humidity is between 32-44 and 21- 24%. Wind direction is north-east, north-west and wind speed is between 6.0-13.0 kmph with gusts expected to be 3-4 kmph high speed than normal.

## Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the India Meteorological Department, there is a no weather warning in next five days.

## Likely impacts of weather warnings on Agriculture and associated Agromet advisories

In view of the prevailing weather conditions, farmers are advised to undertake irrigation in standing crops such as okra, ridge gourd, pumpkin, bottle gourd, cucumber, maize, black gram, and green gram. Furthermore, the weather is favorable for the harvesting and threshing of mature crops like wheat, gram and pigeon pea.

## General Advisory:

Farmers are advised to ensure that the weather is favorable for harvesting and threshing of Rabi crops. Maintain proper moisture by sowing moong and doing irrigation work in maize, urd and groundnut crops as per requirement. During harvesting of mature crops, farmers/laborers must take water bottles with them to the fields. Send field harvested vegetables like tomato, cauliflower, green leafy, cucumber other cucurbit vegetables to your nearest market for direct marketing. For pesticides, herbicides and herbicide chemicals, wash the equipment separately with clean water only. Or use and do not spray or spray pesticides, insecticides and herbicides while standing in the opposite direction of the wind.

| |
|---|
| Spraying should be done in the evening, if possible, after spraying, hands should be washed thoroughly with soap or hand sanitizer before eating and after washing clothes. |

**SMS Advisory:**

| |
|---|
| Temperature and wind speed are expected to be higher than normal in the coming week, hence farmers should maintain proper moisture by lightly irrigating the crops standing in the fields. |

**Crop Specific Advisory:**

| Crop (Stage) | Crop Specific Advisory |
|---|---|
| WHEAT (Drying / Threshing) | Perform harvesting work by threshing the mature crops of wheat. After harvesting the wheat crop, the bundle must be tied so that the lock does not fly due to strong winds and the threshing should be done in the evening and at night when the wind is cool. For storage, the wheat should be dried under hard sun so that the moisture content does not exceed 8 -10%. Disinfect the spruce or storage house with insecticides before storage. |
| MAIZE (Silking) | Keep the maize crops free from weeds and irrigate them at intervals of 12-15 days as per the requirement and maintain proper moisture. There is a possibility of outbreak of stem borer / shoot fly pest in maize crop, so for its prevention, imamactin benzoate 5% Sg 4 g / ha or dimethoate 30% EC 1.0 liter / ha at 600-700 liter Sprinkle by making. |
| BLACK GRAM (Vegetative ) | Keep the black gram crops free from weeds and irrigate them at intervals of 15-20 days as per the requirement and maintain proper moisture. There is a possibility of an outbreak of thrips / green moth in the northern crop, so for prevention, spraying of Oxidematan-methyl 25% EC or Dimethoate 30% EC at the rate of 1.0 liter / ha in 600-700 liters of water is sprayed. |
| GREEN GRAM (Vegetative ) | In green gram crops, 25-30 days after the first irrigation sowing, and the remaining irrigation should be irrigated at intervals of 12 -15 days as per the requirement and maintain proper moisture. |

**Horticulture Specific Advisory:**

| Horticulture (Stage) | Horticulture Specific Advisory |
|---|---|
| ONION (Physiological Maturity) | In the onion crop for prevention of damping off (wetting) disease spraying of Carbendazim (1 g / liter of water) or Mancozeb (2.5 g / liter of water) is sprayed at 15 days intrawall during evening hours. During the summer season, weeding should be done in the crops of vegetables, tomato, brinjal and chilli. The best time to harvest onions is one week after 50% of the leaves fall. |
| MANGO (Fruit formation) | Do irrigation work in orchards of mango, guava, lemon plum, grapes, papaya and litchi etc. To prevent mango fruit from falling, irrigate the gardens or spray the solution in 15 ppm of acetic acid or 4 ml plenofix per 9 liters of water. To prevent cuckoo disorder or internal rot disease in mango pods, make borax 0.8% solution in 0.8 g / liter water and spray 2 at 10 days interval. |

**Live Stock Specific Advisory:**

| Live Stock | Live Stock Specific Advisory |
|---|---|
| BUFFALO | Considering the current weather, farmers are advised not to tie the animals in the open during the night. Tie the animals in a shade time place during the day. Do not tie the animals under the tree because this week the chances of falling trees twigs are high due to the speed of the winds. Vaccination in animals for prevention of hoof-mouth disease. Give sufficient quantity of grains to the animals with green and dry fodder. Animals must be given clean and fresh water 3-4 times a day. Do not tie pregnant animals to the sloping place. The best time to give anthelmintic drugs for the prevention of insects in the stomach of animals. |

**Poultry Specific Advisory:**

| Poultry | Poultry Specific Advisory |
|---|---|
| HEN | Farmers are advised to add dietary supplements; vitamins and energy feed ingredients to the hens as well as add calcium content to the hen. Give medicine for the prevention of worms in the stomach of hen (Divmirg). To protect the chickens from heat, arrange curtains, fans and ventilation in the hen house. To reduce the temperature of the hen house, cover the roof of the shed with straw. |

## Likely impacts of weather warnings (General)

| |
|---|
| According to the weather forecast received from the India Meteorological Department, there is a no weather warning in next five days. |

## Impact based advisories (General)

| |
|---|
| In view of the prevailing weather conditions, farmers are advised to undertake irrigation in standing crops such as okra, ridge gourd, pumpkin, bottle gourd, cucumber, maize, black gram, and green gram. Furthermore, the weather is favorable for the harvesting and threshing of mature crops like wheat, gram and pigeon pea. |

**Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.**

**Mausam MobileApp link: https://play.google.com/store/apps/details/**

**Meghdoot MobileApp link: https://play.google.com/store/apps/details**

**Damini MobileApp link : https://play.google.com/store/apps/details**