Gramin Krishi Mausam Sewa

Agromet Advisory Bulletin for KANNAUJ District

Chander Shekhar Azad University of Agriculture & Technology

Kanpur, Uttar Pradesh

Agromet Advisory Bulletin

Date : 2026-11-03

Weather Forecast of District KANNAUJ (Uttar Pradesh) Issued On : 2026-11-03 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-11-04	2026-11-05	2026-11-06	2026-11-07	2026-11-08
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	31.0	32.0	32.0	32.0	32.0
Tmin(°C)	19.0	19.0	19.0	19.0	19.0
RH-I(%)	49	44	42	46	52
RH-II(%)	21	19	21	22	22
Wind Speed(kmph)	6	7	5	5	4
Wind Direction(Degree)	313	276	288	266	292
Cloud Cover(Octa)	0	0	0	0	0
Warning	No Warning	No Warning	No Warning	No Warning	No Warning

Forecast Summary:

According to the weather forecast received from the Indian Meteorological Department, due to clear sky weather in the next five day there is a no possibility of rain. The maximum temperature is between 31.0-32.0°C, which is likely to be 2-3°C higher normal and the minimum temperature is between 18.0-19.0°C, which is likely to be 3-4°C higher normal. The maximum and minimum range of relative humidity is between 42-52 and 18-19%. Wind direction is north-west, south-west and wind speed is between 4.0-7.0 km/h with winds expected to be 3-4 Km/h higher normal.

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast of the India Meteorological Department, there is a no warning in the next five days.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

With the weather remaining completely dry and clear, conditions are highly favorable for the harvesting and threshing of the paddy crop, as well as for the sowing of Rabi crops (wheat, mustard, and potatoes). Given the increase in humidity due to light morning fog, regularly monitor vegetable crops for pests and fungal diseases. Farmers are advised to ensure seed treatment before sowing and to use 'Pusa Decomposer' in their fields instead of burning crop residue (stubble).

General Advisory:

Due to the completely dry weather, this is the most suitable time for sowing wheat, mustard, potatoes, and peas. Be sure to treat the seeds before sowing. Complete the harvesting and threshing of the standing paddy crop immediately. Store the harvested produce in a safe, dry place to protect it from moisture, and ensure the grain is thoroughly dried before storage (keeping the moisture content below 12%). Do not burn crop residue (stubble) in the fields under any circumstances; instead, spray 'Pusa Decomposer' to break down the residue within the field itself, which will help enhance soil fertility. Given the morning fog and increasing humidity, regularly monitor late-sown crops—such as green chilies, tomatoes, and cauliflower—for the presence of pests and fungal diseases.

SMS Advisory:

Take advantage of the dry weather to complete the paddy harvest, begin sowing Rabi crops immediately, and do not burn stubble in your fields.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
RICE (Drying / Threshing)	Harvest the paddy crops and after drying for 2 -3 days after harvesting, and after harvesting, dry the sun for 3-4 days and store the seeds.
WHEAT (Sowing)	According to the weather, the temperature is favorable for sowing of wheat in the irrigated condition, so the sowing of wheat should start from 10 November. Prepare field for sowing wheat and regional species PBW-723, HPBW-01, DBW-39, K.-9107, K.-1006, K.-402, K.-607, D. .BW-90, WB-02, NW-5054, HD-3043, U.P-2382, H.U. W-468, PBW-443, HD-2967 etc.

Crop (Stage)	Crop Specific Advisory
	Recommendation species for erosive land: - KRL-210, KRL-213, K.- 1317 etc. for the sowing of any species of manure and seeds should be arranged.
RAPESEED (Flowering)	There is a possibility of infestation of hairy caterpillar in the rapeseed crop, so for its prevention, Emamectin benzoate 5% SG at the rate of 200 g / ha is sprayed in 500-600 liters of water.
MUSTARD (Sowing)	Saw fly in mustard crop - These pests cause more damage in 7 to 10 days of germination. To prevent this, methyl parathion 2 percent or malathion 5 percent or carbaryl 5 percent powder at the rate of 25 kg per hectare in the morning or evening. It on clear weather for sowing work at the rate of 3-4 kg seed/ha for sowing any one of the recommended varieties of rye/mustard- Pusa Mustard-34, Pusa Mustard-32, Varuna (V-WRR-2), C5-64 (CS2005-143), Pusa OOM-36 (PDZ-15), BMP-II (DIMR), Azad Mahak, Surekha (KMR-16-02), Rohini, Narendra Rai- 8501, Maya, Vaibhav etc.
FIELD PEA (Sowing)	For sowing of any varieties of pea crop recommended Variety - rachana, Malaviya pea-15, Sapna etc., sow at the rate of 100-125 kg seed / hectare.
CHICK PEA (Sowing)	For sowing of any varieties of chickpea, crop recommendation species- Awarodhi, Pusa-256, Pusa-362, K.W.R-108, K-850, K.D.G.-1168 etc. Small grain seeds of 75-80 kg/ hectare for large grain seeds, sow at the rate of 90-100 kg of seed / hectare.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
POTATO (Sowing)	Main varieties of potato - Kufri Bahar, Kufri Anand, Kufri Badshah, Kufri Sinduri, Kufri Sutlej, Kufri Evergreen Lalima, Kufri Arun, Kufri and Kufri Pukhraj, Kufri Garim, Kufri Lima, Kufri Lalit, Kufri Sangam, Kufri Neelkanth, Kufri Ganga, Kufri Thar-3, Kufri Mohan, Kufri Neelkanth, Kufri Surya, Kufri Chipsona-1, Kufri Chipsona-3, Kufri Chipsona-4 And the weather is favorable for sowing of Kufri, Friesona etc. To prevent chicken pox in potato crop, spray 100 grams of streptocycline or boric acid in 10 liters of water and spray the tubers on the seeds and do not treat the germinated seeds.
VEGETABLE PEA (Sowing)	Sow vegetables like peas, fennel, celery, beetroot, garlic, coriander, spinach, soya, fenugreek, radish, carrot, turnip and leafy vegetables. Raise nursery of tomato, onion, chilli, cauliflower and cabbage. If there is a possibility of water stagnation in the field, then plant cauliflower, brinjal, tomato, chilli etc. on the ridges. If there is an infestation of fruit borer/leaf borer insect in vegetable crops, then for its control, make a solution of neem oil in 1.5 ml/litre water and spray it 3-4 times at an interval of 8-10 days.
PAPAYA (Transplanting)	Plant new plants in place of destroyed plants in new orchards of mango, guava, amla, lemon etc. After hoeing the banana/papaya gardens, add 25-30 cm high soil around the stem. Install wooden or boss stands in banana/papaya fruit trees. To prevent leaf cutting insect in mango, sprinkle 2% carbaryl 1.5% dust.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	Farmers are advised not to tie pregnant buffaloes/cows at sloping places. Feed nutritious fodder and grains to pregnant buffaloes/cows and feed colostrum to the newborn baby for three days. Keep animals in a clean place. Be sure to vaccinate animals to protect them from strangulation. In view of the changing weather, do not tie animals in open places at night and ensure to take care of the health and illness of the animals. Provide adequate amount of grains along with green and dry fodder to the animals. Make sure to feed clean and fresh water to animals 3-4 times a day.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Farmers are advised to Clean every day at the place where the chickens live. Arrange for clean water and balanced diet to the hens. This is not the right time to raise new chicks for eggs but it is a suitable time for broiler chickens. Add supplementary food, vitamins and energy food material to the chickens' diet and also give calcium material to the chickens. Chlorination of water must be done. Give deworming medicine to kill the worms in the stomach of the chickens. Keep the chicken house dry and keep turning the bedding.

Likely impacts of weather warnings (General)

The weather will remain entirely dry and clear, with no adverse weather-related impacts expected on the general public or livestock. Visibility may be slightly affected during the morning and night hours due to light to moderate fog; therefore, please exercise caution while traveling. A gradual drop in temperature will lead to cooler nights, so the public should remain vigilant to protect themselves against seasonal illnesses. Livestock owners are advised to keep milch animals in a secure shed rather than tying them out in the open during the night, helping them avoid the stress caused by changing weather conditions.

Impact based advisories (General)

Light to moderate fog during the morning hours may reduce visibility on roads; therefore, drivers should use fog lights and maintain a controlled speed. Falling temperatures at night and in the morning may exacerbate health issues for those suffering from colds, flu, or respiratory conditions—please ensure you wear light, warm clothing during these times. To protect livestock from the cold during this changing weather, do not keep them tied in the open at night, and ensure their enclosures remain dry and well-ventilated. Take advantage of the clear, sunny weather to complete the harvesting of the mature paddy crop immediately, and begin sowing Rabi crops (such as wheat, potatoes, and mustard) without delay.

Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: https://play.google.com/store/apps/details/

Meghdoot MobileApp link: https://play.google.com/store/apps/details

Damini MobileApp link : https://play.google.com/store/apps/details