



Gramin Krishi Mausam Sewa
Agromet Advisory Bulletin for HATHRAS District
Chander Shekhar Azad University of Agriculture & Technology
Kanpur, Uttar Pradesh



Agromet Advisory Bulletin

Date : 2026-05-12

Weather Forecast of District HATHRAS (Uttar Pradesh) Issued On : 2026-05-12 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-05-13	2026-05-14	2026-05-15	2026-05-16	2026-05-17
Rainfall(mm)	6.0	3.0	0.0	0.0	0.0
Tmax(°C)	37.0	39.0	41.0	41.0	41.0
Tmin(°C)	24.0	25.0	26.0	26.0	26.0
RH-I(%)	72	70	57	53	50
RH-II(%)	44	42	42	42	42
Wind Speed(kmph)	8	11	4	5	11
Wind Direction(Degree)	121	94	280	315	286
Cloud Cover(Octa)	5	5	2	1	1
Warning	Thunderstorms & Lightning, Squall etc; Strong Surface Winds	No Warning	No Warning	No Warning	No Warning

Forecast Summary:

According to the weather forecast received from the Indian Meteorological Department, due to light to moderate cloud cover in the next five days, there is a possibility light rain dated on 13-14 May, 2026 with thunderstorms & lightning, strong surface winds. The maximum temperature is between 37.0-41.0 °C, which is likely to be near its normal and the minimum temperature is between 24.0-26.0 °C, which is likely to be 1-2 °C higher normal. The maximum and minimum range of relative humidity is between 50-72 and 42-44%. Wind direction is south-east, north-east, north-west and wind speed is between 4.0-11.0 kmph, with gusts expected to be 3-4 kmph higher speed than normal.

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the India Meteorological Department, a warning has been issued for 13-14 May, 2026, light rain, thunderstorms accompanied by lightning, strong surface winds.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

In view of the weather conditions, farmers working in the fields are advised not to take shelter or allow themselves or their animals to take shelter under trees in case of heavy winds/storms or rain. Suspend irrigation activities for standing crops such as summer maize, black gram (Urad), and green gram (Moong). Furthermore, harvest mature crops such as wheat and pigeon pea (Arhar) and store them in a safe location.

General Advisory:

Farmers are advised to avoid stepping out of their homes unnecessarily between 11:00 AM and 4:00 PM to protect themselves from the heat (heatwave) and hot winds. While working in the fields, they should keep their heads covered with a cloth (Gamcha) and ensure they carry a water bottle with them. In view of the severe heat, livestock should be given water 3–4 times a day and bathed in the morning and evening. When using chemical pesticides, fungicides, or herbicides, use separate equipment for each category; alternatively, thoroughly wash the equipment with clean water before switching between different chemicals. Furthermore, do not spray or dust these chemicals while standing against the direction of the wind. Whenever possible, spraying operations should be carried out during the evening hours. After spraying, and before eating or drinking, one must thoroughly wash their hands with soap or hand wash; additionally, one should wash their clothes and take a bath.

SMS Advisory:

Farmers are advised to postpone irrigation work. Do not seek shelter or allow yourself or your animals to take refuge under trees during heavy rain and strong winds/storms.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
MAIZE (Silking)	Maize crop is going on at the stage of cumin extract /cob formation/grain filling which is sensitive to moisture deficiency. Therefore, maintain proper moisture by giving light irrigation as per requirements. There is a possibility of outbreak of stem borer / shoot fly pest in maize crop, so for its prevention, solution of Emamectin Benzoate 5% SG at the rate of 200 g/ha or Dimethoate 30% EC 1.0 l/ha in 600-700 liters of water make a spray.
BLACK GRAM (Flowering)	Black gram crop is progressing from flower to pod formation stage which is sensitive to moisture deficiency. Therefore, maintain proper moisture by applying light irrigation as per requirement. There is a possibility of outbreak of pod borer in black gram crop, so to prevent it, apply Emamectin Benzoate 5% SG 220 g / hectare or Dimethoate 30% EC. Spray at the rate of 1.0 liters/hectare in 600-700 liters of water when the sky is clear.
GREEN GRAM (Vegetative)	In green gram crop, maintain proper moisture by giving light irrigation at intervals of 10 -12 days as per requirement. Thrips/green leaf pest is likely to be seen in green gram crop, so for its prevention, spray Oxidemetan-methyl 25% EC or Dimethoate 30% EC at the rate of 1.0 liter/ha by making a solution in 600-700 liters of water.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
BRINJAL (Physiological Maturity)	Harvest the finished crops like Pumpkin, Gourd, Taro, Bitter gourd, Cucumber, Cucurbit, Watermelon and Melon etc. and send them to the market. For the prevention of fruit borer / leaf borer pest in vegetable crops, make a solution of neem oil in 1.5-2.0 ml / liter of water and do 3-4 sprays at an interval of 8 -10 days. Do weeding and hoeing in the standing crops of vegetables and do irrigation work at an interval of 8-10 days in the evening.
MANGO (Physiological Maturity)	Dig pits for planting new gardens. Do irrigation work in orchards of mango, guava, lemon plum, grapes, papaya and litchi etc. Irrigate the orchards to prevent fall of mango fruit or apply 4.5 SL of 20 ml alpha naphthalene acetic acid make a solution per liter of water and spray it. To prevent cuckoo disorder or internal rot disease in mango pods, make borax 0.8% solution in 0.8 g / liter water and spray 2 at 10 days interval.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	Considering the current weather, farmers are advised to tie the animals in the open during the night. Tie the animals in a shade time place during the day. Do not tie the animals under the tree because this week the chances of falling trees twigs are high due to the speed of the winds. FMD for the prevention of foot-mouth disease to animal's vaccine and VQ for prevention of Lagadia

Live Stock	Live Stock Specific Advisory
	fever. Give sufficient quantity of grains to the animals with green and dry fodder. Animals must be given clean and fresh water 3-4 times a day. Do not tie pregnant animals to the sloping place. The best time to give anthelmintic drugs for the prevention of insects in the stomach of animals.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Farmers are advised to add dietary supplements; vitamins and energy feed ingredients to the hens as well as add calcium content to the hen. Give medicine for the prevention of worms in the stomach of hen (Divmirg). To protect the chickens from heat, arrange curtains, fans and ventilation in the hen house. To save the chickens from the heat, sprinkle water on the curtains engaged in the chicken house so that the coolness remains.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation)	Others (Soil / Land Preparation) Specific Advisory
AGRI FIELD	Deep plow the fields empty of wheat and other crops with disc plow so that the eggs and larvae of harmful insects hidden inside the ground are destroyed in the sun.

Likely impacts of weather warnings (General)

According to the weather forecast received from the India Meteorological Department, a warning has been issued for 13-14 May, 2026, light rain, thunderstorms accompanied by lightning, strong surface winds.

Impact based advisories (General)

In view of the weather conditions, farmers working in the fields are advised not to take shelter or allow themselves or their animals to take shelter under trees in case of heavy winds/storms or rain. Suspend irrigation activities for standing crops such as summer maize, black gram (Urad), and green gram (Moong). Furthermore, harvest mature crops such as wheat and pigeon pea (Arhar) and store them in a safe location.

Farmers are advised to download Unified "Mausam" and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: <https://play.google.com/store/apps/details/>

Meghdoot MobileApp link: <https://play.google.com/store/apps/details>

Damini MobileApp link : <https://play.google.com/store/apps/details>