



Agromet Advisory Bulletin

Date : 2026-10-06

Weather Forecast of District AURAIYA (Uttar Pradesh) Issued On : 2026-10-06 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-10-07	2026-10-08	2026-10-09	2026-10-10	2026-10-11
Rainfall(mm)	1.0	1.0	0.0	0.0	0.0
Tmax(°C)	34.0	34.0	35.0	34.0	34.0
Tmin(°C)	24.0	25.0	25.0	24.0	24.0
RH-I(%)	81	84	84	81	79
RH-II(%)	47	48	43	46	43
Wind Speed(kmph)	9	8	7	6	10
Wind Direction(Degree)	138	121	168	114	152
Cloud Cover(Octa)	3	2	0	0	0
Warning	No Warning	Thunderstorms & Lightning, Squall etc; Strong Surface Winds	No Warning	No Warning	No Warning

Forecast Summary:

According to the India Meteorological Department's forecast, light rain accompanied by gusty winds and thunderstorms is likely at the local level during the first two days due to partly cloudy skies, while no rainfall is expected during the remaining days as the weather will be clear. The maximum temperature is expected to range between 34.0°C to 35.0°C potentially 1–2°C above normal), and the minimum temperature is likely to range between 24.0°C and 25.0°C (potentially 3–4°C above normal). Maximum and minimum relative humidity levels will range between 79–84% and 43–48%, respectively. Wind directions will be north-west, south-east and south-west with speeds ranging from 6.0 to 10.0 km/h; wind gusts may exceed the sustained wind speed by 2–3 km/h.

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the India Meteorological Department (IMD), there is a first two days warning of thunderstorms, strong winds light rain.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Farmers are advised to carry out the harvesting, threshing, and sowing of mature crops only when there is no rainfall. Do not tether livestock in open areas or let them out to graze, and prevent them from entering waterlogged areas. Keep animals in a dry, safe place and provide them with clean drinking water; avoid tethering them under trees or near electric poles, as falling branches or electric shocks from the poles could prove fatal.

General Advisory:

In the absence of rain, proceed with the harvesting, threshing, and sowing operations for mature Kharif crops such as paddy, maize, and sesame. Carry out the sowing of Rabi crops—such as gram, peas, lentils, mustard, and linseed—only after treating the seeds. Harvest timely-sown maize and paddy crops when 85% of the crop has turned golden in color.

SMS Advisory:

Farmers are advised to carry out the harvesting and threshing of mature Kharif crops and the sowing of Rabi crops in the absence of rain.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
RICE (Soft Dough)	Harvest the crops when the earrings start showing 80% golden color and after harvesting, dry the lock in the sun and do the threshing work on clear weather. The current weather is favorable for pests, so after flowering, on an average, 2 - 3 odorous pests / hill are visible in the paddy crop, then for its control, Imidacoloroprid 17. 8% SL 350 ml / ha or Burfozin 25% SP 750 ml Make a solution in 500 -600 liters of water at the rate of / hectare and spray it on clear weather.

Crop (Stage)	Crop Specific Advisory
GROUNDNUT (Physiological Maturity)	Dig groundnuts only when the veins appear on the top of the skin and the inner part becomes brown in color and the grain of the groundnut becomes pink. Groundnut crop is likely to show white girdar / termite pest infestation, hence fipronil 0.3% GR at the rate of 20 kg / ha topdressing do in clear weather.
BLACK GRAM (Harvesting)	Dry the crop thoroughly in the sun after last week's rain and then thresh it. The weather is favorable for harvesting/threshing of mature black gram/green gram crops.
RAPSEED (Vegetative)	Within 20-22 sowing of rapeseed crop, weeding should be done within 10-22 centimeters distance from plant to plant.
MUSTARD (Sowing)	For sowing, select one of the recommended mustard varieties—such as Varuna, Rohini, Narendra Rai-8501, Maya, Vaibhav, or hybrids like Pioneer 45S46 and Shriram DCM-1666—and sow at a rate of 3–4 kg of seed per hectare.
CHICK PEA (Sowing)	For sowing of any varieties of chickpea, crop recommendation species-Awarodhi, Pusa-256, Pusa-362, K.W.R-108, K-850, K.D.G.-1168 etc. Small grain seeds of 75-80 kg/ hectare for large grain seeds, sow at the rate of 90-100 kg of seed / hectare keep the sowing work.
SUGARCANE (Grand growth)	Destroy the plants affected by red rot disease by removing them along with the roots and spray the plants with a solution of thiophanate methyl 0.2 percent. If symptoms of bacterial rot disease are seen in sugarcane crop, cut and destroy the infected plant and for its control, spray 0.01% Streptomycin when weather is favorable.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
POTATO (Sowing)	Prepare the fields for potato sowing and start sowing. To protect the potato crop from pox disease, dissolve 100 grams of streptomycin or boric acid in 10 liters of water and spray it on the seeds and do not treat the germinated seeds. Take out the potato seeds from the cold storage 7-10 days before sowing. If sprouts appear in the seeds stored in the cold storage, then sort them out and separate them. Sow early potato varieties like Kufri Ashoka, Kufri Chandramukhi, Kufri Jawahar etc.
CAULIFLOWER (Transplanting)	Sow vegetables like peas, fennel, celery, beetroot, garlic, coriander, spinach, soya, fenugreek, radish, carrot, turnip and leafy vegetables. Raise nursery of tomato, onion, chilli, cauliflower and cabbage. If there is a possibility of water stagnation in the field, then plant cauliflower, brinjal, tomato, chilli etc. on the ridges. If there is an infestation of fruit borer/leaf borer insect in vegetable crops, then for its control, make a solution of neem oil in 1.5 ml/litre water and spray it 3-4 times at an interval of 8-10 days when the sky is clear.
BANANA (Flowering)	Try to complete the planting of fruit plants like mango, guava, amla, lemon etc. as soon as possible. Plant new plants in place of the destroyed plants in the new bag. Make a stand of 25-30 meters high soil around the stem by folding banana / papaya bags. For prevention of mango leaf moth, apply 2% carburil 1.5% dust .

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	Farmers are advised to Pregnant buffaloes/cows should not be tied to the slope, feed nutritious fodder and feed to pregnant buffaloes/cows and feed milk to the newborn child for three days. Keep the animals in a clean place and smoke the animals to prevent fly and mosquito. Get animals vaccinated for the prevention of cracking-mouth disease. Animals suffering from this disease should wash their wounds with potassium permanganate. Give sufficient quantity of grains with green and dry fodder to the animals. Do not feed the stagnant water of the ponds or other places, due to which the animals are prone to liver flu and stomach worms. The animals must give clean and fresh water 3-4 times a day.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Farmers are advised to Clean every day at the place where the chickens live. Arrange for clean water and balanced diet to the hens. This is not the right time to raise new chicks for eggs but it is a suitable time for broiler chickens. Add supplementary food, vitamins and energy food material to the chickens' diet and also give calcium material to the chickens. Chlorination of water must be done. Give deworming medicine to kill the worms in the stomach of the chickens. Keep the chicken house dry and keep turning the bedding. If there is excess moisture in the chicken house, turn on the fan.

Likely impacts of weather warnings (General)

Prolonged exposure to sunlight and high humidity significantly increase the risk of heat stress and infectious diseases in livestock. High humidity and heat reduce the appetite of buffaloes and cows, potentially leading to a sudden drop in milk production. Stagnant water following recent monsoon rains, combined with rising temperatures, creates an environment where bacteria thrive in fodder and water sources. This can lead to the spread of ailments such as intestinal worms, skin infections, and Foot-and-

Mouth Disease among the animals. The combination of a sudden rise in humid heat and a gradual drop in minimum temperatures creates conditions characterized by intense heat during the day and a slight chill at night or in the early morning. This fluctuating weather affects human health; venturing out in strong sunlight can lead to dehydration. Furthermore, this season sees a rapid surge in the prevalence of mosquitoes and viruses responsible for diseases like Chikungunya, Dengue, Malaria, and viral flu.

Impact based advisories (General)

Do not tether animals in direct sunlight during the day (between 11 AM and 4 PM); keep them in a well-ventilated, shaded shed instead. Ensure drinking water troughs are kept clean and provide fresh, clean water at least 3–4 times a day. If vaccinations for Hemorrhagic Septicemia (HS) or Black Quarter (BQ) have not been administered, get the animals vaccinated. At night, use neem leaf smoke in the animal shed to protect against mosquitoes and flies. Carry a water bottle when stepping out. Consume ORS, lemonade, or buttermilk to prevent electrolyte depletion in the body. Do not allow rainwater from recent showers to stagnate around the house or on rooftops. Use mosquito nets or repellents while sleeping. Avoid drinking cold water from the refrigerator or sitting directly in air-conditioned (AC) environments immediately after coming in from the heat; these habits are primary causes of colds, coughs, and lung infections. This dry week is ideal for preparing Rabi crops and for the harvesting

Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: <https://play.google.com/store/apps/details/>

Meghdoot MobileApp link: <https://play.google.com/store/apps/details>

Damini MobileApp link : <https://play.google.com/store/apps/details>