**Gramin Krishi Mausam Sewa**

**Agromet Advisory Bulletin for KANPUR-NAGAR District**

**Chander Shekhar Azad University of Agriculture & Technology**

Kanpur, Uttar Pradesh

# Agromet Advisory Bulletin

**Date : 2026-04-07**

**Weather Forecast of District KANPUR-NAGAR (Uttar Pradesh) Issued On : 2026-04-07 (Valid Till 08:30 IST of the next 5 days)**

| Parameter | 2026-04-08 | 2026-04-09 | 2026-04-10 | 2026-04-11 | 2026-04-12 |
|---|---|---|---|---|---|
| Rainfall(mm) | 9.0 | 9.0 | 0.0 | 0.0 | 0.0 |
| Tmax(°C) | 33.0 | 33.0 | 34.0 | 36.0 | 37.0 |
| Tmin(°C) | 20.0 | 18.0 | 19.0 | 21.0 | 22.0 |
| RH-I(%) | 64 | 81 | 76 | 53 | 47 |
| RH-II(%) | 42 | 39 | 35 | 28 | 28 |
| Wind Speed(kmph) | 5 | 11 | 5 | 10 | 11 |
| Wind Direction(Degree) | 74 | 60 | 8 | 332 | 294 |
| Cloud Cover(Octa) | 6 | 5 | 1 | 0 | 0 |
| Warning | Thunderstorms & Lightning, Squall etc; Strong Surface Winds | Thunderstorms & Lightning, Squall etc; Hailstorm; Strong Surface Winds | No Warning | No Warning | No Warning |

## Forecast Summary:

According to the weather forecast from the India Meteorological Department, due to light to moderate cloud cover in the next three days, there is a possibility of light to moderate rain with local level at high winds speeds, thunderstorms & Lightning, hailstorm dated on 08-09 April, 2026. The maximum temperature is between 33.0-37.0°C, which is likely to be 1-2°C lower normal and the minimum temperature is between 18.0-22.0°C, which is likely to be 1-2°C higher normal. The maximum and minimum range of relative humidity is between 47-81 and 28-42%. The wind direction will be from the north-east, north-west and the wind speed will be between 5.0-11.0 km/hour, which is expected to be 3-4 km/hour higher than normal.

## Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the India Meteorological Department, there is a light to moderate rain, Thunderstorms & Lightning, Strong Surface Winds, Hailstorm weather warning dated on 08-09 April, 2026.

## Likely impacts of weather warnings on Agriculture and associated Agromet advisories

In view of the prevailing weather conditions, suspend the sowing and irrigation of crops such as okra, ridge gourd, pumpkin, bottle gourd, cucumber, snake cucumber, summer maize, black gram, and green gram. Furthermore, harvest mature crops—such as wheat, chickpea, pigeonpea and store them in a safe location.

## General Advisory:

| |
|---|
| Farmers are advised to make proper arrangements for the drainage of excess rainwater. They should gather harvested crops and cover them with polythene sheets; if harvested crops have become wet due to rain, they should be thoroughly dried in the sun before threshing, after which the grains should be stored safely. Irrigation activities for standing Zaid crops should be skiped. Do not spray insecticides, fungicides, or herbicides while standing against the direction of the wind. Spraying should ideally be carried out in the evening; furthermore—if possible—hands should be thoroughly washed with soap after spraying, before eating, and after washing clothes. |

## SMS Advisory:

| |
|---|
| Farmers are advised to thoroughly dry harvested crops in the sun if they have become wet before threshing, and to postpone irrigation and sowing activities. |

## Crop Specific Advisory:

| Crop (Stage) | Crop Specific Advisory |
|---|---|
| WHEAT (Drying / Threshing) | In view of the possibility of rain, Farmers are advised to harvested crop should be threshing before rain or keep safer place. In the wheat crop, when the outbreak of mice is seen, use fodder made of zinc phosphide or aluminum phosphide cakes collectively try to prevent mice. |
| MAIZE (Vegetative ) | In view of the possibility of rain, postpone irrigation. In maize crop, do the first irrigation after 20-25 days of sowing and do weeding and hoeing after the oat arrives. For the prevention of stem borer pest in maize crop, apply phorate 10% g 10 kg or carbofuran 3% g 20 kg / ha after 10 -15 days after sowing. |
| BLACK GRAM (Ear Emergence) | Rainfall possibility during this week, farmers are advised to postpone irrigation. |
| GREEN GRAM (Sowing) | Rainfall forecast during this week farmers are advised sowing should be done with recommended castes of Zaid Moong - Narendra Moong-1, Emperor (PDM-139), Malviya Janapriya (HUM-6), Meha, Malviya Janakalyani, Malviya Jan Chetna, Sweta, Swati, Virat, Shikha, IPM-2-3, IPM-2-14, Malviya Jyoti etc. For sowing of any one variety with 20-25 kg/ha seed. |

## Horticulture Specific Advisory:

| Horticulture (Stage) | Horticulture Specific Advisory |
|---|---|
| OKRA/ LADYFINGER (Sowing) | In view of the possibility of rain, postpone the work of irrigation. If seen purple spot disease in onion crop affects the condition at 28 - 30 ° C temperature and 80 - 90% relative humidity, spray copper oxychloride 0.3% (3 g/liter water) or Mancozeb at the rate of 0.25% (0.25 g/liter of water) 3-4 spray at 15-20 day interval in the clear sky weather. Sown vegetables such as ladyfinger, zucchini, cucumber, bitter gourd, gourd and pumpkin etc. sown during summer season. Transplant the seedlings of tomato, brinjal and chilli. |
| MANGO (Fruit formation) | Irrigate the orchards or spray with acetic acid solution of 15 ppm or 4 ml Planofix per 9 liters of water to prevent drop of mango fruits. There is a possibility of showing the outbreak of midge pest in mango blossom, so for its prevention, spray Fenitrothian in 1.0 ml / liter of water and spray it. |

## Live Stock Specific Advisory:

| Live Stock | Live Stock Specific Advisory |
|---|---|
| BUFFALO | It is the appropriate time to give anthelmintic medicine to the animals for the prevention of worms in the stomach. Make OT mixture of 750 grams mustard oil, 100 grams turmeric, 50 grams dry ginger, 50 grams cumin, 500 grams jaggery after feeding the animals for three days in the morning and evening. In view of the prevailing weather, farmers are advised not to tie the animals in the open during the night. Tie the animals in a shaded place or in the shade of a tree during the day. Give sufficient amount of grains along with green and dry fodder to the animals. The animals must be given clean and fresh water 3-4 times a day. |

**Poultry Specific Advisory:**

| Poultry | Poultry Specific Advisory |
|---|---|
| HEN | Farmers are advised to add dietary supplements; vitamins and energy feed ingredients to the hens as well as add calcium content to the chickens. Give medicine for the prevention of worms in the stomach of chickens (Divmirg). |

**Likely impacts of weather warnings (General)**

| |
|---|
| According to the weather forecast received from the India Meteorological Department, there is a light to moderate rain, Thunderstorms & Lightning, Strong Surface Winds, Hailstorm weather warning dated on 08-09 April, 2026. |

**Impact based advisories (General)**

| |
|---|
| In view of the prevailing weather conditions, suspend the sowing and irrigation of crops such as okra, ridge gourd, pumpkin, bottle gourd, cucumber, snake cucumber, summer maize, black gram, and green gram. Furthermore, harvest mature crops—such as wheat, chickpea, pigeonpea and store them in a safe location. |

**Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.**

**Mausam MobileApp link: https://play.google.com/store/apps/details/**

**Meghdoot MobileApp link: https://play.google.com/store/apps/details**

**Damini MobileApp link : https://play.google.com/store/apps/details**