Gramin Krishi Mausam Sewa

Agromet Advisory Bulletin for UNNAO District

Chander Shekhar Azad University of Agriculture & Technology

Kanpur, Uttar Pradesh

Agromet Advisory Bulletin

Date : 2026-04-13

Weather Forecast of District UNNAO (Uttar Pradesh) Issued On : 2026-04-13 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-04-14	2026-04-15	2026-04-16	2026-04-17	2026-04-18
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	36.0	38.0	40.0	41.0	42.0
Tmin(°C)	20.0	21.0	22.0	23.0	23.0
RH-I(%)	56	51	54	58	57
RH-II(%)	31	30	30	29	29
Wind Speed(kmph)	10	10	10	6	6
Wind Direction(Degree)	325	322	335	7	23
Cloud Cover(Octa)	0	0	0	1	2
Warning	No Warning	No Warning	No Warning	No Warning	No Warning

Forecast Summary:

According to the weather forecast received from the India Meteorological Department, due to clear sky weather in the next three days and light cloud cover in remaining days, there is a no possibility of rain . The maximum temperature is between 36.0-42.0°C, which is likely to be near 2-3°C higher normal and the minimum temperature is between 20.0-23.0°C, which is likely to be 1-2°C higher normal. The maximum and minimum range of relative humidity is between 51-58 and 29-31%. The wind direction will be from the north-west, north-east and the wind speed will be between 6.0-10.0 km/hour, which is expected to be 2-3 km/hour higher than normal.

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the India Meteorological Department, there is a no weather warning in next five days.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

In view of the prevailing weather conditions, farmers are advised to undertake irrigation in standing crops such as okra, ridge gourd, pumpkin, bottle gourd, cucumber, maize, black gram, and green gram. Furthermore, the weather is favorable for the harvesting and threshing of mature crops like wheat, gram, peas, lentils, and pigeon pea.

General Advisory:

Farmers are advised to undertake the harvesting and threshing of mature crops—such as wheat, mustard, gram, peas, and lentils—as well as the sowing of green gram (Moong), and irrigation for

maize, black gram (Urad), and groundnut crops. During the harvesting of mature crops, farmers and laborers must carry a water bottle with them to the fields. Harvest ready vegetables—including tomatoes, cauliflower, leafy greens, cucumbers, and other cucurbit crops—should be sent directly to the nearest market or *mandi* for sale. When handling insecticides, fungicides, and herbicides, use separate equipment—or dedicate specific tools solely for this purpose—and ensure that spraying is never performed while standing downwind (against the direction of the wind). Spraying should ideally be conducted in the evening; furthermore, after spraying—and specifically before eating or changing clothes—hands must be thoroughly washed using soap or a hand wash solution.

SMS Advisory:

The weather conditions are favorable for the harvesting and threshing of mature crops. Irrigation should be carried out in standing Zaid crops as and when required.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
WHEAT (Drying / Threshing)	Perform harvesting work by threshing the mature crops of wheat. After harvesting the wheat crop, the bundle must be tied so that the lock does not fly due to strong winds and the threshing should be done in the evening and at night when the wind is cool.
MAIZE (Vegetative)	Keep the maize crops free from weeds. If an infestation of shoot fly/stem borer insect is seen in the maize crop, for its prevention, spray Dimethoate 30% EC 1.0 litre/hectare or Chlorantraniliprole 400 to 500 ml/hectare or Quinalphos 25% EC 2.0 litre/hectare by making a solution in 500-700 litres of water. For prevention of stem borer insect in the maize crop, spray Forate 10% G 10 kg or Carbofuran 3% G 20 kg or Quinalphos 25% EC 2.0 litre/hectare by making a solution in 500-600 litres of water 10-15 days after sowing.
BLACK GRAM (Vegetative)	Keep the crops of black gram free from weeds. There is a possibility of an outbreak of thrips / green moth in the urd crop, so for its prevention, oxidation-methyl 25% EC or dimethoate 30% EC at the rate of 1.0 liter / ha dissolved in 600-700 liters of water.
GREEN GRAM (Sowing)	Zaid green gram recommended castes- Narendra Moong-1, Samrat (PDM-139), Malaviya Janapriya (HUM-6), Meha, Malaviya Janakalyani, Malaviya Jan Chetna, Sweta, Swati, Virat, Shikha, IPM-2-3, IPM-2-14, Malaviya Jyoti etc. Make sowing work by arranging manure and seeds for sowing of any one variety. For sowing of Zaid green gram, sow at the rate of 20-25 kg seed / hectare.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
ONION (Vegetative)	Irrigate vegetable crops planted/sown during the summer season as needed. Thrips are likely to infest onion crops. To prevent this, spray Methyl-O-Dimethan 25% EC at the rate of 1.5 liters per hectare, diluted in 600-700 liters of water.
MANGO (Fruit formation)	Do irrigation work in the orchards of mango, guava, lemon plum, grapes, papaya and litchi etc. To prevent mango fruit drop, irrigate the orchard or apply to alpha naphthalene acetic acid on 4.5 SL of 20 ml per liter of water spray by making a solution . Spray 0.8 percent borax in 0.8 gm/liter of water twice at an interval of 10 days to prevent coalescence or internal rot disease in mango fruits.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	It is the appropriate time to give anthelmintic medicine to the animals for the prevention of worms in the stomach. FMD for the prevention of foot-mouth disease to animals vaccine and VQ for prevention of Lagadia fever. Get vaccinated with the vaccine. In view of the prevailing weather, farmers are advised tie the animals in a shaded place or in the shade of a tree during the day. Feed animals with green fodder and easily digestible mineral mixture and salt. The animals must be given clean and cold water 3-4 times a day. Do not leave the animals to graze from 11 am to 4 pm. bathe the animals in the morning and evening.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Farmers are advised to add dietary supplements; vitamins and energy feed ingredients to the hens as well as add calcium content to the hen. Give medicine for the prevention of worms in the stomach of hen (Divmirg). To protect the chickens from heat, arrange curtains, fans and ventilation in the hen house. To reduce the temperature of the hen house, cover the roof of the shed with straw.

Likely impacts of weather warnings (General)

According to the weather forecast received from the India Meteorological Department, there is a no weather warning in next five days.

Impact based advisories (General)

In view of the prevailing weather conditions, farmers are advised to undertake irrigation in standing crops such as okra, ridge gourd, pumpkin, bottle gourd, cucumber, maize, black gram, and green gram. Furthermore, the weather is favorable for the harvesting and threshing of mature crops like wheat, gram, peas, lentils, and pigeon pea.

Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: https://play.google.com/store/apps/details/

Meghdoot MobileApp link: https://play.google.com/store/apps/details

Damini MobileApp link : https://play.google.com/store/apps/details