



Gramin Krishi Mausam Sewa
Agromet Advisory Bulletin for HARDOI District
Chander Shekhar Azad University of Agriculture & Technology
 Kanpur, Uttar Pradesh



Agromet Advisory Bulletin

Date : 2026-09-18

Weather Forecast of District HARDOI (Uttar Pradesh) Issued On : 2026-09-18 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-09-19	2026-09-20	2026-09-21	2026-09-22	2026-09-23
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	33.0	34.0	35.0	35.0	36.0
Tmin(°C)	26.0	26.0	26.0	27.0	27.0
RH-I(%)	85	80	80	77	72
RH-II(%)	53	49	45	43	39
Wind Speed(kmph)	6	1	3	3	3
Wind Direction(Degree)	330	90	14	330	315
Cloud Cover(Octa)	5	2	1	0	0
Warning	No Warning	No Warning	No Warning	No Warning	No Warning

Forecast Summary:

According to the weather forecast received from the Indian Meteorological Department, light cloud cover in the first three days and clear sky weather in remaining days but there is no possibility rain. The maximum temperature is between 34.0- 38.0°C, which is likely to be 2-3°C higher normal and the minimum temperature is between 26.0-27.0°C, which is likely to be 2-3°C higher normal. The maximum and minimum range of relative humidity is between 72-85 and 39-53%. Wind direction is north-west, north-east and wind speed is between 1.0-6.0km/h with winds expected to be 1-2 Km/h normal.

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast of the India Meteorological Department, there is no warning in next five days

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Do the work of irrigation of standing Kharif crops as per requirement. Do the spraying of pesticides and insecticides, weeding and harvesting and threshing of mature Kharif crops. Do not allow water logging in maize, pulses and oilseed crops. Regularly check crops for insects and diseases after rains. Do not tie animals in open places or leave them outside for grazing and do not allow animals to go to waterlogged places. Keep the animals in a dry and safe place and give clean water to the animals. Do not tie the animals under trees or near electric poles, because the animals can die due to electric shock from falling branches or poles.

General Advisory:

Break/harvesting/threshing of mature crops standing in the field and sowing of rapeseed should be done and if the crop harvested in the field has got wet, then threshing should be done after drying it thoroughly in the sun. Prevent animals from going to flowing water/waterlogged areas. Do not allow animals to drink stagnant rain water. During the rainy season, the possibility of lice/lice infestation in animals becomes stronger. To prevent this, sprinkle lime in the animal enclosure so that the eggs and

larvae of the lice/lice are destroyed. During the rainy season, when going out of the house at night, carry a torch and to avoid snakes, scorpions etc., close the walls and holes of the house and sprinkle a little kerosene or phenyl, the smell of which drives away the snakes.

SMS Advisory:

Harvest/throw mature crops of maize and sow rapeseed in the absence of rain and monitor the diseases/pests in the crops standing in the field.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
RICE (Flowering)	The paddy crop is at the stage of ear budding (cabbage stage) and flowering stage, which are sensitive to lack of moisture, hence in case of no rain, maintain proper moisture by doing light irrigation as per requirement. If infestation of leaf roll, stem borer, green hopper and hispa insect is seen in paddy crop, then to control it, use 150 to 200 grams of Cartap Hydrochloride (50 SP) in 200 liters of water. Make a solution in water and spray per acre. Imidacloprid 17.8 percent SL for the prevention of stink bug and soldier bug. 125ml Or Azadirachtin 0.15 percent EC. Of 2.5 litres. Per quantity. Make a solution in 500-600 liters of water and spray in clear weather.
MAIZE (Physiological Maturity)	Harvest the ripe cobs of the maize crop when 75% of the cobs appear golden in color, dry them well in the sun and separate the grains from the cobs by hand or machine.
SESAME (GINGELLY/TIL) (Pod Development)	There is a possibility of showing an outbreak of leaf and pod larva pest in sesame crop, so for its prevention, use Oxydimethane-methyl 50% EC. Spray the solution in 500-600 liters of water at the rate of 1.0 liter/ha in clear weather.
GROUNDNUT (Pod Development)	At the time of pod formation in groundnut crops, maintain adequate moisture in the fields by doing light irrigation. There is a possibility of infestation of white giard/termite insect in groundnut crop, hence for its prevention use Fipronil 0.3% GR. Spread at the rate of 20 kg/hectare.
GREEN GRAM (Pod initiation)	There is a possibility of infestation of pod borer caterpillars in the urad/mung bean crop, hence to prevent this, spray Nibouli 5% or Quinalphos 25 EC at the rate of 1.25 liters/hectare in 500-600 liters of water.
RAPSEED (Sowing)	For sowing any one of the recommended varieties of rapeseed - Ta-36, Ta-9 Bhavani, PT-303, PT-30 etc., arrangement of fertilizer and seeds at the rate of 3-4 kg seeds/hectare do the sowing work .
SUGARCANE (Vegetative)	If the length of sugarcane has become five feet, then do the first tying at a height of two and half feet. For the first tying, tie two bunches of sugarcane in a line together with the dry leaves below. Destroy the plants affected by red rot disease by removing them along with the roots and spray the plants with a solution of thiophanate methyl 0.2 percent. If symptoms of bacterial rot disease are seen in sugarcane crop, cut and destroy the infected plant and for its control, spray 0.01% Streptomycin when weather is favorable.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
POTATO (Sowing)	Start sowing early potato varieties like Kufri Chandra Mukhi and Kufri Ashoka. Take out the potato seeds from cold storage 7-10 days before sowing. If sprouts emerge from seeds stored in cold storage, sort them out. If the potato seeds have not been treated before storing them in the cold storage, then immediately after removing the potatoes from the cold storage and sorting them, treat the potato tubers with 3 percent boric acid solution for 30 minutes and dry them in a shady place.
CABBAGE (Transplanting)	Sow radish, carrot, beetroot, turnip and leafy vegetables on ridges in the absence of rain. Transplant prepared seedlings of vegetables grown in Kharif like brinjal, chilli, tomato and early cauliflower etc. If there is a possibility of water stagnation in the field, then plant early cauliflower, brinjal, tomato, chilli etc. on the ridges. If the infestation of fruit borer/leaf borer insects is visible in vegetable crops, then to control it, make a solution of Neem oil in

Horticulture (Stage)	Horticulture Specific Advisory
	1.5 ml/litre of water and spray it 3-4 times at an interval of 8-10 days when the sky is clear.
GUAVA (Transplanting)	Try to complete the transplantation of fruit plants like mango, guava, amla, lemon etc. as soon as possible. Plant new plants in place of destroyed plants in new gardens. After hoeing the banana/papaya gardens add 25-30 cm high soil around the stem and make it stand. To prevent leaf cutting insect in mango, sprinkle 2% Carbaryl 1.5% dust when the sky is clear.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	Farmers are advised to take precautions as the possibility of lice/lice infestation in animals increases during the rainy season. To prevent this, sprinkle lime in the animal enclosure so that the eggs and larvae of the lice/lice are destroyed. Do not tie animals in open spaces/under trees during rains. Tie animals in the open only at night when the sky is clear. Tie the animals in a shady place during the day. Do not feed stagnant water from ponds or other places to animals, as this increases the risk of animals getting liver flu and stomach worms. This is the best time to give deworming medicine to prevent worms in the stomach of animals. Provide adequate amount of grains along with green and dry fodder to the animals. Make sure to feed clean and fresh water to animals 3-4 times a day. Do not tie pregnant animals to a sloping place. This is the right time to give deworming medicine to animals to prevent stomach worms.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Clean the chicken shelter every day. Provide clean water and balanced diet to the chickens. This time is not suitable for raising new chicks for eggs but this time is suitable for broiler chickens. Add dietary supplements, vitamins and energy food ingredients to the chickens' food as well as give the chickens calcium. Do chlorination of water. Give deworming medicine to kill stomach worms of chickens. Keep the hen house dry and keep turning the bedding. If there is excess moisture in the hen house, run a fan.

Likely impacts of weather warnings (General)

Waterlogging in low-lying areas disrupting daily activities and local commuting. Increased risk of road accidents due to reduced visibility and slippery roads. Potential for loss of life, injuries, and property damage caused by thunderstorms, lightning strikes and strong winds. Risk of casualties and damage to property from falling trees, branches, hoardings, and electric poles; temporary disruption to power, communication, and other essential services. Impact on residential areas, markets, schools, and public facilities due to localized flooding or waterlogging; increased risk of waterborne and mosquito-borne diseases caused by high humidity and waterlogging. Damage to crops, horticultural produce, livestock shelters, and agricultural infrastructure. Adverse effects on commercial activities, construction work, and the operations of outdoor workers.

Impact based advisories (General)

Regularly monitor the weather forecasts and warnings issued. During thunderstorms and lightning, avoid open fields, tall trees, electric poles, and water bodies. Travel only when necessary and avoid waterlogged roads, bridges, and culverts. Do not stand under weak structures, hoardings, or trees during strong winds. Inspect and secure rooftops, windows, and drainage systems around your home. Use electrical appliances with caution and keep flashlights and other essential emergency supplies ready in case of power outages. Take special care of children, elderly persons, pregnant women, and individuals with health conditions. Use clean and safe drinking water to prevent water-borne diseases. Keep livestock in safe shelters and ensure adequate fodder and clean drinking water for them.
--

Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: <https://play.google.com/store/apps/details/>

Meghdoot MobileApp link: <https://play.google.com/store/apps/details>

Damini MobileApp link : <https://play.google.com/store/apps/details>