Gramin Krishi Mausam Sewa

Agromet Advisory Bulletin for ETAWAH District

Chander Shekhar Azad University of Agriculture & Technology

Kanpur, Uttar Pradesh

Agromet Advisory Bulletin

Date : 2026-03-30

Weather Forecast of District ETAWAH (Uttar Pradesh) Issued On : 2026-03-30 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-03-31	2026-04-01	2026-04-02	2026-04-03	2026-04-04
Rainfall(mm)	6.0	7.0	0.0	0.0	5.0
Tmax(°C)	36.0	34.0	37.0	37.0	35.0
Tmin(°C)	21.0	22.0	21.0	22.0	21.0
RH-I(%)	71	70	68	63	77
RH-II(%)	41	42	42	39	46
Wind Speed(kmph)	7	14	5	3	7
Wind Direction(Degree)	144	321	34	315	297
Cloud Cover(Octa)	4	5	3	4	3
Warning	Thunderstorms & Lightning, Squall etc; Hailstorm; Strong Surface Winds	Thunderstorms & Lightning, Squall etc	No Warning	No Warning	No Warning

Forecast Summary:

According to the weather forecast from the India Meteorological Department, due to light to moderate cloud cover in the next five days, there is a possibility of light to moderate rain with local level at high winds speeds, thunderstorms & Lightning dated on 31 Mar, 01 & 04 Apr, 2026. The maximum temperature is between 34.0- 37.0°C, which is likely to be 2-3°C higher normal and the minimum temperature is between 21.0-22.0°C, which is likely to be 4-5°C higher normal. The maximum and minimum range of relative humidity is between 63-77 and 39-42%. The wind direction will be from the south-east, north-west, north-east and the wind speed will be between 3.0-14.0 km/hour, which is expected to be 3-4 km/hour higher than normal.

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the India Meteorological Department, there is a light rain, Thunderstorms & Lightning, Strong Surface Winds weather warning dated on 31March, 01 & 04 April, 2026.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

In view of the prevailing weather conditions, suspend the sowing and irrigation of crops such as okra, ridge gourd, pumpkin, bottle gourd, cucumber, snake cucumber, summer maize, black gram, and green

gram. Furthermore, harvest or dig up mature crops—such as mustard and potatoes—and store them in a safe location.

General Advisory:

Farmers are advised to make proper arrangements for the drainage of excess rainwater. They should gather harvested crops and cover them with polythene sheets; if harvested crops have become wet due to rain, they should be thoroughly dried in the sun before threshing, after which the grains should be stored safely. Irrigation activities for standing Zaid crops should be suspended. Do not spray insecticides, fungicides, or herbicides while standing against the direction of the wind. Spraying should ideally be carried out in the evening; furthermore—if possible—hands should be thoroughly washed with soap after spraying, before eating, and after washing clothes.

SMS Advisory:

Farmers are advised to thoroughly dry any harvested crops in the sun if they have become wet before threshing, and to postpone irrigation and sowing activities.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
WHEAT (Maturity and Harvesting)	In view of the possibility of rain, Farmers are advised to thoroughly dry harvested crops in keep safer place. In the wheat crop, when the outbreak of mice is seen, use fodder made of zinc phosphide or aluminum phosphide cakes collectively try to prevent mice.
CHICK PEA (Maturity and Harvesting)	To make proper arrangements for the drainage of excess water. In the chickpea crop, there is an outbreak of pod borer in the stage of grain filling. There is a possibility of outbreaks of pod borer insects so for it's prevent this, spraying Qunalphos 25 EC@ 2 liters per hectare or Emamectin benzoate 5 % sg 180 - 200 gram per hectare in 500 to 600 liters of water in the clear sky weather.
MAIZE (Vegetative)	In view of the possibility of rain, postpone the work of irrigation . In maize crop, do the first irrigation after 20-25 days of sowing and do weeding and hoeing after the oat arrives. For the prevention of stem borer pest in maize crop, apply phorate 10% g 10 kg or carbofuran 3% g 20 kg / ha after 10 -15 days after sowing.
BLACK GRAM (Sowing)	In view of the possibility of rain, postpone the work of sowing. Do to Zaid black gram is variety - Urd-Ta-9, Narendra Urd-1, Azad Urd-1, Azad Urd-2, Shekhar-2, Sujata, PU-40 etc arrange of fertilizer and seed for sowing of any one of the varieties. for sowing of black gram seeds at the rate of 25-30 kg of seed / hectare .
GREEN GRAM (Sowing)	In view of the possibility of rain, postpone the work of sowing. Recommended castes of Zaid Moong - Narendra Moong-1, Emperor (PDM-139), Malviya Janapriya (HUM-6), Meha, Malviya Janakalyani, Malviya Jan Chetna, Sweta, Swati, Virat, Shikha, IPM-2-3, IPM-2-14, Malviya Jyoti etc. For sowing of any one species, , do sowing work by arranging fertilizers and seeds. For sowing of Zaid moong crop, sow seed at the rate of 20-25 kg/ha.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
OKRA/ LADYFINGER (Sowing)	In view of the possibility of rain, postpone the work of irrigation / sowing. Outbreak of purple spot disease in onion crop affects the condition at 28 - 30 ° C temperature and 80 - 90% relative humidity, so copper oxychloride 0.3% (3 g/liter water) or Mancozeb at the rate of 0.25% (0.25 g/liter of water) and

Horticulture (Stage)	Horticulture Specific Advisory
	spray 3-4 at a 15-20 day interval in the clear sky weather . Sown vegetables such as ladyfinger, zucchini, cucumber, bitter gourd, gourd and pumpkin etc. sown during summer season. Transplant the seedlings of tomato, brinjal and chilli.
MANGO (Flowering)	Irrigate the orchards or spray with acetic acid solution of 15 ppm or 4 ml Planofix per 9 liters of water to prevent drop of mango fruits. There is a possibility of showing the outbreak of midge pest in mango blossom, so for its prevention, spray Fenitrothian in 1.0 ml / liter of water and spray it.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	It is the appropriate time to give anthelmintic medicine to the animals for the prevention of worms in the stomach. Make OT mixture of 750 grams mustard oil, 100 grams turmeric, 50 grams dry ginger, 50 grams cumin, 500 grams jaggery after feeding the animals for three days in the morning and evening. In view of the prevailing weather, farmers are advised not to tie the animals in the open during the night. Tie the animals in a shaded place or in the shade of a tree during the day. Give sufficient amount of grains along with green and dry fodder to the animals. The animals must be given clean and fresh water 3-4 times a day.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Farmers are advised to add dietary supplements; vitamins and energy feed ingredients to the hens as well as add calcium content to the chickens. Give medicine for the prevention of worms in the stomach of chickens (Divmirg).

Likely impacts of weather warnings (General)

According to the weather forecast received from the India Meteorological Department, there is a light rain, Thunderstorms & Lightning, ; Strong Surface Winds weather warning.

Impact based advisories (General)

In view of the prevailing weather conditions, suspend the sowing and irrigation of crops such as okra, ridge gourd, pumpkin, bottle gourd, cucumber, snake cucumber, summer maize, black gram, and green gram. Furthermore, harvest or dig up mature crops—such as mustard and potatoes—and store them in a safe location.

Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: https://play.google.com/store/apps/details/

Meghdoot MobileApp link: https://play.google.com/store/apps/details

Damini MobileApp link : https://play.google.com/store/apps/details