



Agromet Advisory Bulletin

Date : 2026-08-11

Weather Forecast of District HARDOI (Uttar Pradesh) Issued On : 2026-08-11 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-08-12	2026-08-13	2026-08-14	2026-08-15	2026-08-16
Rainfall(mm)	0.0	0.0	8.0	12.0	8.0
Tmax(°C)	33.0	36.0	35.0	31.0	32.0
Tmin(°C)	26.0	27.0	28.0	27.0	27.0
RH-I(%)	94	85	88	95	94
RH-II(%)	60	55	61	76	70
Wind Speed(kmph)	13	3	6	5	4
Wind Direction(Degree)	81	21	245	193	285
Cloud Cover(Octa)	4	4	7	8	7
Warning	NA	NA	NA	NA	NA

Forecast Summary:

According to the weather forecast received from the India Meteorological Department, due to light to moderate cloud cover in the next five days but there is a possibility of light to moderate rain third, fourth & fifth days with thunderstorms & lightning, squall, strong winds and remaining day is clear sky weather due to no rain. The maximum temperature is between 31.0- 36.0°C, which is likely to be 1-2°C higher normal and the minimum temperature is between 26.0-28.0°C, which is likely to be 1-2°C higher normal. The maximum and minimum range of relative humidity is between 85-94 and 55-76%. Wind direction is north-east, south-west, north-west and wind speed is between 3.0-13.0 kmph with gusts expected to be 4-5 kmph higher than normal speeds.

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the India Meteorological Department (IMD), there is a third, fourth, fifth days warning of thunderstorms, strong winds light to moderate rain.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Carry out sowing and other agricultural activities when there is no rain and the sky is clear. Apply fertilizers, insecticides, and fungicides only when the weather clears up. Regularly monitor crops for pests and diseases after the rain. Keep livestock in a dry, safe area and provide them with clean drinking water. Do not tether animals under trees or near electricity poles, as falling branches or electric shocks from the poles could prove fatal to the animals.

General Advisory:

Postpone activities such as sowing crops (like Urad and Moong), applying pesticides, fungicides, or fertilizers, and irrigating fields. Strengthen the embankments (bunds) of paddy fields to retain rainwater within the fields. Regularly monitor crops for pests and diseases after the rain. Do not allow livestock to drink stagnant rainwater; keep animals in a dry, safe area and provide them with clean drinking water. Avoid tethering animals under trees or near electric poles. During the rainy season, the risk of tick and louse infestations in livestock increases significantly; to prevent this, sprinkle lime in the animal enclosure to destroy their eggs and larvae. When stepping outside at night during the rainy season, carry a torch; to protect against snakes and scorpions, seal cracks and holes in house walls and spray a small amount of kerosene or phenyl, as the odor repels snakes.

SMS Advisory:

Never take shelter under trees, telephone and electricity poles during rain. In such a situation, there is a high possibility of electric current flowing in the poles and lightning striking.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
RICE (Transplanting)	For controlling broad and narrow leaf weeds in paddy crop, use Bispyribac Sodium 10% SC at the rate of 0.20 litre per hectare, 15-20 days after transplanting, by dissolving it in 500-600 litres of water at appropriate moisture and spray when sky is clear. If Khaira disease is seen in paddy, then spray 20-25 kg zinc sulphate and 2.5 kg lime solution in 800 liters of water for its control. Topdressing of urea @ 50 to 60 kg per hectare is done 25 to 30

Crop (Stage)	Crop Specific Advisory
	days after transplanting of paddy. Keep in mind that weeds should be removed before top dressing and while topdressing, there should not be more than 2 to 3 cm of water in the field. If infestation of root worm pest is seen in paddy crop, spray Cartap hydrochloride 4% granular chemical at the rate of 20 to 25 kg per hectare in three to four cm stagnant water.
MAIZE (Vegetative)	The second topdressing of urea in maize crop should be done at the rate of 60 to 70 kg per hectare at the time of tasseling stage in case of no rain. In maize crop, there is a possibility of outbreak of stem borer / shoot fly pest, so for its prevention, Emamectin Benzoate 5% Sg 200 g / ha at 500-600 liters of water solution Sprinkle on clean weather.
SESAME (GINGELLY/TIL) (Vegetative)	In sesame crop, do the second weeding 35-40 days after sowing in case of no rain. There is a possibility of outbreak of leaf and fruit bollworm in sesame crop, so for its prevention, Emamectin Benzoate 5 % SG 200 g / hectare or Quinalphos 25 % EC 1. 500-600 liters at the rate of 25 liters / hectare. Make a solution in water and spray on clear weather.
GROUNDNUT (Vegetative)	In case of no rain, after weeding the peanut crops 20 to 25 days after sowing, add 100 kg gypsum per hectare and then do light hoeing and cover the plants with soil. Groundnut crop is likely to show white gidar / termite pest infestation, hence fipronil 0.3% GR at the rate of 20 kg / ha topdressing of clear sky.
BLACK GRAM (Vegetative)	For the control of weeds in the black gram crop, Imazethapyr 10 SL at the rate of 1.0 liter / hectare, spray the solution in 500-600 liters of water after 15-20 days of sowing and spray it on clear sky.
GREEN GRAM (Sowing)	For the control of weeds in the green gram crop, Imazethapyr 10 SL at the rate of 1.0 liter / hectare, spray the solution in 500-800 liters of water after 15-20 days of sowing and spray it on clear sky. The recommended varieties of green gram sown in Kharif - Narendra Moong-1, Pant Moong-1, Pant Moong-3, Pant Moong-4, Samrat, Malaviya Janchetna, Malaviya Janapriya, Malaviya Jagriti Shweta, Swati, Virat and IPM -2-14 etc For the sowing of any one variety. The sowing work at the rate of 12 -15 kg seed / hectare and try to complete it soon the event of no rain.
SUGARCANE (Grand growth)	If the length of sugarcane has become five feet, then do the first tying at a height of two and half feet. For the first tying, tie two bunches of sugarcane in a line together with the dry leaves below. Destroy the plants affected by red rot disease by removing them along with the roots and spray the plants with a solution of thiophanate methyl 0.2 percent. Infestation of mite insect is being seen in sugarcane, for its control, mix 750 ml of Profenophos 40 percent + Cyper 4 percent insecticide in 625 liters of water and spray at the rate of per hectare.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
CAULIFLOWER (Transplanting)	Suspend irrigation/sowing work in standing vegetable crop in view of the possibility of rain. Set up nurseries of vegetables to be sown in Kharif like cabbage, knotweed, broccoli etc., along with planting ready saplings of early cauliflower, brinjal, chilli. If there is a possibility of water stagnation in the field, then plant the early cauliflower, brinjal, tomato, chilli etc. on the ridges. For prevention of fruit borer/leaf borer pest in vegetable crops, prepare a solution of 1.5-2.0 ml/liter of water with neem oil and spray it 3-4 times at an interval of 8-10 days. Weeding-hoeing should be done in standing vegetable crops when sky is clear.
MANGO (Planting)	Plant new plants in pits already filled with mango, guava, amla, lemon etc. Plant new plants in place of the destroyed plants in the new bag. Make a stand of 25-30 meters high soil around the stem by folding banana / papaya bags. For prevention of mango leaf moth, apply 2% carburil 1.5% dust on the clear sky.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	Farmers are advised to During the rainy season, the possibility of lice infestation in animals increases. To prevent this, spray lime in the animal shed so that the eggs and larvae of lice are destroyed. Do not tie animals in open spaces/under trees during rains. Tie animals in the open only when the sky is clear at night. Tie animals in a shady place during the day. Vaccinate animals with FMD vaccine to prevent foot and mouth disease and VQ vaccine to prevent dengue fever. Give sufficient amount of grains to animals along with green and dry fodder. Make sure to give clean and fresh water to animals 3-4 times a day. Do not tie pregnant animals at a sloping place. This is the right time to give deworming medicine to animals to prevent stomach worms. The department is operating a Mobile Veterinary Unit scheme to provide veterinary services at the doorsteps of farmers and livestock owners. Farmers and livestock owners can contact the toll-free helpline number 1962 to avail the benefits of this scheme.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Protect the chickens from dampness and waterlogging during the current weather conditions. Ensure proper ventilation in the poultry shed and provide clean drinking water along with a balanced diet. Maintain cleanliness around the shed to control mosquitoes and flies, and ensure timely vaccination for all

Poultry	Poultry Specific Advisory
	chickens. Immediately isolate any chickens that appear sick or lethargic, and seek veterinary advice promptly if necessary.

Likely impacts of weather warnings (General)

Waterlogging in low-lying areas disrupting daily activities and local commuting. Increased risk of road accidents due to reduced visibility and slippery roads. Potential for loss of life, injuries, and property damage caused by thunderstorms, lightning strikes and strong winds. Risk of casualties and damage to property from falling trees, branches, hoardings, and electric poles; temporary disruption to power, communication, and other essential services. Impact on residential areas, markets, schools, and public facilities due to localized flooding or waterlogging; increased risk of waterborne and mosquito-borne diseases caused by high humidity and waterlogging. Damage to crops, horticultural produce, livestock shelters, and agricultural infrastructure. Adverse effects on commercial activities, construction work, and the operations of outdoor workers.

Impact based advisories (General)

Regularly monitor the weather forecasts and warnings issued. During thunderstorms and lightning, avoid open fields, tall trees, electric poles, and water bodies. Travel only when necessary and avoid waterlogged roads, bridges, and culverts. Do not stand under weak structures, hoardings, or trees during strong winds. Inspect and secure rooftops, windows, and drainage systems around your home. Use electrical appliances with caution and keep flashlights and other essential emergency supplies ready in case of power outages. Take special care of children, elderly persons, pregnant women, and individuals with health conditions. Use clean and safe drinking water to prevent water-borne diseases. Keep livestock in safe shelters and ensure adequate fodder and clean drinking water for them.

Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: <https://play.google.com/store/apps/details/>

Meghdoot MobileApp link: <https://play.google.com/store/apps/details>

Damini MobileApp link : <https://play.google.com/store/apps/details>