



Gramin Krishi Mausam Sewa
Agromet Advisory Bulletin for UNNAO District
Chander Shekhar Azad University of Agriculture & Technology
Kanpur, Uttar Pradesh



Agromet Advisory Bulletin

Date : 25-09-2026

Weather Forecast of UNNAO(Uttar Pradesh) Issued On : 2026-09-25(Valid Till 08:30 IST of the next 5 days)

Parameter	2026-09-26	2026-09-27	2026-09-28	2026-09-29	2026-09-30
Rainfall(mm)	82.0	61.0	0.0	0.0	0.0
Tmax(°C)	25.0	25.0	26.0	30.0	32.0
Tmin(°C)	22.0	22.0	22.0	23.0	24.0
RH-I(%)	95	96	97	94	90
RH-II(%)	90	91	80	67	56
Wind Speed(kmph)	21	23	12	6	8
Wind Direction(Degree)	68	47	169	196	260
Cloud cover(octa)	8	8	5	2	1
Warning	Heavy Rain; Thunderstorms & Lightning, Squall etc; Strong Surface Winds	Heavy Rain; Thunderstorms & Lightning, Squall etc; Strong Surface Winds	No Warning	No Warning	No Warning

Forecast Summary:

According to the weather forecast received from the Indian Meteorological Department, due to moderate to dense cloud cover in the next five day there is a possibility of moderate to heavy rain with thunderstorms & lightning at local level between dated on 26-27 September, 2026. The maximum temperature is between 25.0-32.0°C, which is likely to be 4-5°C lower normal and the minimum temperature is between 22.0-23.0°C, which is likely to be 1-2°C lower normal. The maximum and minimum range of relative humidity is between 90-97 and 56-91%. Wind direction is north-east, south-east, south-west and wind speed is between 6.0-23.0 km/h with winds expected to be 10-12 Km/h higher normal.

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast of the India Meteorological Department, a warning of moderate to heavy rain with strong winds, thunderstorm and lightning.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Farmers are advised to postpone irrigation and sowing activities in view of the likelihood of rain. Mature crops in the fields should be harvested and threshed, and the grain stored in a safe place. Harvested crops should be gathered on elevated ground within the field and covered with polythene sheets. After the rain, the crop should be thoroughly dried in the sun before threshing. Do not tether livestock in open areas or let them out to graze, and prevent them from entering waterlogged areas. Keep the animals in a dry and safe place and give clean water to the animals. Do not tie the animals under trees or near electric poles, because the animals can die due to electric shock from falling branches or poles.

General Advisory:

Farmers are advised to postpone irrigation and sowing operations in view of the likelihood of rain, and to complete the harvesting and threshing of mature crops as soon as possible. If the crop harvested in the field has got wet, then threshing should be done after drying it thoroughly in the sun. Prevent animals from going to flowing water/waterlogged areas. Do not allow animals to drink stagnant rain water. During the rainy season, the possibility of lice/lice infestation in animals becomes stronger. To prevent this, sprinkle lime in the animal enclosure so that the eggs and larvae of the lice/lice are destroyed. During the rainy season, when going out of the house at night, carry a torch and to avoid snakes, scorpions etc., close the walls and holes of the house and sprinkle a little kerosene or phenyl, the smell of which drives away the snakes.

SMS Advisory:

In view of the likelihood of rain, postpone the harvesting, threshing, and sowing of mature crops; additionally, make proper arrangements for draining excess rainwater.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
RICE (Millking)	Farmers are advised to postpone irrigation operations in view of the potential for rain. The rice crop is in the stage of emergence of ears and flowering/milking, which is sensitive to moisture deficiency, so in the absence of rain, maintain proper moisture by doing light irrigation as per requirement. The current weather is favorable for pests, so on an average, 2 - 3 odorous pests / hill are visible in the paddy crop after flowering, then for its control, Imidacoloroprid 17. 8% SL 350 ml / ha or Burfozin 25% SP 750 ml Make a solution in 500 -600 liters of water at the rate of / hectare and spray it on clear weather. If the outbreak of stem borer insect is seen in the paddy crop, then Imidacloprid 17.8 EC for its control. SL at the rate of 25 g/ha make a solution in 500-600 liters of water and spray it in clear weather.
MAIZE (Maturity and Harvesting)	In view of the potential for rain, complete harvesting and threshing of mature crops as soon as possible. Harvest the ripe cobs of the maize crop when 75% of the cobs appear golden in color, dry them well in the sun and separate the grains from the cobs by hand or machine.
SESAME (GINGELLY/TIL) (Physiological Maturity)	In view of the potential for rain, complete harvesting and threshing of mature crops as soon as possible. There is a possibility of showing an outbreak of leaf and pod larva pest in sesame crop, so for its prevention, Emamectin Benzoate 5 % SG 200 g / hectare or Quinalphos 25 % EC at the rate of 1. 25 liters / hectare make a solution in 500-600 liters water and spray on clear weather.
GROUNDNUT (Pod Development)	Farmers are advised to postpone irrigation activities in view of the likelihood of rain and to make proper arrangements for the drainage of rainwater. Groundnut crop is likely to show white gidar / termite pest infestation, hence fipronil 0.3% GR at the rate of 20 kg / ha topdressing of clear sky.
BLACK GRAM (Pod Development)	Farmers are advised to postpone irrigation activities in view of the likelihood of rain and to make proper arrangements for the drainage of rainwater. To prevent the pod borer larvae in standing black gram/green gram crops, which bore into the pods and eat the grains, spray 750 ml of Cypermethrin 10 EC per hectare by dissolving it in 500-600 liters of water in clear weather.
RAPESEED (Sowing)	Farmers are advised to postpone sowing operations in view of the potential for rain. For sowing any one of the recommended varieties of rapeseed - Ta-36, Ta-9 Bhavani, PT-303, PT-30 etc., arrangement of fertilizer and seeds at the rate of 3-4 kg seeds/hectare do the sowing work .
SUGARCANE (Vegetative)	Farmers are advised to postpone irrigation operations in view of the potential for rain. If the length of sugarcane has become five feet, then do the first tying at a height of two and half feet. For the first tying, tie two bunches of sugarcane in a line together with the dry leaves below. Destroy the plants affected by red rot disease by removing them along with the roots and spray the plants with a solution of thiophanate methyl 0.2 percent. If symptoms of bacterial rot disease are seen in sugarcane crop, cut and destroy the infected plant and for its control, spray 0.01% Streptomycin when weather is favorable.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
POTATO (Sowing)	Farmers are advised to postpone sowing operations in view of the potential for rain. Start sowing of early varieties of potato like Kufri Chandra Mukhi and Kufri Ashoka. Take out the potato seeds from cold storage 7-10 days before sowing. If sprouts appear in the seeds stored in cold storage, then sort them and separate them. If the potato seeds have not been treated before storing them in the cold storage, then immediately after taking out the potatoes from the cold storage and sorting them, treat the potato tubers with 3 percent boric acid solution for 30 minutes and dry them in a shady place.
CABBAGE (Transplanting)	Farmers are advised to postpone irrigation and sowing operations in view of the potential for rain. Sow radish, carrot, beetroot, turnip and leafy vegetables on the ridges. Transplanting of saplings planted brinjal, chilli, tomato and early cauliflower. If there is a possibility of water stagnation in the field, then plant the early cauliflower, brinjal, tomato, chilli etc. on the ridges. If there is an outbreak of fruit borer / leaf borer in vegetable crops, make a solution of neem oil in 1.5 ml / liter of water to control it and spray it 3-4 times after clearing the sky at intervals of 6-10 days.
GUAVA (Transplanting)	Try to complete the planting of fruit plants like mango, guava, amla, lemon etc. as soon as possible. Plant new plants in place of the destroyed plants in the new bag. Make a stand of 25-30 meters high soil around the stem by folding banana / papaya bags. For prevention of mango leaf moth, apply 2% carbaryl 1.5% dust on the clear sky.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	Farmers are advised to take precautions as the possibility of lice/lice infestation in animals increases during the rainy season. To prevent this, sprinkle lime in the animal enclosure so that the eggs and larvae of the lice/lice are destroyed. Do not tie animals in open spaces/under trees during rains. Tie animals in the open only at night when the sky is clear. Tie the animals in a shady place during the day. Do not feed stagnant water from ponds or other places to animals, as this increases the risk of animals getting liver flu and stomach worms. This is the best time to give deworming medicine to prevent worms in the stomach of animals. Provide adequate amount of grains along with green and dry fodder to the animals. Make sure to feed clean and fresh water to animals 3-4 times a day. Do not tie pregnant animals to a sloping place. This is the right time to give deworming medicine to animals to prevent stomach worms.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Clean the chicken shelter every day. Provide clean water and balanced diet to the chickens. This time is not suitable for raising new chicks for eggs but this time is suitable for broiler chickens. Add dietary supplements, vitamins and energy food ingredients to the chickens' food as well as give the chickens calcium. Do chlorination of water. Give deworming medicine to kill stomach worms of chickens. Keep the hen house dry and keep turning the bedding. If there is excess moisture in the hen house, run a fan.

Likely impacts of weather warnings (General)

Waterlogging in low-lying areas disrupting daily activities and local commuting. Increased risk of road accidents due to reduced visibility and slippery roads. Potential for loss of life, injuries, and property damage caused by thunderstorms, lightning strikes and strong winds. Risk of casualties and damage to

property from falling trees, branches, hoardings, and electric poles; temporary disruption to power, communication, and other essential services. Impact on residential areas, markets, schools, and public facilities due to localized flooding or waterlogging; increased risk of waterborne and mosquito-borne diseases caused by high humidity and waterlogging. Damage to crops, horticultural produce, livestock shelters, and agricultural infrastructure. Adverse effects on commercial activities, construction work, and the operations of outdoor workers.

Impact based advisories (General)

Regularly monitor the weather forecasts and warnings issued. During thunderstorms and lightning, avoid open fields, tall trees, electric poles, and water bodies. Travel only when necessary and avoid waterlogged roads, bridges, and culverts. Do not stand under weak structures, hoardings, or trees during strong winds. Inspect and secure rooftops, windows, and drainage systems around your home. Use electrical appliances with caution and keep flashlights and other essential emergency supplies ready in case of power outages. Take special care of children, elderly persons, pregnant women, and individuals with health conditions. Use clean and safe drinking water to prevent water-borne diseases. Keep livestock in safe shelters and ensure adequate fodder and clean drinking water for them.

Farmers are advised to download Unified  Mausam  and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: <https://play.google.com/store/apps/details/>

Meghdoot MobileApp link: <https://play.google.com/store/apps/details>

Damini MobileApp link : <https://play.google.com/store/apps/details>