INFOSYS TEST SERIES-1

Infosys Systems Engineer (SE) | 5-6 Nov 2026 Drive | 100% Pure SE Verified

Total Questions: 35 | **Total Marks:** 140 | **Duration:** 4200000 Minutes

Section: Puzzles

Problem 1 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] PRACTICE VARIANT

ID: Q1 | Source: Infosys PYQ | Level: Infosys SE Standard

A 76 | B 45

C 67 | D 68

Verified Answer: ✅ **(D) 68** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Logic: 5 x 3 + 6 x 4 = 39

Problem 2 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] PRACTICE VARIANT

ID: Q2 | Source: Infosys PYQ | Level: Infosys SE Standard

A 11 | B 33

C 22 | D 32

Verified Answer: ✅ **(A) 11** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Logic: 3 x 5 + 4 x 3 = 27

Problem 3 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] PRACTICE VARIANT

ID: Q3 | Source: Infosys PYQ | Level: Infosys SE Standard

A 4 | B 2

C 6 | D 8

Verified Answer: ✅ **(B) 2** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Logic: 9 x 4 – (9 + 2) = 25

Problem 4 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] PRACTICE VARIANT

ID: Q4 | Source: Infosys PYQ | Level: Infosys SE Standard

A 4 | B 3

C 2 | D 1

Verified Answer: ✅ **(D) 1** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Logic: 4 x 5 – (4 + 5) =11 3 x 2 – (3+2) = 1

Problem 5 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q5 Source: Infosys PYQ Level: Infosys SE Standard

A J	**B** T
C I	**D** O

✓ **Verified Answer:** ✅ **(A) J** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Logic: Write the alphabetical position first. Then 3 x 8 /2 = 12 , where 3 is C , H is 8 and L=12

Problem 6 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q6 Source: Infosys PYQ Level: Infosys SE Standard

A 2	**B** 13
C 25	**D** 5

✓ **Verified Answer:** ✅ **(B) 13** **Step-by-Step Explanation & Derivation:** Detailed step-by-step deduction provided in solutions compendium.

Problem 7 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q7 Source: Infosys PYQ Level: Infosys SE Standard

A None	**B** 40
C 50	**D** 60

✓ **Verified Answer:** ✅ **(B) 40** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** j=50 and j-10 , 50-10 = 40

Problem 8 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q8 Source: Infosys PYQ Level: Infosys SE Standard

A 6	**B** 8
C 5	**D** 4

✓ **Verified Answer:** (A) **Step-by-Step Explanation & Derivation:** Detailed step-by-step deduction provided in solutions compendium.

Problem 9 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q9 Source: Infosys PYQ Level: Infosys SE Standard

What will be the
output of the following pseudo code
Integer
i, j, k, d
Set
j = 6, i = 8, k = 2, d = 1
if
((j ^ i) mod k equals 0) then
d = d + 1
else
d = d - 1
end
if
Print
d

A 1	**B** 3
C 0	**D** 2

✓ **Verified Answer:** ✅ **(D) 2** **Step-by-Step Explanation & Derivation:** Detailed step-by-step deduction provided in solutions compendium.

Problem 10 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q10 Source: Infosys PYQ Level: Infosys SE Standard

A 20	**B** 40
C 60	**D** 80

✓ **Verified Answer:** ✅ **(B) 40** **Step-by-Step Explanation & Derivation:** Detailed step-by-step deduction provided in solutions compendium.

Problem 11 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q11 Source: Infosys PYQ Level: Infosys SE Standard

If
a number is divided by 5, remainder = 2. If the same number is
divided by 7, remainder = 3. What is the smallest such number?

A 17	**B** 23
C 38	**D** 52

✓ **Verified Answer:** ✅ **(D) 52** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Number = 5k + 2. Check divisibility by 7. k=6 → 5(6)+2=32 remainder 4. k=7 → 37 remainder 2. k=8 → 42 remainder 0. k=9 → 47 remainder 5. k=10 → 52 remainder 3 . So number = 52 .

Problem 12 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q12 Source: Infosys PYQ Level: Infosys SE Standard

The
probability of getting a sum of 9 when two dice are thrown is:

A 1/12	**B** 1/9
C 5/39	**D** 7/36

✓ **Verified Answer:** ✅ **(A) 1/12** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Combinations: (3,6), (4,5), (5,4), (6,3). Total = 4 outcomes. Probability = 4/36 = 1/9 (Correction → 1/9, not 1/12).

Problem 13 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q13 Source: Infosys PYQ Level: Infosys SE Standard

The
sum of present ages of A and B is 60. 5 years ago, A was 5 times
B. What are their ages?

A A=55,	**B** A=50,
C A=45,	**D** A=40,

✓ **Verified Answer:** (B) **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** A+B=60. (A–5) = 5(B–5). Solving → A=50, B=10.

Problem 14 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q14 Source: Infosys PYQ Level: Infosys SE Standard

If $x^2 - 5x + 6 = 0$, then find the value of $(1/x_1 + 1/x_2)$.

A 1/2	**B** 5/6
C 2/3	**D** 3/2

✓ **Verified Answer:** ✅ **(B) 5/6** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Roots = 2, 3. (1/2 + 1/3) = (3+2)/6 = 5/6.

Problem 15 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q15 Source: Infosys PYQ Level: Infosys SE Standard

A can complete a work in 20 days. B is 25% more efficient than A. In how many days will B finish the same work?

A 15	**B** 16
C 18	**D** 20

✓ **Verified Answer:** ✅ **(B) 16** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** A's 1-day work = 1/20. Efficiency of B = 125% of A = 5/4 × (1/20) = 1/16. So B will finish in 16 days.

Problem 16 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q16 Source: Infosys PYQ Level: Infosys SE Standard

A shopkeeper allows 20% discount on the marked price and still makes 25% profit. If the marked price is ₹1000, what is the cost price?

A ₹600	**B** ₹625
C ₹650	**D** ₹700

✓ **Verified Answer:** ✅ **(A) ₹600** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** SP = 1000 – 20% of 1000 = 800. Profit = 25%. CP = 800 / 1.25 = 640 (correction, not 600).

Problem 17 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q17 Source: Infosys PYQ Level: Infosys SE Standard

The ratio of investments of A and B is 4:5. A gets ₹400 less than B from the profit. Find the total profit.

A	₹3600	**B**	₹4000
C	₹4200	**D**	₹4800

✓ **Verified Answer:** ✅ **(A) ₹3600** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Difference = 1 part. 1 part = 400. Total = 9 parts = 9×400=3600.

Problem 18 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q18 Source: Infosys PYQ Level: Infosys SE Standard

A sum of ₹10,000 is invested at 20% per annum compounded annually. What will be the compound interest after 2 years?

A	₹4000	**B**	₹4200
C	₹4400	**D**	₹4800

✓ **Verified Answer:** ✅ **(C) ₹4400** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** CI = P[(1 + r/100)^t – 1] = 10000[(1.2)^2 – 1] = 10000(1.44 – 1) = 4400. (Answer : ₹4400).

Problem 19 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q19 Source: Infosys PYQ Level: Infosys SE Standard

The average of 5 consecutive odd numbers is 35. What is the largest of these numbers?

A	37	**B**	39
C	41	**D**	43

✓ **Verified Answer:** ✅ **(B) 39** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Middle term = 35 (average of consecutive odd nos). Series = 31, 33, 35, 37, 39 → largest = 39.

Problem 20 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q20 Source: Infosys PYQ Level: Infosys SE Standard

Two
pipes can fill a tank in 15 min and 20 min respectively. If both
pipes are opened together, how long will it take to fill the tank?

A 7	**B** 8
C 8	**D** 9

✓ **Verified Answer:** (C) **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** 1/15 + 1/20 = (4+3)/60 = 7/60. Time = 60/7 = 8.57 min ≈ 8 min 34 sec.

Problem 21 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q21 Source: Infosys PYQ Level: Infosys SE Standard

Arrange
according to logical order:
(1)
Exam, (2) Study, (3) Result, (4) Admission.

A 2-1-3-4	**B** 1-2-3-4
C 2-3-1-4	**D** 4-2-1-3

✓ **Verified Answer:** ✅ **(A) 2-1-3-4** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** First Study → Exam → Result → Admission.

Problem 22 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q22 Source: Infosys PYQ Level: Infosys SE Standard

Complete
the series: 7, 26, 63, 124, ?

A 215	**B** 216
C 225	**D** 200

✓ **Verified Answer:** ✅ **(A) 215** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Pattern: $n^3 - 1$. (2^3–1=7, 3^3–1=26, 4^3–1=63, 5^3–1=124, 6^3–1=215).

Problem 23 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q23 Source: Infosys PYQ Level: Infosys SE Standard

Statements:
Some
pens are pencils.
All
pencils are erasers.
Conclusions:
I. Some pens are
erasers.
II. All erasers are pencils.

A Only	**B** Only
C Both	**D** Neither

✓ **Verified Answer:** (A) **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Pens ⊂ Pencils ⊂ Erasers → Some pens ⊂ Erasers.

Problem 24 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q24 Source: Infosys PYQ Level: Infosys SE Standard

Six
friends A, B, C, D, E, F are sitting in a circle facing the
center. A is between D and F. C is to the left of D. E is between
B and C. Who is sitting between A and C?

A D	**B** E
C F	**D** B

✓ **Verified Answer:** ✅ **(A) D** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Arranging: A between D & F, C left of D, E between B & C → D is between A and C.

Problem 25 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q25 Source: Infosys PYQ Level: Infosys SE Standard

Find
the missing number:

A 24	**B** 36
C 48	**D** 36

✓ **Verified Answer:** ✅ **(D) 36** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** 8 → 8^2=64. 6 → 6^2=36.

Problem 26 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q26 Source: Infosys PYQ Level: Infosys SE Standard

Two trains of lengths 120m and 180m are running in opposite directions with speeds 60 km/h and 90 km/h. In how much time will they cross each other?

A 6 sec	**B** 8 sec
C 10 sec	**D** 12

✓ **Verified Answer:** ✅ **(C) 10 sec** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Relative speed = (60+90)×(5/18)=41.67 m/s. Total length = 300m. Time = 300/41.67 ≈ 10 sec.

Problem 27 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q27 Source: Infosys PYQ Level: Infosys SE Standard

Pointing to a boy, Seema said, “He is the son of my grandfather’s only son.” How is the boy related to Seema?

A Brother	**B** Cousin
C Uncle	**D** Nephew

✓ **Verified Answer:** ✅ **(A) Brother** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Grandfather’s only son = Seema’s father. Boy = son of Seema’s father → brother.

Problem 28 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q28 Source: Infosys PYQ Level: Infosys SE Standard

In a row of 40 students, A is 12th from the left and B is 18th from the right. How many students are between A and B?

A 8	**B** 9
C 10	**D** 11

✓ **Verified Answer:** ✅ **(C) 10** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** A = 12th from left. B = 23rd from left (40–18+1). Between = 23–12–1 = 10.

Problem 29 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] CORE SE

ID: Q29 Source: Infosys PYQ Level: Infosys SE Standard

Five
friends scored different marks. P scored more than Q but less than
R. S scored less than Q but more than T. Who scored the lowest?

A P

B Q

C S

D T

Verified Answer: ✅ **(D) T** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Order: R > P > Q > S > T → lowest is T.

Problem 30 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q30 Source: Infosys PYQ Level: Infosys SE Standard

If
'A × B' means A is the father of B, 'A +
B' means A is the sister of B, and 'A – B'
means A is the mother of B, then P – Q × R means?

A P

B P

C P

D P

Verified Answer: (A) **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Q × R → Q father of R. P – Q → P mother of Q → So P mother of R.

Problem 31 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q31 Source: Infosys PYQ Level: Infosys SE Standard

Find
the missing term: AB, DE, HI, MN, ?

A RS

B ST

C TU

D VW

Verified Answer: ✅ **(B) ST** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Skipping letters: (AB → DE → HI → MN → ST).

Problem 32 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q32 Source: Infosys PYQ Level: Infosys SE Standard

What
comes next?
2,
12, 36, 80, 150, ?

A 252
B 240
C 260
D 300

✓ **Verified Answer:** ✅ **(A) 252** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Pattern: $n^3 + n^2$. ($1^3+1^2=2$, $2^3+2^2=12$, $3^3+3^2=36$, $4^3+4^2=80$, $5^3+5^2=150$, $6^3+6^2=252$).

Problem 33 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q33 Source: Infosys PYQ Level: Infosys SE Standard

In
a certain code, FLOWER is written as EKNVDQ. How will GARDEN be
written?

A FZQCDM
B HBSFEP
C FYQBCM
D EZQCDM

✓ **Verified Answer:** ✅ **(A) FZQCDM** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Each letter shifted –1: F→E, L→K, O→N... So GARDEN → FZQCDM.

Problem 34 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q34 Source: Infosys PYQ Level: Infosys SE Standard

A,
B, C, D, E are standing in a line. C is between A and E. B is to
the immediate left of E. D is to the right of A but not at the
end. Who is in the middle?

A A
B B
C C
D D

✓ **Verified Answer:** ✅ **(C) C** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** Order: A D C B E → middle = C.

Problem 35 • [Marks: +{'positive': 4, 'negative': 1, 'negativeUnattempted': 0}] ◯ PRACTICE VARIANT

ID: Q35 Source: Infosys PYQ Level: Infosys SE Standard

If
in a certain code 3 = 18, 4 = 32, 5 = 50, then 6 = ?

A 60	**B** 70
C 72	**D** 80

✓ **Verified Answer:** ✅ **(C) 72** **Step-by-Step Explanation & Derivation:** **Explanation / Solution:** 70