

# Plan

- Directions of Fea(LPP)
- Extreme directions of Fea(LPP)
- Representation Theorem for Fea(LPP)
- Necessary and sufficient conditions for the existence of optimal solutions
- Optimal solutions in atleast one corner point

- $Fea(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}.$

- $Fea(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}.$
- **Observation 5:** Suppose if a LPP has an **unbounded feasible region**, then there exists a vector  $\mathbf{d} \neq \mathbf{0}$  such that starting from any  $\mathbf{x} \in Fea(LPP)$  and moving in the positive direction of  $\mathbf{d}$ , always gives elements of  $Fea(LPP)$ .

- $Fea(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}.$
- **Observation 5:** Suppose if a LPP has an **unbounded feasible region**, then there exists a vector  $\mathbf{d} \neq \mathbf{0}$  such that starting from any  $\mathbf{x} \in Fea(LPP)$  and moving in the positive direction of  $\mathbf{d}$ , always gives elements of  $Fea(LPP)$ .

That is for any  $\mathbf{x} \in Fea(LPP)$ ,  $\mathbf{x} + \alpha \mathbf{d} \in Fea(LPP)$  for all  $\alpha \geq 0$ ,

- $Fea(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}.$
- **Observation 5:** Suppose if a LPP has an **unbounded feasible region**, then there exists a vector  $\mathbf{d} \neq \mathbf{0}$  such that starting from any  $\mathbf{x} \in Fea(LPP)$  and moving in the positive direction of  $\mathbf{d}$ , always gives elements of  $Fea(LPP)$ .  
That is for any  $\mathbf{x} \in Fea(LPP)$ ,  $\mathbf{x} + \alpha \mathbf{d} \in Fea(LPP)$  for all  $\alpha \geq 0$ ,  
then  $\mathbf{d} \neq \mathbf{0}$  is called a **direction** of  $S = Fea(LPP)$ .

- $Fea(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ .
- **Observation 5:** Suppose if a LPP has an **unbounded feasible region**, then there exists a vector  $\mathbf{d} \neq \mathbf{0}$  such that starting from any  $\mathbf{x} \in Fea(LPP)$  and moving in the positive direction of  $\mathbf{d}$ , always gives elements of  $Fea(LPP)$ .

That is for any  $\mathbf{x} \in Fea(LPP)$ ,  $\mathbf{x} + \alpha \mathbf{d} \in Fea(LPP)$  for all  $\alpha \geq 0$ ,

then  $\mathbf{d} \neq \mathbf{0}$  is called a **direction** of  $S = Fea(LPP)$ .

Throughout our discussion,  $\mathbf{d} = [d_1, \dots, d_n]^T$  will denote a column vector.

- $Fea(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ .
- **Observation 5:** Suppose if a LPP has an **unbounded feasible region**, then there exists a vector  $\mathbf{d} \neq \mathbf{0}$  such that starting from any  $\mathbf{x} \in Fea(LPP)$  and moving in the positive direction of  $\mathbf{d}$ , always gives elements of  $Fea(LPP)$ .

That is for any  $\mathbf{x} \in Fea(LPP)$ ,  $\mathbf{x} + \alpha \mathbf{d} \in Fea(LPP)$  for all  $\alpha \geq 0$ ,

then  $\mathbf{d} \neq \mathbf{0}$  is called a **direction** of  $S = Fea(LPP)$ .

Throughout our discussion,  $\mathbf{d} = [d_1, \dots, d_n]^T$  will denote a column vector.

- **Definition:** Given a non empty **convex set**  $S$ ,  $S \subset \mathbb{R}^n$ ,  $\mathbf{d} \neq \mathbf{0}$  is called a **direction** of  $S$  if for all  $\mathbf{x} \in S$ ,  $\mathbf{x} + \alpha \mathbf{d} \in S$  for all  $\alpha \geq 0$ .

- $Fea(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ .
- **Observation 5:** Suppose if a LPP has an **unbounded feasible region**, then there exists a vector  $\mathbf{d} \neq \mathbf{0}$  such that starting from any  $\mathbf{x} \in Fea(LPP)$  and moving in the positive direction of  $\mathbf{d}$ , always gives elements of  $Fea(LPP)$ .

That is for any  $\mathbf{x} \in Fea(LPP)$ ,  $\mathbf{x} + \alpha \mathbf{d} \in Fea(LPP)$  for all  $\alpha \geq 0$ ,

then  $\mathbf{d} \neq \mathbf{0}$  is called a **direction** of  $S = Fea(LPP)$ .

Throughout our discussion,  $\mathbf{d} = [d_1, \dots, d_n]^T$  will denote a column vector.

- **Definition:** Given a non empty **convex set**  $S$ ,  $S \subset \mathbb{R}^n$ ,  $\mathbf{d} \neq \mathbf{0}$  is called a **direction** of  $S$  if for all  $\mathbf{x} \in S$ ,  $\mathbf{x} + \alpha \mathbf{d} \in S$  for all  $\alpha \geq 0$ .
- If  $\mathbf{d}$  is a **direction** of a convex set  $S$ , then for all  $\gamma > 0$ ,  $\mathbf{x} + \alpha \mathbf{d} = \mathbf{x} + \left(\frac{\alpha}{\gamma}\right) \gamma \mathbf{d} \in S$  for all  $\alpha > 0$ ,



- $Fea(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ .
- **Observation 5:** Suppose if a LPP has an **unbounded feasible region**, then there exists a vector  $\mathbf{d} \neq \mathbf{0}$  such that starting from any  $\mathbf{x} \in Fea(LPP)$  and moving in the positive direction of  $\mathbf{d}$ , always gives elements of  $Fea(LPP)$ .

That is for any  $\mathbf{x} \in Fea(LPP)$ ,  $\mathbf{x} + \alpha \mathbf{d} \in Fea(LPP)$  for all  $\alpha \geq 0$ ,

then  $\mathbf{d} \neq \mathbf{0}$  is called a **direction** of  $S = Fea(LPP)$ .

Throughout our discussion,  $\mathbf{d} = [d_1, \dots, d_n]^T$  will denote a column vector.

- **Definition:** Given a non empty **convex set**  $S$ ,  $S \subset \mathbb{R}^n$ ,  $\mathbf{d} \neq \mathbf{0}$  is called a **direction** of  $S$  if for all  $\mathbf{x} \in S$ ,  $\mathbf{x} + \alpha \mathbf{d} \in S$  for all  $\alpha \geq 0$ .
- If  $\mathbf{d}$  is a **direction** of a convex set  $S$ , then for all  $\gamma > 0$ ,  $\mathbf{x} + \alpha \mathbf{d} = \mathbf{x} + \left(\frac{\alpha}{\gamma}\right) \gamma \mathbf{d} \in S$  for all  $\alpha > 0$ ,  
 $\Rightarrow \gamma \mathbf{d}$  is again a **direction** for all  $\gamma > 0$ .

- Directions  $\mathbf{d}_1, \mathbf{d}_2$  of  $S$  are said to be **distinct** if  $\mathbf{d}_1 \neq \gamma \mathbf{d}_2$  for any  $\gamma > 0$   
( or equivalently  $\mathbf{d}_2 \neq \beta \mathbf{d}_1$  for any  $\beta > 0$ ).

- Directions  $\mathbf{d}_1, \mathbf{d}_2$  of  $S$  are said to be **distinct** if  $\mathbf{d}_1 \neq \gamma \mathbf{d}_2$  for any  $\gamma > 0$   
( or equivalently  $\mathbf{d}_2 \neq \beta \mathbf{d}_1$  for any  $\beta > 0$ ).
- **Example 2: (revisited)** Consider the problem,  
 $\text{Min } -x + 2y$

- Directions  $\mathbf{d}_1, \mathbf{d}_2$  of  $S$  are said to be **distinct** if  $\mathbf{d}_1 \neq \gamma \mathbf{d}_2$  for any  $\gamma > 0$   
( or equivalently  $\mathbf{d}_2 \neq \beta \mathbf{d}_1$  for any  $\beta > 0$ ).
- **Example 2: (revisited)** Consider the problem,  
Min  $-x + 2y$   
subject to  
 $x + 2y \geq 1$

- Directions  $\mathbf{d}_1, \mathbf{d}_2$  of  $S$  are said to be **distinct** if  $\mathbf{d}_1 \neq \gamma \mathbf{d}_2$  for any  $\gamma > 0$   
( or equivalently  $\mathbf{d}_2 \neq \beta \mathbf{d}_1$  for any  $\beta > 0$ ).
- **Example 2: (revisited)** Consider the problem,  
Min  $-x + 2y$   
subject to  
 $x + 2y \geq 1$   
 $-x + y \leq 1,$

- Directions  $\mathbf{d}_1, \mathbf{d}_2$  of  $S$  are said to be **distinct** if  $\mathbf{d}_1 \neq \gamma \mathbf{d}_2$  for any  $\gamma > 0$   
( or equivalently  $\mathbf{d}_2 \neq \beta \mathbf{d}_1$  for any  $\beta > 0$ ).
- **Example 2: (revisited)** Consider the problem,  
Min  $-x + 2y$   
subject to  
 $x + 2y \geq 1$   
 $-x + y \leq 1,$   
 $x \geq 0, y \geq 0.$

- Directions  $\mathbf{d}_1, \mathbf{d}_2$  of  $S$  are said to be **distinct** if  $\mathbf{d}_1 \neq \gamma \mathbf{d}_2$  for any  $\gamma > 0$   
( or equivalently  $\mathbf{d}_2 \neq \beta \mathbf{d}_1$  for any  $\beta > 0$ ).
- **Example 2: (revisited)** Consider the problem,  
Min  $-x + 2y$   
subject to  
 $x + 2y \geq 1$   
 $-x + y \leq 1,$   
 $x \geq 0, y \geq 0.$
- $\mathbf{d}_1 = [1, 1]^T, \mathbf{d}'_1 = [2, 2]^T, \dots$

- Directions  $\mathbf{d}_1, \mathbf{d}_2$  of  $S$  are said to be **distinct** if  $\mathbf{d}_1 \neq \gamma \mathbf{d}_2$  for any  $\gamma > 0$   
( or equivalently  $\mathbf{d}_2 \neq \beta \mathbf{d}_1$  for any  $\beta > 0$ ).
- **Example 2: (revisited)** Consider the problem,  
Min  $-x + 2y$   
subject to  
 $x + 2y \geq 1$   
 $-x + y \leq 1,$   
 $x \geq 0, y \geq 0.$
- $\mathbf{d}_1 = [1, 1]^T, \mathbf{d}'_1 = [2, 2]^T, \dots$   
are all equal as directions of  $Fea(LPP)$ .
- Similarly  $\mathbf{d}_2 = [1, 0]^T, \mathbf{d}'_2 = [2, 0]^T, \dots$



- Directions  $\mathbf{d}_1, \mathbf{d}_2$  of  $S$  are said to be **distinct** if  $\mathbf{d}_1 \neq \gamma \mathbf{d}_2$  for any  $\gamma > 0$   
( or equivalently  $\mathbf{d}_2 \neq \beta \mathbf{d}_1$  for any  $\beta > 0$ ).
- **Example 2: (revisited)** Consider the problem,  
Min  $-x + 2y$   
subject to  
 $x + 2y \geq 1$   
 $-x + y \leq 1,$   
 $x \geq 0, y \geq 0.$
- $\mathbf{d}_1 = [1, 1]^T, \mathbf{d}'_1 = [2, 2]^T, \dots$   
are all equal as directions of  $Fea(LPP)$ .
- Similarly  $\mathbf{d}_2 = [1, 0]^T, \mathbf{d}'_2 = [2, 0]^T, \dots$   
are all equal as directions of  $Fea(LPP)$ .

- Directions  $\mathbf{d}_1, \mathbf{d}_2$  of  $S$  are said to be **distinct** if  $\mathbf{d}_1 \neq \gamma \mathbf{d}_2$  for any  $\gamma > 0$   
( or equivalently  $\mathbf{d}_2 \neq \beta \mathbf{d}_1$  for any  $\beta > 0$ ).
- **Example 2: (revisited)** Consider the problem,  
Min  $-x + 2y$   
subject to  
 $x + 2y \geq 1$   
 $-x + y \leq 1,$   
 $x \geq 0, y \geq 0.$
- $\mathbf{d}_1 = [1, 1]^T, \mathbf{d}'_1 = [2, 2]^T, \dots$   
are all equal as directions of  $Fea(LPP)$ .
- Similarly  $\mathbf{d}_2 = [1, 0]^T, \mathbf{d}'_2 = [2, 0]^T, \dots$   
are all equal as directions of  $Fea(LPP)$ .
- Whereas  $\mathbf{d}_1 = [1, 1]^T, \mathbf{d}_2 = [1, 0]^T$  give two **distinct directions**.

- **Result:** The set of all directions of  
 $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$

- **Result:** The set of all directions of  $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$  is given by

- **Result:** The set of all directions of  
 $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$  is given by  
 $D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, \quad A_{m \times n} \mathbf{d} \leq \mathbf{0}, \quad \mathbf{d} \geq \mathbf{0}\}$  or by

- **Result:** The set of all directions of

$S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$  is given by

$D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, \quad A_{m \times n} \mathbf{d} \leq \mathbf{0}, \quad \mathbf{d} \geq \mathbf{0}\}$  or by

$D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, \quad \mathbf{a}_i^T \mathbf{d} \leq 0, \text{ for all } i = 1, 2, \dots, m, \quad \mathbf{d} \geq \mathbf{0}\}.$

- Result:** The set of all directions of  $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$  is given by  $D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, A_{m \times n} \mathbf{d} \leq \mathbf{0}, \mathbf{d} \geq \mathbf{0}\}$  or by  $D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, \mathbf{a}_i^T \mathbf{d} \leq 0, \text{ for all } i = 1, 2, \dots, m, \mathbf{d} \geq \mathbf{0}\}$ .
- If  $\mathbf{d}_1, \mathbf{d}_2$  are directions of  $S$ , then  $\mathbf{d} = \alpha \mathbf{d}_1 + \beta \mathbf{d}_2$  will again be a direction of  $S$ , for any  $\alpha, \beta$  non negative

- Result:** The set of all directions of  $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$  is given by  $D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, A_{m \times n} \mathbf{d} \leq \mathbf{0}, \mathbf{d} \geq \mathbf{0}\}$  or by  $D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, \mathbf{a}_i^T \mathbf{d} \leq 0, \text{ for all } i = 1, 2, \dots, m, \mathbf{d} \geq \mathbf{0}\}$ .
- If  $\mathbf{d}_1, \mathbf{d}_2$  are directions of  $S$ , then  $\mathbf{d} = \alpha \mathbf{d}_1 + \beta \mathbf{d}_2$  will again be a direction of  $S$ , for any  $\alpha, \beta$  non negative (as long as both  $\alpha, \beta$  are not equal to zero, or  $\alpha + \beta \neq 0$ ).



- Result:** The set of all directions of  $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$  is given by  $D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, A_{m \times n} \mathbf{d} \leq \mathbf{0}, \mathbf{d} \geq \mathbf{0}\}$  or by  $D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, \mathbf{a}_i^T \mathbf{d} \leq 0, \text{ for all } i = 1, 2, \dots, m, \mathbf{d} \geq \mathbf{0}\}$ .
- If  $\mathbf{d}_1, \mathbf{d}_2$  are directions of  $S$ , then  $\mathbf{d} = \alpha \mathbf{d}_1 + \beta \mathbf{d}_2$  will again be a direction of  $S$ , for any  $\alpha, \beta$  non negative (as long as both  $\alpha, \beta$  are not equal to zero, or  $\alpha + \beta \neq 0$ ).
- The set of all directions of  $S = \text{Fea}(LPP)$  is a **convex set**.

- **Definition:** A direction  $\mathbf{d}$  of  $S$  is called an **extreme direction** of  $S$ ,

- **Definition:** A direction  $\mathbf{d}$  of  $S$  is called an **extreme direction** of  $S$ , if it **cannot** be written as a **positive linear combination** of **two distinct directions** of  $S$ ,

- **Definition:** A direction  $\mathbf{d}$  of  $S$  is called an **extreme direction** of  $S$ , if it **cannot** be written as a **positive linear combination** of **two distinct directions** of  $S$ , that is, if  $\mathbf{d}$  an **extreme direction** of  $S$  and

- **Definition:** A direction  $\mathbf{d}$  of  $S$  is called an **extreme direction** of  $S$ , if it **cannot** be written as a **positive linear combination** of **two distinct directions** of  $S$ , that is, if  $\mathbf{d}$  an **extreme direction** of  $S$  and  $\mathbf{d} = \alpha \mathbf{d}_1 + \beta \mathbf{d}_2$ , for  $\alpha, \beta > 0$  and  $\mathbf{d}_1, \mathbf{d}_2 \in D$  then  $\mathbf{d}_1 = \gamma \mathbf{d}_2$  for some  $\gamma > 0$ .

- **Definition:** A direction  $\mathbf{d}$  of  $S$  is called an **extreme direction** of  $S$ , if it **cannot** be written as a **positive linear combination** of **two distinct directions** of  $S$ , that is, if  $\mathbf{d}$  an **extreme direction** of  $S$  and  $\mathbf{d} = \alpha \mathbf{d}_1 + \beta \mathbf{d}_2$ , for  $\alpha, \beta > 0$  and  $\mathbf{d}_1, \mathbf{d}_2 \in D$  then  $\mathbf{d}_1 = \gamma \mathbf{d}_2$  for some  $\gamma > 0$ .
- If  $D$  denotes the set of all directions of  $S$  (  $D = \phi$  if  $S$  is bounded ), then

- **Definition:** A direction  $\mathbf{d}$  of  $S$  is called an **extreme direction** of  $S$ , if it **cannot** be written as a **positive linear combination** of **two distinct directions** of  $S$ , that is, if  $\mathbf{d}$  an **extreme direction** of  $S$  and  $\mathbf{d} = \alpha \mathbf{d}_1 + \beta \mathbf{d}_2$ , for  $\alpha, \beta > 0$  and  $\mathbf{d}_1, \mathbf{d}_2 \in D$  then  $\mathbf{d}_1 = \gamma \mathbf{d}_2$  for some  $\gamma > 0$ .
- If  $D$  denotes the set of all directions of  $S$  ( $D = \emptyset$  if  $S$  is bounded ), then  $D' = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \geq \mathbf{0}, A\mathbf{d} \leq \mathbf{0}, \sum_i d_i = 1\}$  is a set of all **distinct directions** of  $S$ .

- Definition:** A direction  $\mathbf{d}$  of  $S$  is called an **extreme direction** of  $S$ , if it **cannot** be written as a **positive linear combination** of **two distinct directions** of  $S$ , that is, if  $\mathbf{d}$  an **extreme direction** of  $S$  and  $\mathbf{d} = \alpha \mathbf{d}_1 + \beta \mathbf{d}_2$ , for  $\alpha, \beta > 0$  and  $\mathbf{d}_1, \mathbf{d}_2 \in D$  then  $\mathbf{d}_1 = \gamma \mathbf{d}_2$  for some  $\gamma > 0$ .
- If  $D$  denotes the set of all directions of  $S$  ( $D = \emptyset$  if  $S$  is bounded), then  $D' = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \geq \mathbf{0}, A\mathbf{d} \leq \mathbf{0}, \sum_i d_i = 1\}$  is a set of all **distinct directions** of  $S$ .
- Also each  $\mathbf{d} \in D$  is of the form  $\mathbf{d} = \alpha \mathbf{d}'$  for some  $\mathbf{d}' \in D'$  where  $\alpha = \sum_i d_i (> 0)$ .
- $D' = \left\{ \mathbf{d} \in \mathbb{R}^n : \mathbf{d} \geq \mathbf{0}, \begin{bmatrix} & & A & \\ 1 & 1, \dots, & 1 \\ -1 & -1, \dots, & -1 \end{bmatrix} \mathbf{d} \leq \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} \right\}.$



- **Definition:** A direction  $\mathbf{d}$  of  $S$  is called an **extreme direction** of  $S$ , if it **cannot** be written as a **positive linear combination** of **two distinct directions** of  $S$ , that is, if  $\mathbf{d}$  an **extreme direction** of  $S$  and  $\mathbf{d} = \alpha \mathbf{d}_1 + \beta \mathbf{d}_2$ , for  $\alpha, \beta > 0$  and  $\mathbf{d}_1, \mathbf{d}_2 \in D$  then  $\mathbf{d}_1 = \gamma \mathbf{d}_2$  for some  $\gamma > 0$ .

- If  $D$  denotes the set of all directions of  $S$  ( $D = \emptyset$  if  $S$  is bounded), then

$D' = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \geq \mathbf{0}, A\mathbf{d} \leq \mathbf{0}, \sum_i d_i = 1\}$  is a set of all **distinct directions** of  $S$ .

- Also each  $\mathbf{d} \in D$  is of the form  $\mathbf{d} = \alpha \mathbf{d}'$  for some  $\mathbf{d}' \in D'$  where  $\alpha = \sum_i d_i (> 0)$ .

- $D' = \left\{ \mathbf{d} \in \mathbb{R}^n : \mathbf{d} \geq \mathbf{0}, \begin{bmatrix} & & A & \\ 1 & 1, \dots, & 1 \\ -1 & -1, \dots, & -1 \end{bmatrix} \mathbf{d} \leq \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} \right\}.$

The set  $D'$  now looks exactly like the feasible region of a LPP.

- Since  $D'$  is like  $Fea(LPP) = S$ , if  $D' \neq \phi$ ,  $D'$  must have at least one **extreme point**.

- Since  $D'$  is like  $Fea(LPP) = S$ , if  $D' \neq \phi$ ,  $D'$  must have at least one extreme point.
- **Result:**  $\mathbf{d}$  is an extreme direction of  $S$  if and only if  $\mathbf{d}' = \frac{\mathbf{d}}{\sum_i d_i}$  is an extreme point of  $D'$ .

- Since  $D'$  is like  $Fea(LPP) = S$ , if  $D' \neq \phi$ ,  $D'$  must have at least one **extreme point**.
- **Result:**  $\mathbf{d}$  is an **extreme direction** of  $S$  if and only if  $\mathbf{d}' = \frac{\mathbf{d}}{\sum_i d_i}$  is an **extreme point** of  $D'$ .
- If  $D \neq \phi$ , then  $Fea(LPP) = S(\neq \phi)$  must have at least one **extreme direction**.

- Since  $D'$  is like  $Fea(LPP) = S$ , if  $D' \neq \phi$ ,  $D'$  must have at least one **extreme point**.
- **Result:**  $\mathbf{d}$  is an **extreme direction** of  $S$  if and only if  $\mathbf{d}' = \frac{\mathbf{d}}{\sum_i d_i}$  is an **extreme point** of  $D'$ .
- If  $D \neq \phi$ , then  $Fea(LPP) = S(\neq \phi)$  must have at least one **extreme direction**.
- If  $Fea(LPP) = S \neq \phi$  is **unbounded** then  $D \neq \phi$  and  $S$  must have at least one **extreme direction**.

- Since  $D'$  is like  $Fea(LPP) = S$ , if  $D' \neq \phi$ ,  $D'$  must have at least one **extreme point**.
- **Result:**  $\mathbf{d}$  is an **extreme direction** of  $S$  if and only if  $\mathbf{d}' = \frac{\mathbf{d}}{\sum_i d_i}$  is an **extreme point** of  $D'$ .
- If  $D \neq \phi$ , then  $Fea(LPP) = S(\neq \phi)$  must have at least one **extreme direction**.
- If  $Fea(LPP) = S \neq \phi$  is **unbounded** then  $D \neq \phi$  and  $S$  must have at least one **extreme direction**.
- The number of **distinct extreme directions** of  $S$  is **finite** (why?).

- The extreme directions of  $S$  which are extreme points of  $D'$  (after suitable normalization) will lie on  $n$  LI hyperplanes defining  $D'$ .

- The **extreme directions of  $S$**  which are **extreme points** of  $D'$  (after suitable normalization) will lie on  **$n$  LI hyperplanes** defining  $D'$ .
- Since  $\mathbf{d} \in \mathbb{R}^n$ ,  $\mathbf{d} \neq \mathbf{0}$  cannot be orthogonal to  $n$  **LI** vectors,



- The **extreme directions of  $S$**  which are **extreme points** of  $D'$  (after suitable normalization) will lie on  **$n$  LI hyperplanes** defining  $D'$ .
- Since  $\mathbf{d} \in \mathbb{R}^n$ ,  $\mathbf{d} \neq \mathbf{0}$  cannot be orthogonal to  $n$  **LI** vectors, so  $\mathbf{d}$  cannot lie on  **$n$  LI hyperplanes** of the  $(m + n)$  hyperplanes given by,

- The **extreme directions of  $S$**  which are **extreme points** of  $D'$  (after suitable normalization) will lie on  **$n$  LI hyperplanes** defining  $D'$ .
- Since  $\mathbf{d} \in \mathbb{R}^n$ ,  $\mathbf{d} \neq \mathbf{0}$  cannot be orthogonal to  $n$  **LI** vectors, so  $\mathbf{d}$  cannot lie on  **$n$  LI hyperplanes** of the  $(m + n)$  hyperplanes given by,  
 $\{\mathbf{d} \in \mathbb{R}^n : \mathbf{a}_i^T \mathbf{d} = 0\}$  for  $i = 1, 2, \dots, m$ , and

- The **extreme directions of  $S$**  which are **extreme points** of  $D'$  (after suitable normalization) will lie on  **$n$  LI hyperplanes** defining  $D'$ .
- Since  $\mathbf{d} \in \mathbb{R}^n$ ,  $\mathbf{d} \neq \mathbf{0}$  cannot be orthogonal to  $n$  **LI** vectors, so  $\mathbf{d}$  cannot lie on  **$n$  LI hyperplanes** of the  $(m + n)$  hyperplanes given by,

$$\{\mathbf{d} \in \mathbb{R}^n : \mathbf{a}_i^T \mathbf{d} = 0\} \text{ for } i = 1, 2, \dots, m, \text{ and}$$

$$\{\mathbf{d} \in \mathbb{R}^n : -\mathbf{e}_j^T \mathbf{d} = 0\} \text{ for } j = 1, 2, \dots, n.$$

- The **extreme directions of  $S$**  which are **extreme points** of  $D'$  (after suitable normalization) will lie on  **$n$  LI hyperplanes** defining  $D'$ .
- Since  $\mathbf{d} \in \mathbb{R}^n$ ,  $\mathbf{d} \neq \mathbf{0}$  cannot be orthogonal to  $n$  **LI** vectors, so  $\mathbf{d}$  cannot lie on  **$n$  LI hyperplanes** of the  $(m + n)$  hyperplanes given by,
 
$$\{\mathbf{d} \in \mathbb{R}^n : \mathbf{a}_i^T \mathbf{d} = 0\} \text{ for } i = 1, 2, \dots, m, \text{ and}$$

$$\{\mathbf{d} \in \mathbb{R}^n : -\mathbf{e}_j^T \mathbf{d} = 0\} \text{ for } j = 1, 2, \dots, n.$$
- If  $\mathbf{d} \in D'$ , is an **extreme direction** of  $S$  then it should lie on  **$(n - 1)$  LI hyperplanes** of the above mentioned  **$(m + n)$  hyperplanes**,

- The **extreme directions of  $S$**  which are **extreme points** of  $D'$  (after suitable normalization) will lie on  **$n$  LI hyperplanes** defining  $D'$ .
- Since  $\mathbf{d} \in \mathbb{R}^n$ ,  $\mathbf{d} \neq \mathbf{0}$  cannot be orthogonal to  $n$  **LI** vectors, so  $\mathbf{d}$  cannot lie on  **$n$  LI hyperplanes** of the  $(m + n)$  hyperplanes given by,
 
$$\{\mathbf{d} \in \mathbb{R}^n : \mathbf{a}_i^T \mathbf{d} = 0\} \text{ for } i = 1, 2, \dots, m, \text{ and}$$

$$\{\mathbf{d} \in \mathbb{R}^n : -\mathbf{e}_j^T \mathbf{d} = 0\} \text{ for } j = 1, 2, \dots, n.$$
- If  $\mathbf{d} \in D'$ , is an **extreme direction** of  $S$  then it should lie on  **$(n - 1)$  LI hyperplanes** of the above mentioned  **$(m + n)$  hyperplanes**, and the hyperplane
 
$$\{\mathbf{d} \in \mathbb{R}^n : [1, 1, \dots, 1]\mathbf{d} = 1\},$$
 which gives a collection of  **$n$  LI hyperplanes**.

- The **extreme directions of  $S$**  which are **extreme points** of  $D'$  (after suitable normalization) will lie on  **$n$  LI hyperplanes** defining  $D'$ .
- Since  $\mathbf{d} \in \mathbb{R}^n$ ,  $\mathbf{d} \neq \mathbf{0}$  cannot be orthogonal to  $n$  **LI** vectors, so  $\mathbf{d}$  cannot lie on  **$n$  LI hyperplanes** of the  $(m + n)$  hyperplanes given by,
 
$$\{\mathbf{d} \in \mathbb{R}^n : \mathbf{a}_i^T \mathbf{d} = 0\} \text{ for } i = 1, 2, \dots, m, \text{ and}$$

$$\{\mathbf{d} \in \mathbb{R}^n : -\mathbf{e}_j^T \mathbf{d} = 0\} \text{ for } j = 1, 2, \dots, n.$$
- If  $\mathbf{d} \in D'$ , is an **extreme direction** of  $S$  then it should lie on  **$(n - 1)$  LI hyperplanes** of the above mentioned  **$(m + n)$  hyperplanes**, and the hyperplane
 
$$\{\mathbf{d} \in \mathbb{R}^n : [1, 1, \dots, 1]\mathbf{d} = 1\},$$
 which gives a collection of  **$n$  LI hyperplanes**.

- **Exercise:** Check that if a  $\mathbf{d} \in D$  lies on  $(n-1)$  LI hyperplanes (out of the  $(m+n)$  defining hyperplanes of  $D$ ) given by  $\{H_1, \dots, H_{n-1}\}$ , then  $\{H, H_1, \dots, H_{n-1}\}$  is LI where  $H = \{\mathbf{d} \in \mathbb{R}^n : [1, 1, \dots, 1]\mathbf{d} = 1\}$ .

- **Exercise:** Check that if a  $\mathbf{d} \in D$  lies on  $(n - 1)$  LI hyperplanes (out of the  $(m + n)$  defining hyperplanes of  $D$ ) given by  $\{H_1, \dots, H_{n-1}\}$ , then  $\{H, H_1, \dots, H_{n-1}\}$  is LI where  $H = \{\mathbf{d} \in \mathbb{R}^n : [1, 1, \dots, 1]\mathbf{d} = 1\}$ .
- Any  $\mathbf{d} \in D$ , which lies on  $(n - 1)$  LI hyperplanes out of the  $(m + n)$  hyperplanes given by



- **Exercise:** Check that if a  $\mathbf{d} \in D$  lies on  $(n - 1)$  LI hyperplanes (out of the  $(m + n)$  defining hyperplanes of  $D$ ) given by  $\{H_1, \dots, H_{n-1}\}$ , then  $\{H, H_1, \dots, H_{n-1}\}$  is LI where  $H = \{\mathbf{d} \in \mathbb{R}^n : [1, 1, \dots, 1]\mathbf{d} = 1\}$ .
- Any  $\mathbf{d} \in D$ , which lies on  $(n - 1)$  LI hyperplanes out of the  $(m + n)$  hyperplanes given by  $\{\mathbf{d} \in \mathbb{R}^n : \mathbf{a}_i^T \mathbf{d} = 0\}$  for  $i = 1, 2, \dots, m$ , and

- **Exercise:** Check that if a  $\mathbf{d} \in D$  lies on  $(n - 1)$  LI hyperplanes (out of the  $(m + n)$  defining hyperplanes of  $D$ ) given by  $\{H_1, \dots, H_{n-1}\}$ , then  $\{H, H_1, \dots, H_{n-1}\}$  is LI where  $H = \{\mathbf{d} \in \mathbb{R}^n : [1, 1, \dots, 1]\mathbf{d} = 1\}$ .
- Any  $\mathbf{d} \in D$ , which lies on  $(n - 1)$  LI hyperplanes out of the  $(m + n)$  hyperplanes given by  $\{\mathbf{d} \in \mathbb{R}^n : \mathbf{a}_i^T \mathbf{d} = 0\}$  for  $i = 1, 2, \dots, m$ , and  $\{\mathbf{d} \in \mathbb{R}^n : -\mathbf{e}_j^T \mathbf{d} = 0\}$  for  $j = 1, 2, \dots, n$ , is an **extreme direction** of  $S$ .

- **Example 2: (revisited)** Consider the problem,  
 $\text{Min } -x + 2y$   
subject to

- **Example 2: (revisited)** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

- **Example 2: (revisited)** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

- **Example 2: (revisited)** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

- **Example 2: (revisited)** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

Note that here the set of all **directions** of  $S$  is given by

- **Example 2: (revisited)** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

Note that here the set of all **directions** of  $S$  is given by

- $D = \{\mathbf{d} \in \mathbb{R}^2 : [-1, -2]\mathbf{d} \leq 0, [-1, 1]\mathbf{d} \leq 0, \mathbf{d} \geq \mathbf{0}\}.$

Also if  $\mathbf{d} \in D$  is an **extreme direction** of  $S$  then it has to lie on **exactly one** of the hyperplanes given by



- **Example 2: (revisited)** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

Note that here the set of all **directions** of  $S$  is given by

- $D = \{\mathbf{d} \in \mathbb{R}^2 : [-1, -2]\mathbf{d} \leq 0, [-1, 1]\mathbf{d} \leq 0, \mathbf{d} \geq \mathbf{0}\}.$

Also if  $\mathbf{d} \in D$  is an **extreme direction** of  $S$  then it has to lie on **exactly one** of the hyperplanes given by

- (i)  $\{\mathbf{d} \in \mathbb{R}^2 : [-1, -2]\mathbf{d} = 0\},$

- **Example 2: (revisited)** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

Note that here the set of all **directions** of  $S$  is given by

- $D = \{\mathbf{d} \in \mathbb{R}^2 : [-1, -2]\mathbf{d} \leq 0, [-1, 1]\mathbf{d} \leq 0, \mathbf{d} \geq \mathbf{0}\}.$

Also if  $\mathbf{d} \in D$  is an **extreme direction** of  $S$  then it has to lie on **exactly one** of the hyperplanes given by

- (i)  $\{\mathbf{d} \in \mathbb{R}^2 : [-1, -2]\mathbf{d} = 0\},$
- (ii)  $\{\mathbf{d} \in \mathbb{R}^2 : [-1, 1]\mathbf{d} = 0\},$

- **Example 2: (revisited)** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

Note that here the set of all **directions** of  $S$  is given by

- $D = \{\mathbf{d} \in \mathbb{R}^2 : [-1, -2]\mathbf{d} \leq 0, [-1, 1]\mathbf{d} \leq 0, \mathbf{d} \geq \mathbf{0}\}.$

Also if  $\mathbf{d} \in D$  is an **extreme direction** of  $S$  then it has to lie on **exactly one** of the hyperplanes given by

- (i)  $\{\mathbf{d} \in \mathbb{R}^2 : [-1, -2]\mathbf{d} = 0\},$
- (ii)  $\{\mathbf{d} \in \mathbb{R}^2 : [-1, 1]\mathbf{d} = 0\},$
- (iii)  $\{\mathbf{d} \in \mathbb{R}^2 : d_1 = 0\},$

- **Example 2: (revisited)** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

Note that here the set of all **directions** of  $S$  is given by

- $D = \{\mathbf{d} \in \mathbb{R}^2 : [-1, -2]\mathbf{d} \leq 0, [-1, 1]\mathbf{d} \leq 0, \mathbf{d} \geq \mathbf{0}\}.$

Also if  $\mathbf{d} \in D$  is an **extreme direction** of  $S$  then it has to lie on **exactly one** of the hyperplanes given by

- (i)  $\{\mathbf{d} \in \mathbb{R}^2 : [-1, -2]\mathbf{d} = 0\},$
- (ii)  $\{\mathbf{d} \in \mathbb{R}^2 : [-1, 1]\mathbf{d} = 0\},$
- (iii)  $\{\mathbf{d} \in \mathbb{R}^2 : d_1 = 0\},$
- (iv)  $\{\mathbf{d} \in \mathbb{R}^2 : d_2 = 0\}.$

- Note that there exists no  $d \geq 0, d \neq 0$  such that  $[-1, -2]d = 0$ .

- Note that there exists no  $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$  such that  $[-1, -2]\mathbf{d} = 0$ .
- Also if  $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$ , satisfies the condition  $d_1 = 0$ , then  $[-1, 1]\mathbf{d} \leq 0$  cannot be satisfied, hence such a  $\mathbf{d}$  does not belong to  $D$ .

- Note that there exists no  $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$  such that  $[-1, -2]\mathbf{d} = 0$ .
- Also if  $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$ , satisfies the condition  $d_1 = 0$ , then  $[-1, 1]\mathbf{d} \leq 0$  cannot be satisfied, hence such a  $\mathbf{d}$  does not belong to  $D$ .
- Hence if  $\mathbf{d} \in D$ , is an **extreme direction** of  $S$  then it lies on of following Hyperplanes:

- Note that there exists no  $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$  such that  $[-1, -2]\mathbf{d} = 0$ .
- Also if  $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$ , satisfies the condition  $d_1 = 0$ , then  $[-1, 1]\mathbf{d} \leq 0$  cannot be satisfied, hence such a  $\mathbf{d}$  does not belong to  $D$ .
- Hence if  $\mathbf{d} \in D$ , is an **extreme direction** of  $S$  then it lies on of following Hyperplanes:
- $\{\mathbf{d} \in \mathbb{R}^2 : [-1, 1]\mathbf{d} = 0\}, \{\mathbf{d} \in \mathbb{R}^2 : d_2 = 0\}$ .



- Note that there exists no  $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$  such that  $[-1, -2]\mathbf{d} = \mathbf{0}$ .
- Also if  $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$ , satisfies the condition  $d_1 = 0$ , then  $[-1, 1]\mathbf{d} \leq \mathbf{0}$  cannot be satisfied, hence such a  $\mathbf{d}$  does not belong to  $D$ .
- Hence if  $\mathbf{d} \in D$ , is an **extreme direction** of  $S$  then it lies on of following Hyperplanes:
- $\{\mathbf{d} \in \mathbb{R}^2 : [-1, 1]\mathbf{d} = \mathbf{0}\}, \{\mathbf{d} \in \mathbb{R}^2 : d_2 = 0\}$ .
- $\mathbf{d}' = [1, 1]^T$  and any **positive scalar multiple** of  $\mathbf{d}'$ , are **extreme directions** of  $S = \text{Fea}(LPP)$ .

- Note that there exists no  $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$  such that  $[-1, -2]\mathbf{d} = \mathbf{0}$ .
- Also if  $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$ , satisfies the condition  $d_1 = 0$ , then  $[-1, 1]\mathbf{d} \leq \mathbf{0}$  cannot be satisfied, hence such a  $\mathbf{d}$  does not belong to  $D$ .
- Hence if  $\mathbf{d} \in D$ , is an **extreme direction** of  $S$  then it lies on of following Hyperplanes:
- $\{\mathbf{d} \in \mathbb{R}^2 : [-1, 1]\mathbf{d} = \mathbf{0}\}, \{\mathbf{d} \in \mathbb{R}^2 : d_2 = 0\}$ .
- $\mathbf{d}' = [1, 1]^T$  and any **positive scalar multiple** of  $\mathbf{d}'$ , are **extreme directions** of  $S = \text{Fea}(LPP)$ .
- $\mathbf{d}'' = [1, 0]^T$  and any **positive scalar multiple** of  $\mathbf{d}''$ , are **extreme directions** of  $S = \text{Fea}(LPP)$ .

- Note that there exists no  $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$  such that  $[-1, -2]\mathbf{d} = 0$ .
- Also if  $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$ , satisfies the condition  $d_1 = 0$ , then  $[-1, 1]\mathbf{d} \leq 0$  cannot be satisfied, hence such a  $\mathbf{d}$  does not belong to  $D$ .
- Hence if  $\mathbf{d} \in D$ , is an **extreme direction** of  $S$  then it lies on of following Hyperplanes:
- $\{\mathbf{d} \in \mathbb{R}^2 : [-1, 1]\mathbf{d} = 0\}, \{\mathbf{d} \in \mathbb{R}^2 : d_2 = 0\}$ .
- $\mathbf{d}' = [1, 1]^T$  and any **positive scalar multiple** of  $\mathbf{d}'$ , are **extreme directions** of  $S = \text{Fea}(LPP)$ .
- $\mathbf{d}'' = [1, 0]^T$  and any **positive scalar multiple** of  $\mathbf{d}''$ , are **extreme directions** of  $S = \text{Fea}(LPP)$ .
- These are the only **extreme directions** of  $S = \text{Fea}(LPP)$ .

- **Theorem:**

If  $S = \text{Fea}(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$  is nonempty, then  $S$  has at least one **extreme point**.

- **Theorem:**

If  $S = \text{Fea}(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$  is nonempty, then  $S$  has at least one **extreme point**.

- **Remark:** Note that the above result is **not necessarily true** for all **polyhedral sets**.

For example take any **single half space**, or say a **straight line** in  $\mathbb{R}^n$ , which are **polyhedral sets**, but does not have any **extreme point**.

- **Theorem:**

If  $S = \text{Fea}(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$  is nonempty, then  $S$  has at least one **extreme point**.

- **Remark:** Note that the above result is **not necessarily true** for all **polyhedral sets**.

For example take any **single half space**, or say a **straight line** in  $\mathbb{R}^n$ , which are **polyhedral sets**, but does not have any **extreme point**.

- The theorem works for  $\text{Fea}(LPP)$  because of the **non negativity constraints**, that is because  $\text{Fea}(LPP)$  is given a supply of  **$n$  LI hyperplanes**, **defining hyperplanes**.

- **Exercise:** Can you find a nonempty polyhedral set  $S$ ,  $S \subset \mathbb{R}^2$  which has two defining hyperplanes but does not have any extreme point?

- **Exercise:** Can you find a nonempty polyhedral set  $S$ ,  $S \subset \mathbb{R}^2$  which has two defining hyperplanes but does not have any extreme point?
- **Exercise:** Can you find a nonempty polyhedral set  $S$ ,  $S \subset \mathbb{R}^3$  which has two LI defining hyperplanes but does not have any extreme point?



- **Exercise:** Can you find a nonempty polyhedral set  $S$ ,  $S \subset \mathbb{R}^2$  which has two defining hyperplanes but does not have any extreme point?
- **Exercise:** Can you find a nonempty polyhedral set  $S$ ,  $S \subset \mathbb{R}^3$  which has two LI defining hyperplanes but does not have any extreme point?
- **Exercise:** Is it possible to find a nonempty polyhedral set  $S$ ,  $S \subset \mathbb{R}^3$  which has three LI defining hyperplanes ( not necessarily the nonnegativity constraints) but does not have any extreme point.

- **Exercise:** Can you find a nonempty polyhedral set  $S$ ,  $S \subset \mathbb{R}^2$  which has two defining hyperplanes but does not have any extreme point?
- **Exercise:** Can you find a nonempty polyhedral set  $S$ ,  $S \subset \mathbb{R}^3$  which has two LI defining hyperplanes but does not have any extreme point?
- **Exercise:** Is it possible to find a nonempty polyhedral set  $S$ ,  $S \subset \mathbb{R}^3$  which has three LI defining hyperplanes (not necessarily the nonnegativity constraints) but does not have any extreme point.
- **Definition:** Given  $S$ , a nonempty subset of  $\mathbb{R}^n$ , and  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k \in S$ ,  $\sum_{i=1}^k \lambda_i \mathbf{x}_i$ , is called a convex combination of  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$ , where  $0 \leq \lambda_i \leq 1$  for all  $i = 1, 2, \dots, k$ , and  $\sum_{i=1}^k \lambda_i = 1$ .

- All possible **convex combinations** of **two distinct points** gives a **straight line segment** with those two points as boundary points.

- All possible **convex combinations** of **two distinct points** gives a **straight line segment** with those two points as boundary points.
- All possible convex combinations of **three non colinear points** gives a **triangle** with those points as corner points.

- All possible **convex combinations** of **two distinct points** gives a **straight line segment** with those two points as boundary points.
- All possible convex combinations of **three non colinear points** gives a **triangle** with those points as corner points.
- All possible convex combinations of **four points** no three of which are colinear gives a **quadrilateral**.

- All possible **convex combinations** of **two distinct points** gives a **straight line segment** with those two points as boundary points.
- All possible convex combinations of **three non colinear points** gives a **triangle** with those points as corner points.
- All possible convex combinations of **four points** no three of which are colinear gives a **quadrilateral**.
- **Result:** Given  $\phi \neq S \subset \mathbb{R}^n$ ,  $S$  is a **convex set** if and only if for all  $k \in \mathbb{N}$ , the **convex combination of any  $k$  elements** of  $S$  is again an element of  $S$ .

- **Theorem: (Representation Theorem)**

If  $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$  is nonempty and if  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$  are the extreme points of  $S$  and  $\mathbf{d}_1, \mathbf{d}_2, \dots, \mathbf{d}_r$  are the distinct extreme directions of  $S$  (the set of directions is empty if  $S$  is bounded) then  $\mathbf{x} \in S$  if and only if

- **Theorem: (Representation Theorem)**

If  $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$  is nonempty and if  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$  are the extreme points of  $S$  and  $\mathbf{d}_1, \mathbf{d}_2, \dots, \mathbf{d}_r$  are the distinct extreme directions of  $S$  (the set of directions is empty if  $S$  is bounded) then  $\mathbf{x} \in S$  if and only if

$$\mathbf{x} = \sum_{i=1}^k \lambda_i \mathbf{x}_i + \sum_{j=1}^r \mu_j \mathbf{d}_j$$



- **Theorem: (Representation Theorem)**

If  $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$  is nonempty and if  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$  are the extreme points of  $S$  and  $\mathbf{d}_1, \mathbf{d}_2, \dots, \mathbf{d}_r$  are the distinct extreme directions of  $S$  (the set of directions is empty if  $S$  is bounded) then  $\mathbf{x} \in S$  if and only if

$$\mathbf{x} = \sum_{i=1}^k \lambda_i \mathbf{x}_i + \sum_{j=1}^r \mu_j \mathbf{d}_j$$

where  $0 \leq \lambda_i \leq 1$  for all  $i = 1, 2, \dots, k$ ,  $\sum_i \lambda_i = 1$ ,

- **Theorem: (Representation Theorem)**

If  $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$  is nonempty and if  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$  are the extreme points of  $S$  and  $\mathbf{d}_1, \mathbf{d}_2, \dots, \mathbf{d}_r$  are the distinct extreme directions of  $S$  (the set of directions is empty if  $S$  is bounded) then  $\mathbf{x} \in S$  if and only if

$$\mathbf{x} = \sum_{i=1}^k \lambda_i \mathbf{x}_i + \sum_{j=1}^r \mu_j \mathbf{d}_j$$

where  $0 \leq \lambda_i \leq 1$  for all  $i = 1, 2, \dots, k$ ,  $\sum_i \lambda_i = 1$ ,  
and  $\mu_j \geq 0$ , for all  $j = 1, 2, \dots, r$ .

- **Theorem: (Representation Theorem)**

If  $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$  is nonempty and if  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$  are the extreme points of  $S$  and  $\mathbf{d}_1, \mathbf{d}_2, \dots, \mathbf{d}_r$  are the distinct extreme directions of  $S$  (the set of directions is empty if  $S$  is bounded) then  $\mathbf{x} \in S$  if and only if

$$\mathbf{x} = \sum_{i=1}^k \lambda_i \mathbf{x}_i + \sum_{j=1}^r \mu_j \mathbf{d}_j$$

where  $0 \leq \lambda_i \leq 1$  for all  $i = 1, 2, \dots, k$ ,  $\sum_i \lambda_i = 1$ , and  $\mu_j \geq 0$ , for all  $j = 1, 2, \dots, r$ .

- That is,  $\mathbf{x} \in S \Leftrightarrow \mathbf{x}$  can be written as a convex combination of the extreme points of  $S$  plus a non negative linear combination of the extreme directions of  $S$ .

- **Observation 6:** If  $S = \text{Fea}(LPP)$  is a nonempty bounded set then any  $\mathbf{x} \in S$  is a convex combination of the extreme points of  $S$ .

- **Observation 6:** If  $S = \text{Fea}(LPP)$  is a nonempty **bounded set** then any  $\mathbf{x} \in S$  is a **convex combination** of the extreme points of  $S$ .
- **Observation 7:** Since  $D'$ , the set of distinct directions of  $S$  (if it is nonempty) is a **bounded set** ( $\mathbf{d} \geq \mathbf{0}$  and  $\sum_{i=1}^n d_i = 1$ ),

- **Observation 6:** If  $S = \text{Fea}(LPP)$  is a nonempty bounded set then any  $\mathbf{x} \in S$  is a convex combination of the extreme points of  $S$ .
- **Observation 7:** Since  $D'$ , the set of distinct directions of  $S$  (if it is nonempty) is a bounded set ( $\mathbf{d} \geq \mathbf{0}$  and  $\sum_{i=1}^n d_i = 1$ ), so any  $\mathbf{d} \in D'$  is a convex combination of the extreme points of  $D'$ .

- **Observation 6:** If  $S = \text{Fea}(LPP)$  is a nonempty bounded set then any  $\mathbf{x} \in S$  is a convex combination of the extreme points of  $S$ .
- **Observation 7:** Since  $D'$ , the set of distinct directions of  $S$  (if it is nonempty) is a bounded set ( $\mathbf{d} \geq \mathbf{0}$  and  $\sum_{i=1}^n d_i = 1$ ),  
so any  $\mathbf{d} \in D'$  is a convex combination of the extreme points of  $D'$ .
- So any direction  $\mathbf{d} \in D$  can be written as a nonnegative linear combination of the extreme directions of  $S$ .

- **Observation 8:** Note that if there exists a  $\mathbf{d} \in D$  such that  $\mathbf{c}^T \mathbf{d} < 0$  then the LPP(\*)



- **Observation 8:** Note that if there exists a  $\mathbf{d} \in D$  such that  $\mathbf{c}^T \mathbf{d} < 0$  then the LPP(\*)  
(\*) Min  $\mathbf{c}^T \mathbf{x}$ ,

- **Observation 8:** Note that if there exists a  $\mathbf{d} \in D$  such that  $\mathbf{c}^T \mathbf{d} < 0$  then the LPP(\*)  
( (\*) Min  $\mathbf{c}^T \mathbf{x}$ , subject to  $A\mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$  )

- **Observation 8:** Note that if there exists a  $\mathbf{d} \in D$  such that  $\mathbf{c}^T \mathbf{d} < 0$  then the LPP(\*)  
( (\*) Min  $\mathbf{c}^T \mathbf{x}$ , subject to  $A\mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$  )  
does not have an optimal solution.

- **Observation 8:** Note that if there exists a  $\mathbf{d} \in D$  such that  $\mathbf{c}^T \mathbf{d} < 0$  then the LPP(\*)  
( (\*) Min  $\mathbf{c}^T \mathbf{x}$ , subject to  $A\mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$  )  
does not have an optimal solution.  
Given  $\mathbf{x} \in S$ , and real  $M$ ,

- **Observation 8:** Note that if there exists a  $\mathbf{d} \in D$  such that  $\mathbf{c}^T \mathbf{d} < 0$  then the LPP(\*)  
( (\*) Min  $\mathbf{c}^T \mathbf{x}$ , subject to  $A\mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$  )  
does not have an optimal solution.  
Given  $\mathbf{x} \in S$ , and real  $M$ ,  
 $\mathbf{c}^T (\mathbf{x} + \alpha \mathbf{d}) = \mathbf{c}^T \mathbf{x} + \alpha \mathbf{c}^T \mathbf{d} < M$ ,  
for  $\alpha > 0$  sufficiently large.

- **Observation 8:** Note that if there exists a  $\mathbf{d} \in D$  such that  $\mathbf{c}^T \mathbf{d} < 0$  then the LPP(\*)  
( (\*) Min  $\mathbf{c}^T \mathbf{x}$ , subject to  $A\mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$  )  
does not have an optimal solution.  
Given  $\mathbf{x} \in S$ , and real  $M$ ,  
 $\mathbf{c}^T (\mathbf{x} + \alpha \mathbf{d}) = \mathbf{c}^T \mathbf{x} + \alpha \mathbf{c}^T \mathbf{d} < M$ ,  
for  $\alpha > 0$  sufficiently large.

- **Exercise:** If  $\mathbf{c}^T \mathbf{d}_j \geq 0$  for all extreme directions  $\mathbf{d}_j$  of the nonempty and unbounded feasible region  $S$  of a LPP, then does it imply that  $\mathbf{c}^T \mathbf{d} \geq 0$  for all directions  $\mathbf{d} \in D$ , of the feasible region  $S$ ?

- **Exercise:** If  $\mathbf{c}^T \mathbf{d}_j \geq 0$  for all extreme directions  $\mathbf{d}_j$  of the nonempty and unbounded feasible region  $S$  of a LPP, then does it imply that  $\mathbf{c}^T \mathbf{d} \geq 0$  for all directions  $\mathbf{d} \in D$ , of the feasible region  $S$ ?

**Ans** is **yes**.



- **Exercise:** If  $\mathbf{c}^T \mathbf{d}_j \geq 0$  for all extreme directions  $\mathbf{d}_j$  of the nonempty and unbounded feasible region  $S$  of a LPP, then does it imply that  $\mathbf{c}^T \mathbf{d} \geq 0$  for all directions  $\mathbf{d} \in D$ , of the feasible region  $S$ ?

**Ans** is **yes**.

- **Observation 9:** From the representation theorem of  $S$  we can see that if  $S \neq \phi$  and  $\mathbf{c}^T \mathbf{d}_j \geq 0$  for all  $j = 1, 2, \dots, r$ , then LPP(\*) has an optimal solution, and atleast one optimal solution is attained at an extreme point of  $S$ .

- **Observation 10:** From the **representation theorem** of  $S$  we can also see that if  $S = \text{Fea}(LPP)$  is nonempty and **bounded** then the  $LPP(*)$  has an **optimal solution** and the **optimal value** is attained in at least one **extreme point**.

- **Observation 10:** From the **representation theorem** of  $S$  we can also see that if  $S = \text{Fea}(LPP)$  is nonempty and **bounded** then the  $LPP(*)$  has an **optimal solution** and the **optimal value** is attained in at least one **extreme point**.

From the above observations we can conclude the following:

- **Observation 10:** From the **representation theorem** of  $S$  we can also see that if  $S = \text{Fea}(LPP)$  is nonempty and **bounded** then the  $LPP(*)$  has an **optimal solution** and the **optimal value** is attained in at least one **extreme point**.

From the above observations we can conclude the following:

- **Conclusion 1:** If  $S = \text{Fea}(LPP) \neq \phi$ , then the  $LPP (*)$  has an **optimal solution** if and only if one of the following is true:

- **Observation 10:** From the **representation theorem** of  $S$  we can also see that if  $S = \text{Fea}(LPP)$  is nonempty and **bounded** then the  $LPP(*)$  has an **optimal solution** and the **optimal value** is attained in at least one **extreme point**.

From the above observations we can conclude the following:

- **Conclusion 1:** If  $S = \text{Fea}(LPP) \neq \emptyset$ , then the  $LPP (*)$  has an **optimal solution** if and only if one of the following is true:
- (i)  $S = \text{Fea}(LPP)$  is **bounded** (also seen before by using extreme value theorem)
- (ii)  $S = \text{Fea}(LPP)$  is **unbounded** and  $\mathbf{c}^T \mathbf{d}_j \geq 0$  for all extreme directions  $\mathbf{d}_j$  of the feasible region  $S$ .

- **Observation 10:** From the **representation theorem** of  $S$  we can also see that if  $S = \text{Fea}(LPP)$  is nonempty and **bounded** then the  $LPP(*)$  has an **optimal solution** and the **optimal value** is attained in at least one **extreme point**.

From the above observations we can conclude the following:

- **Conclusion 1:** If  $S = \text{Fea}(LPP) \neq \emptyset$ , then the  $LPP (*)$  has an **optimal solution** if and only if one of the following is true:
- (i)  $S = \text{Fea}(LPP)$  is **bounded** (also seen before by using extreme value theorem)
- (ii)  $S = \text{Fea}(LPP)$  is **unbounded** and  $\mathbf{c}^T \mathbf{d}_j \geq 0$  for all extreme directions  $\mathbf{d}_j$  of the feasible region  $S$ .

- **Conclusion 2:** If LPP (\*) has an optimal solution then there exists an extreme point of the feasible region  $S$ , which is an optimal solution.

- **Conclusion 2:** If LPP (\*) has an **optimal solution** then there exists an **extreme point** of the feasible region  $S$ , which is an **optimal solution**.
- **Exercise:** Give an example of a **nonlinear** function  $f : S \rightarrow \mathbb{R}$ , where  $S \subset \mathbb{R}$  is a **closed** and **bounded polyhedral** subset of  $\mathbb{R}$ , (what are these sets?) such that  $f$  has a **minimum value** in  $S$  but the minimum value is not attained at any **extreme point** of  $S$ .



- **Conclusion 2:** If LPP (\*) has an optimal solution then there exists an extreme point of the feasible region  $S$ , which is an optimal solution.
- **Exercise:** Give an example of a nonlinear function  $f : S \rightarrow \mathbb{R}$ , where  $S \subset \mathbb{R}$  is a closed and bounded polyhedral subset of  $\mathbb{R}$ , (what are these sets?) such that  $f$  has a minimum value in  $S$  but the minimum value is not attained at any extreme point of  $S$ .
- **Conclusion 3:** If  $S = \text{Fea}(LPP)$  is nonempty,

- **Conclusion 2:** If LPP (\*) has an **optimal solution** then there exists an **extreme point** of the feasible region  $S$ , which is an **optimal solution**.
- **Exercise:** Give an example of a **nonlinear** function  $f : S \rightarrow \mathbb{R}$ , where  $S \subset \mathbb{R}$  is a **closed** and **bounded polyhedral** subset of  $\mathbb{R}$ , (what are these sets?) such that  $f$  has a **minimum value** in  $S$  but the minimum value is not attained at any **extreme point** of  $S$ .
- **Conclusion 3:** If  $S = \text{Fea}(LPP)$  is nonempty, and there exists an  $M \in \mathbb{R}$  such that for all  $\mathbf{x} \in S$ ,  $\mathbf{c}^T \mathbf{x} \geq M$ ,

- **Conclusion 2:** If LPP (\*) has an optimal solution then there exists an extreme point of the feasible region  $S$ , which is an optimal solution.
- **Exercise:** Give an example of a nonlinear function  $f : S \rightarrow \mathbb{R}$ , where  $S \subset \mathbb{R}$  is a closed and bounded polyhedral subset of  $\mathbb{R}$ , (what are these sets?) such that  $f$  has a minimum value in  $S$  but the minimum value is not attained at any extreme point of  $S$ .
- **Conclusion 3:** If  $S = \text{Fea}(LPP)$  is nonempty, and there exists an  $M \in \mathbb{R}$  such that for all  $\mathbf{x} \in S$ ,  $\mathbf{c}^T \mathbf{x} \geq M$ , then the LPP (\*) has an optimal solution.

- To understand the significance of the previous result solve the following problems.

- To understand the significance of the previous result solve the following problems.
- **Exercise:** Give an example of a **linear** function  $f : S \rightarrow \mathbb{R}$ , where  $S \subset \mathbb{R}$  is not a polyhedral subset of  $\mathbb{R}$ , such that  $f(x) \geq 1$  but  $f$  **does not** have a minimum value in  $S$ .

- To understand the significance of the previous result solve the following problems.
- **Exercise:** Give an example of a **linear** function  $f : S \rightarrow \mathbb{R}$ , where  $S \subset \mathbb{R}$  is not a polyhedral subset of  $\mathbb{R}$ , such that  $f(x) \geq 1$  but  $f$  **does not** have a minimum value in  $S$ .
- **Exercise:** Give an example of a **nonlinear** function  $f : S \rightarrow \mathbb{R}$ , where  $S \subset \mathbb{R}$  is a **polyhedral** subset of  $\mathbb{R}$ , such that  $f(x) \geq 1$  but  $f$  **does not** have a minimum value in  $S$ .

- We can come to similar conclusions if we consider a linear programming problem, LPP(\*\*) as

- We can come to similar conclusions if we consider a linear programming problem, LPP(\*\*) as  
(\*\*)  $\text{Max } \mathbf{c}^T \mathbf{x}$



- We can come to similar conclusions if we consider a linear programming problem, LPP(\*\*) as

$$(**) \text{Max } \mathbf{c}^T \mathbf{x}$$

subject to  $\mathbf{Ax} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$ .

- We can come to similar conclusions if we consider a linear programming problem, LPP(\*\*) as  
(\*\*)  $\text{Max } \mathbf{c}^T \mathbf{x}$   
subject to  $\mathbf{Ax} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$ .
- **Conclusion 1a:** If  $S = \text{Fea}(LPP) \neq \phi$ , then the LPP (\*\*) has an optimal solution if and only if one of the following is true:

- We can come to similar conclusions if we consider a linear programming problem, LPP(\*\*) as  
(\*\*)  $\text{Max } \mathbf{c}^T \mathbf{x}$   
subject to  $\mathbf{Ax} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$ .
- **Conclusion 1a:** If  $S = \text{Fea}(LPP) \neq \phi$ , then the LPP (\*\*) has an optimal solution if and only if one of the following is true:
  - (i)  $S = \text{Fea}(LPP)$  is bounded

- We can come to similar conclusions if we consider a linear programming problem, LPP(\*\*) as

$$(**) \text{Max } \mathbf{c}^T \mathbf{x}$$

subject to  $\mathbf{Ax} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$ .

- **Conclusion 1a:** If  $S = \text{Fea}(LPP) \neq \phi$ , then the LPP (\*\*) has an optimal solution if and only if one of the following is true:

(i)  $S = \text{Fea}(LPP)$  is bounded

(ii)  $S = \text{Fea}(LPP)$  is unbounded and  $\mathbf{c}^T \mathbf{d}_j \leq 0$  for all extreme directions  $\mathbf{d}_j$  of the feasible region  $S$ .

- We can come to similar conclusions if we consider a linear programming problem, LPP(\*\*) as

$$(**) \text{Max } \mathbf{c}^T \mathbf{x}$$

subject to  $\mathbf{Ax} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$ .

- **Conclusion 1a:** If  $S = \text{Fea}(LPP) \neq \phi$ , then the LPP (\*\*) has an optimal solution if and only if one of the following is true:

(i)  $S = \text{Fea}(LPP)$  is bounded

(ii)  $S = \text{Fea}(LPP)$  is unbounded and  $\mathbf{c}^T \mathbf{d}_j \leq 0$  for all extreme directions  $\mathbf{d}_j$  of the feasible region  $S$ .

- **Conclusion 2a:** If a LPP (\*\*) has an optimal solution then there exists an extreme point of the feasible region  $S$ , which is an optimal solution.

- We can come to similar conclusions if we consider a linear programming problem, LPP(\*\*) as

$$(**) \text{Max } \mathbf{c}^T \mathbf{x}$$

subject to  $\mathbf{Ax} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$ .

- **Conclusion 1a:** If  $S = \text{Fea}(LPP) \neq \emptyset$ , then the LPP (\*\*) has an optimal solution if and only if one of the following is true:

(i)  $S = \text{Fea}(LPP)$  is bounded

(ii)  $S = \text{Fea}(LPP)$  is unbounded and  $\mathbf{c}^T \mathbf{d}_j \leq 0$  for all extreme directions  $\mathbf{d}_j$  of the feasible region  $S$ .

- **Conclusion 2a:** If a LPP (\*\*) has an optimal solution then there exists an extreme point of the feasible region  $S$ , which is an optimal solution.
- **Conclusion 3a:** If  $S = \text{Fea}(LPP)$  is nonempty, and there exists an  $M \in \mathbb{R}$  such that for all  $\mathbf{x} \in S, \mathbf{c}^T \mathbf{x} \leq M$ ,

- We can come to similar conclusions if we consider a linear programming problem, LPP(\*\*) as

$$(**) \text{Max } \mathbf{c}^T \mathbf{x}$$

subject to  $\mathbf{Ax} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$ .

- **Conclusion 1a:** If  $S = \text{Fea}(LPP) \neq \phi$ , then the LPP (\*\*) has an optimal solution if and only if one of the following is true:

(i)  $S = \text{Fea}(LPP)$  is bounded

(ii)  $S = \text{Fea}(LPP)$  is unbounded and  $\mathbf{c}^T \mathbf{d}_j \leq 0$  for all extreme directions  $\mathbf{d}_j$  of the feasible region  $S$ .

- **Conclusion 2a:** If a LPP (\*\*) has an optimal solution then there exists an extreme point of the feasible region  $S$ , which is an optimal solution.
- **Conclusion 3a:** If  $S = \text{Fea}(LPP)$  is nonempty, and there exists an  $M \in \mathbb{R}$  such that for all  $\mathbf{x} \in S, \mathbf{c}^T \mathbf{x} \leq M$ , then the LPP (\*\*) has an optimal solution.