

Plan

- Basic Feasible Solution (BFS)
- Non degenerate BFS
- Degenerate BFS
- Simplex Algorithm

- Consider LPP(P)
Max or Min $\mathbf{c}^T \mathbf{x}$
subject to $\mathbf{A}_{m \times n} \mathbf{x} = \mathbf{b}_{m \times 1}$, $\mathbf{x} \geq \mathbf{0}$,
where $\text{rank}(\mathbf{A}) = m$. (*)

- Consider LPP(P)
Max or Min $\mathbf{c}^T \mathbf{x}$
subject to $\mathbf{A}_{m \times n} \mathbf{x} = \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0}$,
where $\text{rank}(\mathbf{A}) = m$.
 $\text{Fea}(P) = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{A}\mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}.$

- If (P) is of the form:

- Consider LPP(P)

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{m \times n} \mathbf{x} = \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0},$ (*)

where $\text{rank}(\mathbf{A}) = m.$

$$\text{Fea}(P) = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{A}\mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}.$$

- If (P) is of the form:

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{m \times n} \mathbf{x} \leq \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0},$

- Consider LPP(P)

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{m \times n} \mathbf{x} = \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0}$, (*)

where $\text{rank}(\mathbf{A}) = m$.

$$\text{Fea}(P) = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{A}\mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}.$$

- If (P) is of the form:

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{m \times n} \mathbf{x} \leq \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0}$,

we add some nonnegative variables and transform the set of constraints as,

- Consider LPP(P)

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{m \times n} \mathbf{x} = \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0}, \quad (*)$

where $\text{rank}(\mathbf{A}) = m$.

$$\text{Fea}(P) = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{A}\mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}.$$

- If (P) is of the form:

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{m \times n} \mathbf{x} \leq \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0},$

we add some nonnegative variables and transform the set of constraints as,

$$\mathbf{A}_{m \times n} \mathbf{x} + \mathbf{s}_{m \times 1} = [\mathbf{A} : \mathbf{I}] \begin{bmatrix} \mathbf{x} \\ \mathbf{s} \end{bmatrix} = \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0}, \mathbf{s} \geq \mathbf{0}.$$

- Consider LPP(P)

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{m \times n} \mathbf{x} = \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0}, \quad (*)$

where $\text{rank}(\mathbf{A}) = m$.

$$\text{Fea}(P) = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{A}\mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}.$$

- If (P) is of the form:

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{m \times n} \mathbf{x} \leq \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0},$

we add some nonnegative variables and transform the set of constraints as,

$$\mathbf{A}_{m \times n} \mathbf{x} + \mathbf{s}_{m \times 1} = [\mathbf{A} : \mathbf{I}] \begin{bmatrix} \mathbf{x} \\ \mathbf{s} \end{bmatrix} = \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0}, \mathbf{s} \geq \mathbf{0}.$$

- If suppose we are given a problem of the type:

- Consider LPP(P)

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{m \times n} \mathbf{x} = \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0}$, (*)

where $\text{rank}(\mathbf{A}) = m$.

$$\text{Fea}(P) = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{A}\mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}.$$

- If (P) is of the form:

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{m \times n} \mathbf{x} \leq \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0}$,

we add some nonnegative variables and transform the set of constraints as,

$$\mathbf{A}_{m \times n} \mathbf{x} + \mathbf{s}_{m \times 1} = [\mathbf{A} : \mathbf{I}] \begin{bmatrix} \mathbf{x} \\ \mathbf{s} \end{bmatrix} = \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0}, \mathbf{s} \geq \mathbf{0}.$$

- If suppose we are given a problem of the type:

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{k \times n} \mathbf{x} = \mathbf{b}_{k \times 1}, \mathbf{x} \geq \mathbf{0}$,

where $k > \text{rank}(\mathbf{A}) = m$.

Basic Feasible Solution

- If $A_{k \times n} \mathbf{x} = \mathbf{b}_{m \times 1}$, $\mathbf{x} \geq \mathbf{0}$ is consistent

Basic Feasible Solution

- If $A_{k \times n} \mathbf{x} = \mathbf{b}_{m \times 1}$, $\mathbf{x} \geq \mathbf{0}$ is consistent then throw away appropriately $k - m$ rows of A and the corresponding components of \mathbf{b} to get an **equivalent** system of equations of the form (*), such that $\text{rank}(A) = m$.

Basic Feasible Solution

- If $A_{k \times n} \mathbf{x} = \mathbf{b}_{m \times 1}$, $\mathbf{x} \geq \mathbf{0}$ is consistent then throw away appropriately $k - m$ rows of A and the corresponding components of \mathbf{b} to get an **equivalent** system of equations of the form (*), such that $\text{rank}(A) = m$.
- An $\mathbf{x} \in \text{Fea}(P)$ is called a **basic feasible solution (BFS)** of the LPP if the **columns** of the matrix A corresponding to the **nonzero components** of \mathbf{x} are **LI**.

Basic Feasible Solution

- If $A_{k \times n} \mathbf{x} = \mathbf{b}_{m \times 1}$, $\mathbf{x} \geq \mathbf{0}$ is consistent then throw away appropriately $k - m$ rows of A and the corresponding components of \mathbf{b} to get an **equivalent** system of equations of the form (*), such that $\text{rank}(A) = m$.
- An $\mathbf{x} \in \text{Fea}(P)$ is called a **basic feasible solution (BFS)** of the LPP if the **columns** of the matrix A corresponding to the **nonzero components** of \mathbf{x} are **LI**.
- An \mathbf{x} satisfying $A\mathbf{x} = \mathbf{b}$, such that the columns corresponding to the **nonzero components** are **LI**, is called a **basic solution** of the LPP.

Basic Feasible Solution

- If $A_{k \times n} \mathbf{x} = \mathbf{b}_{m \times 1}$, $\mathbf{x} \geq \mathbf{0}$ is consistent then throw away appropriately $k - m$ rows of A and the corresponding components of \mathbf{b} to get an **equivalent** system of equations of the form (*), such that $\text{rank}(A) = m$.
- An $\mathbf{x} \in \text{Fea}(P)$ is called a **basic feasible solution (BFS)** of the LPP if the **columns** of the matrix A corresponding to the **nonzero components** of \mathbf{x} are **LI**.
- An \mathbf{x} satisfying $A\mathbf{x} = \mathbf{b}$, such that the columns corresponding to the **nonzero components** are **LI**, is called a **basic solution** of the LPP.
- So a **basic solution** may **not** be a **non negative** vector, hence need not be a **feasible solution** of the LPP.

Basic Feasible Solution

- If $A_{k \times n} \mathbf{x} = \mathbf{b}_{m \times 1}$, $\mathbf{x} \geq \mathbf{0}$ is consistent then throw away appropriately $k - m$ rows of A and the corresponding components of \mathbf{b} to get an **equivalent** system of equations of the form (*), such that $\text{rank}(A) = m$.
- An $\mathbf{x} \in \text{Fea}(P)$ is called a **basic feasible solution (BFS)** of the LPP if the **columns** of the matrix A corresponding to the **nonzero components** of \mathbf{x} are **LI**.
- An \mathbf{x} satisfying $A\mathbf{x} = \mathbf{b}$, such that the columns corresponding to the **nonzero components** are **LI**, is called a **basic solution** of the LPP.
- So a **basic solution** may **not** be a **non negative** vector, hence need not be a **feasible solution** of the LPP.
- A **basic feasible solution** of a LPP of the form (1), can have **at most m** strictly positive components ($\text{rank}(A) = m$).

- **Example 1:** Consider the LPP (P)
Max or Min $\mathbf{c}^T \mathbf{x}$
subject to $\mathbf{A}_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}, \mathbf{x} \geq \mathbf{0}$

- **Example 1:** Consider the LPP (P)

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}, \mathbf{x} \geq \mathbf{0}$

where $\mathbf{A} = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 1 & -1 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$

- **Example 1:** Consider the LPP (P)

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}, \mathbf{x} \geq \mathbf{0}$

where $\mathbf{A} = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 1 & -1 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$

- Clearly $\text{rank}(\mathbf{A}) = 2$.

- **Example 1:** Consider the LPP (P)
 Max or Min $\mathbf{c}^T \mathbf{x}$
 subject to $\mathbf{A}_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}, \mathbf{x} \geq \mathbf{0}$
 where $\mathbf{A} = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 1 & -1 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$
- Clearly $\text{rank}(\mathbf{A}) = 2$.
- $\mathbf{x} = [1, 1, 0]^T$ is a **BFS** of (P) since it satisfies the conditions

- **Example 1:** Consider the LPP (P)

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}, \mathbf{x} \geq \mathbf{0}$

where $\mathbf{A} = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 1 & -1 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$

- Clearly $\text{rank}(\mathbf{A}) = 2$.
- $\mathbf{x} = [1, 1, 0]^T$ is a **BFS** of (P) since it satisfies the conditions $\mathbf{A}_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}, \mathbf{x} \geq \mathbf{0}$

- **Example 1:** Consider the LPP (P)

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}, \mathbf{x} \geq \mathbf{0}$

where $\mathbf{A} = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 1 & -1 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$

- Clearly $\text{rank}(\mathbf{A}) = 2$.
- $\mathbf{x} = [1, 1, 0]^T$ is a **BFS** of (P) since it satisfies the conditions $\mathbf{A}_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}, \mathbf{x} \geq \mathbf{0}$ and the columns of \mathbf{A} corresponding to the **positive components** of \mathbf{x} are

- **Example 1:** Consider the LPP (P)

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}, \mathbf{x} \geq \mathbf{0}$

where $\mathbf{A} = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 1 & -1 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$

- Clearly $\text{rank}(\mathbf{A}) = 2$.
- $\mathbf{x} = [1, 1, 0]^T$ is a **BFS** of (P) since it satisfies the conditions $\mathbf{A}_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}, \mathbf{x} \geq \mathbf{0}$ and the columns of \mathbf{A} corresponding to the **positive components** of \mathbf{x} are $[1, 1]^T, [2, 1]^T$, is **LI**.

- $\mathbf{x} = [2, 1, 1]^T$ is a **feasible** solution of (P) since it satisfies

- $\mathbf{x} = [2, 1, 1]^T$ is a **feasible** solution of (P) since it satisfies $A_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}$, $\mathbf{x} \geq \mathbf{0}$.

- $\mathbf{x} = [2, 1, 1]^T$ is a **feasible** solution of (P) since it satisfies $A_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}$, $\mathbf{x} \geq \mathbf{0}$.
But $\mathbf{x} = [2, 1, 1]^T$ is not a **BFS**

- $\mathbf{x} = [2, 1, 1]^T$ is a **feasible** solution of (P) since it satisfies $A_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}$, $\mathbf{x} \geq \mathbf{0}$.

But $\mathbf{x} = [2, 1, 1]^T$ is not a **BFS**

Since the columns of A corresponding to the **positive components** of \mathbf{x} are

- $\mathbf{x} = [2, 1, 1]^T$ is a **feasible** solution of (P) since it satisfies $A_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}$, $\mathbf{x} \geq \mathbf{0}$.

But $\mathbf{x} = [2, 1, 1]^T$ is not a **BFS**

Since the columns of A corresponding to the **positive components** of \mathbf{x} are

$[1, 1]^T, [2, 1]^T, [-1, -1]^T$ is **LD**.

- $\mathbf{x} = [2, 1, 1]^T$ is a **feasible** solution of (P) since it satisfies $A_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}$, $\mathbf{x} \geq \mathbf{0}$.
But $\mathbf{x} = [2, 1, 1]^T$ is not a **BFS**
Since the columns of A corresponding to the **positive components** of \mathbf{x} are $[1, 1]^T, [2, 1]^T, [-1, -1]^T$ is **LD**.
- $\mathbf{x} = [0, 1, -1]^T$ is a **basic solution** of (P).

- $\mathbf{x} = [2, 1, 1]^T$ is a **feasible** solution of (P) since it satisfies $A_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}$, $\mathbf{x} \geq \mathbf{0}$.

But $\mathbf{x} = [2, 1, 1]^T$ is not a **BFS**

Since the columns of A corresponding to the **positive components** of \mathbf{x} are

$[1, 1]^T, [2, 1]^T, [-1, -1]^T$ is **LD**.

- $\mathbf{x} = [0, 1, -1]^T$ is a **basic solution** of (P).
It satisfies $A_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}$.

- $\mathbf{x} = [2, 1, 1]^T$ is a **feasible** solution of (P) since it satisfies $A_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}$, $\mathbf{x} \geq \mathbf{0}$.

But $\mathbf{x} = [2, 1, 1]^T$ is not a **BFS**

Since the columns of A corresponding to the **positive components** of \mathbf{x} are

$[1, 1]^T, [2, 1]^T, [-1, -1]^T$ is **LD**.

- $\mathbf{x} = [0, 1, -1]^T$ is a **basic solution** of (P).

It satisfies $A_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}$.

The columns of A corresponding to the **nonzero components** of \mathbf{x} are

$[2, 1]^T, [-1, -1]^T$, is **LI**.

- $\mathbf{x} = [2, 1, 1]^T$ is a **feasible** solution of (P) since it satisfies $A_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}$, $\mathbf{x} \geq \mathbf{0}$.

But $\mathbf{x} = [2, 1, 1]^T$ is not a **BFS**

Since the columns of A corresponding to the **positive components** of \mathbf{x} are

$[1, 1]^T, [2, 1]^T, [-1, -1]^T$ is **LD**.

- $\mathbf{x} = [0, 1, -1]^T$ is a **basic solution** of (P).

It satisfies $A_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}$.

The columns of A corresponding to the **nonzero components** of \mathbf{x} are

$[2, 1]^T, [-1, -1]^T$, is **LI**.

But \mathbf{x} is **not** a **BFS**, \mathbf{x} is not a feasible solution of (P).

Non degenerate BFS

- A **basic feasible solution** is called a **non degenerate BFS** if it has exactly m positive components.

Non degenerate BFS

- A **basic feasible solution** is called a **non degenerate BFS** if it has exactly m positive components. Otherwise it is said to be a **degenerate BFS**.

Non degenerate BFS

- A **basic feasible solution** is called a **non degenerate BFS** if it has exactly m positive components. Otherwise it is said to be a **degenerate BFS**.
- If \mathbf{x} is a **non degenerate BFS**, then the columns of A corresponding to the **nonzero** (positive) components of \mathbf{x} form a basis of \mathbb{R}^m .

Non degenerate BFS

- A **basic feasible solution** is called a **non degenerate BFS** if it has exactly m positive components. Otherwise it is said to be a **degenerate BFS**.
- If \mathbf{x} is a **non degenerate BFS**, then the columns of A corresponding to the **nonzero** (positive) components of \mathbf{x} form a basis of \mathbb{R}^m .
- If the positive components of \mathbf{x} are x_1, \dots, x_m ,

Non degenerate BFS

- A **basic feasible solution** is called a **non degenerate BFS** if it has exactly m positive components. Otherwise it is said to be a **degenerate BFS**.
- If \mathbf{x} is a **non degenerate BFS**, then the columns of A corresponding to the **nonzero** (positive) components of \mathbf{x} form a basis of \mathbb{R}^m .
- If the positive components of \mathbf{x} are x_1, \dots, x_m , then $\tilde{\mathbf{a}}_i, i = 1, \dots, m$,

Non degenerate BFS

- A **basic feasible solution** is called a **non degenerate BFS** if it has exactly m positive components. Otherwise it is said to be a **degenerate BFS**.
- If \mathbf{x} is a **non degenerate BFS**, then the columns of A corresponding to the **nonzero** (positive) components of \mathbf{x} form a basis of \mathbb{R}^m .
- If the positive components of \mathbf{x} are x_1, \dots, x_m , then $\tilde{\mathbf{a}}_i, i = 1, \dots, m$, forms a **basis** of \mathbb{R}^m ,

Non degenerate BFS

- A **basic feasible solution** is called a **non degenerate BFS** if it has exactly m positive components. Otherwise it is said to be a **degenerate BFS**.
- If \mathbf{x} is a **non degenerate BFS**, then the columns of A corresponding to the **nonzero** (positive) components of \mathbf{x} form a basis of \mathbb{R}^m .
- If the positive components of \mathbf{x} are x_1, \dots, x_m , then $\tilde{\mathbf{a}}_i, i = 1, \dots, m$, forms a **basis** of \mathbb{R}^m , where $\tilde{\mathbf{a}}_i$ is the i th column of A .

Non degenerate BFS

- A **basic feasible solution** is called a **non degenerate BFS** if it has exactly m positive components. Otherwise it is said to be a **degenerate BFS**.
- If \mathbf{x} is a **non degenerate BFS**, then the columns of A corresponding to the **nonzero** (positive) components of \mathbf{x} form a basis of \mathbb{R}^m .
- If the positive components of \mathbf{x} are x_1, \dots, x_m , then $\tilde{\mathbf{a}}_i, i = 1, \dots, m$, forms a **basis** of \mathbb{R}^m , where $\tilde{\mathbf{a}}_i$ is the i th column of A .
- $B_{m \times m} = [\tilde{\mathbf{a}}_1 \dots \tilde{\mathbf{a}}_m]$ with columns $\tilde{\mathbf{a}}_i, i = 1, \dots, m$, of A is called **the basis matrix** corresponding to \mathbf{x} .

Non degenerate BFS

- A **basic feasible solution** is called a **non degenerate BFS** if it has exactly m positive components. Otherwise it is said to be a **degenerate BFS**.
- If \mathbf{x} is a **non degenerate BFS**, then the columns of A corresponding to the **nonzero** (positive) components of \mathbf{x} form a basis of \mathbb{R}^m .
- If the positive components of \mathbf{x} are x_1, \dots, x_m , then $\tilde{\mathbf{a}}_i, i = 1, \dots, m$, forms a **basis** of \mathbb{R}^m , where $\tilde{\mathbf{a}}_i$ is the i th column of A .
- $B_{m \times m} = [\tilde{\mathbf{a}}_1 \dots \tilde{\mathbf{a}}_m]$ with columns $\tilde{\mathbf{a}}_i, i = 1, \dots, m$, of A is called **the basis matrix** corresponding to \mathbf{x} .
- The variables x_1, \dots, x_m are called **basic variables**, and $x_{m+1} = x_{m+2} = \dots = x_n = 0$, are called **non basic variables** of the **BFS** \mathbf{x} .

Degenerate BFS

- If \mathbf{x} is a **degenerate** BFS, and if the columns of A corresponding to the **nonzero components** of \mathbf{x} are $\tilde{\mathbf{a}}_1, \dots, \tilde{\mathbf{a}}_k$, $k < m$,

Degenerate BFS

- If \mathbf{x} is a **degenerate** BFS, and if the columns of A corresponding to the **nonzero components** of \mathbf{x} are $\tilde{\mathbf{a}}_1, \dots, \tilde{\mathbf{a}}_k$, $k < m$, then consider $(m - k)$ **LI** columns of A , such that these $(m - k)$ columns of A together with $\tilde{\mathbf{a}}_1, \dots, \tilde{\mathbf{a}}_k$, form a **basis** of \mathbb{R}^m .

Degenerate BFS

- If \mathbf{x} is a **degenerate** BFS, and if the columns of A corresponding to the **nonzero components** of \mathbf{x} are $\tilde{\mathbf{a}}_1, \dots, \tilde{\mathbf{a}}_k, k < m$, then consider $(m - k)$ **LI** columns of A , such that these $(m - k)$ columns of A together with $\tilde{\mathbf{a}}_1, \dots, \tilde{\mathbf{a}}_k$, form a **basis** of \mathbb{R}^m .
- $B_{m \times m} = [\tilde{\mathbf{a}}_1 \dots \tilde{\mathbf{a}}_m]$ is called **a** basis matrix corresponding to \mathbf{x} .

Degenerate BFS

- If \mathbf{x} is a **degenerate** BFS, and if the columns of A corresponding to the **nonzero components** of \mathbf{x} are $\tilde{\mathbf{a}}_1, \dots, \tilde{\mathbf{a}}_k$, $k < m$, then consider $(m - k)$ **LI** columns of A , such that these $(m - k)$ columns of A together with $\tilde{\mathbf{a}}_1, \dots, \tilde{\mathbf{a}}_k$, form a **basis** of \mathbb{R}^m .
- $B_{m \times m} = [\tilde{\mathbf{a}}_1 \dots \tilde{\mathbf{a}}_m]$ is called **a** basis matrix corresponding to \mathbf{x} .
- The components x_1, \dots, x_m of \mathbf{x} are called **basic variables** corresponding to \mathbf{x} and the basis matrix B .

Degenerate BFS

- If \mathbf{x} is a **degenerate** BFS, and if the columns of A corresponding to the **nonzero components** of \mathbf{x} are $\tilde{\mathbf{a}}_1, \dots, \tilde{\mathbf{a}}_k, k < m$, then consider $(m - k)$ **LI** columns of A , such that these $(m - k)$ columns of A together with $\tilde{\mathbf{a}}_1, \dots, \tilde{\mathbf{a}}_k$, form a **basis** of \mathbb{R}^m .
- $B_{m \times m} = [\tilde{\mathbf{a}}_1 \dots \tilde{\mathbf{a}}_m]$ is called **a** basis matrix corresponding to \mathbf{x} .
- The components x_1, \dots, x_m of \mathbf{x} are called **basic variables** corresponding to \mathbf{x} and the basis matrix B .
- The components $x_{m+1} = \dots = x_n = 0$ are **nonbasic variables** corresponding to \mathbf{x} and the basis matrix B .

- **Example 2:** Consider the LPP (P)

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}, \mathbf{x} \geq \mathbf{0}$

- **Example 2:** Consider the LPP (P)

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}, \mathbf{x} \geq \mathbf{0}$

where $\mathbf{A} = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 1 & 0 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

- **Example 2:** Consider the LPP (P)

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}, \mathbf{x} \geq \mathbf{0}$

where $\mathbf{A} = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 1 & 0 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

- Clearly $\text{rank}(\mathbf{A}) = 2$.

- **Example 2:** Consider the LPP (P)

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}, \mathbf{x} \geq \mathbf{0}$

where $\mathbf{A} = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 1 & 0 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

- Clearly $\text{rank}(\mathbf{A}) = 2$.
- $\mathbf{x} = [0, 1, 1]^T$ is a **non degenerate BFS** of (P) since it satisfies the conditions

- **Example 2:** Consider the LPP (P)

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}, \mathbf{x} \geq \mathbf{0}$

where $\mathbf{A} = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 1 & 0 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

- Clearly $\text{rank}(\mathbf{A}) = 2$.
- $\mathbf{x} = [0, 1, 1]^T$ is a **non degenerate BFS** of (P) since it satisfies the conditions
 $\mathbf{A}_{m \times n} \mathbf{x} = \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0}$

- **Example 2:** Consider the LPP (P)

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}, \mathbf{x} \geq \mathbf{0}$

where $\mathbf{A} = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 1 & 0 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

- Clearly $\text{rank}(\mathbf{A}) = 2$.
- $\mathbf{x} = [0, 1, 1]^T$ is a **non degenerate BFS** of (P) since it satisfies the conditions
 $\mathbf{A}_{m \times n} \mathbf{x} = \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0}$
and the columns of \mathbf{A} corresponding to the positive components of \mathbf{x} are

- **Example 2:** Consider the LPP (P)

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}, \mathbf{x} \geq \mathbf{0}$

where $\mathbf{A} = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 1 & 0 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

- Clearly $\text{rank}(\mathbf{A}) = 2$.
- $\mathbf{x} = [0, 1, 1]^T$ is a **non degenerate BFS** of (P) since it satisfies the conditions

$$\mathbf{A}_{m \times n} \mathbf{x} = \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0}$$

and the columns of \mathbf{A} corresponding to the positive components of \mathbf{x} are

$[2, 1]^T, [-1, 0]^T$, is LI.

- **Example 2:** Consider the LPP (P)

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}, \mathbf{x} \geq \mathbf{0}$

where $\mathbf{A} = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 1 & 0 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

- Clearly $\text{rank}(\mathbf{A}) = 2$.
- $\mathbf{x} = [0, 1, 1]^T$ is a **non degenerate BFS** of (P) since it satisfies the conditions
 $\mathbf{A}_{m \times n} \mathbf{x} = \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0}$
and the columns of \mathbf{A} corresponding to the positive components of \mathbf{x} are
 $[2, 1]^T, [-1, 0]^T$, is LI.
Also the number of **positive components** of \mathbf{x} is equal to
 $2 = \text{rank}(\mathbf{A})$.

- **Example 2:** Consider the LPP (P)

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}, \mathbf{x} \geq 0$

where $\mathbf{A} = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 1 & 0 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

- Clearly $\text{rank}(\mathbf{A}) = 2$.
- $\mathbf{x} = [0, 1, 1]^T$ is a **non degenerate BFS** of (P) since it satisfies the conditions

$$\mathbf{A}_{m \times n} \mathbf{x} = \mathbf{b}_{m \times 1}, \mathbf{x} \geq 0$$

and the columns of \mathbf{A} corresponding to the positive components of \mathbf{x} are

$[2, 1]^T, [-1, 0]^T$, is LI.

Also the number of **positive components** of \mathbf{x} is equal to $2 = \text{rank}(\mathbf{A})$.

- The **basis matrix** for this \mathbf{x} is unique upto a permutation of the columns.

- **Example 2:** Consider the LPP (P)

Max or Min $\mathbf{c}^T \mathbf{x}$

subject to $\mathbf{A}_{2 \times 3} \mathbf{x} = \mathbf{b}_{2 \times 1}, \mathbf{x} \geq 0$

where $\mathbf{A} = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 1 & 0 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

- Clearly $\text{rank}(\mathbf{A}) = 2$.
- $\mathbf{x} = [0, 1, 1]^T$ is a **non degenerate BFS** of (P) since it satisfies the conditions

$$\mathbf{A}_{m \times n} \mathbf{x} = \mathbf{b}_{m \times 1}, \mathbf{x} \geq 0$$

and the columns of \mathbf{A} corresponding to the positive components of \mathbf{x} are

$[2, 1]^T, [-1, 0]^T$, is LI.

Also the number of **positive components** of \mathbf{x} is equal to $2 = \text{rank}(\mathbf{A})$.

- The **basis matrix** for this \mathbf{x} is unique upto a permutation of the columns.

x_2, x_3 are basic variables and x_1 is non basic.

Example 2 (continued)

- $\mathbf{x} = [1, 0, 0]^T$ is also a BFS of (P) since it satisfies

Example 2 (continued)

- $\mathbf{x} = [1, 0, 0]^T$ is also a BFS of (P) since it satisfies
 $A_{2 \times 3} \mathbf{x} = \mathbf{b}_{3 \times 1}$, $\mathbf{x} \geq \mathbf{0}$,

Example 2 (continued)

- $\mathbf{x} = [1, 0, 0]^T$ is also a BFS of (P) since it satisfies $A_{2 \times 3} \mathbf{x} = \mathbf{b}_{3 \times 1}$, $\mathbf{x} \geq \mathbf{0}$,
the columns of A corresponding to the **positive components** of \mathbf{x} is

Example 2 (continued)

- $\mathbf{x} = [1, 0, 0]^T$ is also a BFS of (P) since it satisfies $A_{2 \times 3} \mathbf{x} = \mathbf{b}_{3 \times 1}$, $\mathbf{x} \geq \mathbf{0}$,
the columns of A corresponding to the positive components of \mathbf{x} is $[1, 1]^T$, is LI.

Example 2 (continued)

- $\mathbf{x} = [1, 0, 0]^T$ is also a BFS of (P) since it satisfies $A_{2 \times 3} \mathbf{x} = \mathbf{b}_{3 \times 1}$, $\mathbf{x} \geq \mathbf{0}$,
the columns of A corresponding to the positive components of \mathbf{x} is $[1, 1]^T$, is LI.
- \mathbf{x} is a degenerate BFS since the number of positive components of \mathbf{x} is strictly less 2.

Example 2 (continued)

- $\mathbf{x} = [1, 0, 0]^T$ is also a BFS of (P) since it satisfies $A_{2 \times 3} \mathbf{x} = \mathbf{b}_{3 \times 1}$, $\mathbf{x} \geq \mathbf{0}$,
the columns of A corresponding to the **positive components** of \mathbf{x} is $[1, 1]^T$, is **LI**.
- \mathbf{x} is a **degenerate** BFS since the number of **positive components** of \mathbf{x} is strictly less 2.
- There are **two** different basis matrices B for \mathbf{x} .
(i) B with columns $[1, 1]^T, [2, 1]^T$, **basic variables** x_1, x_2 .

Example 2 (continued)

- $\mathbf{x} = [1, 0, 0]^T$ is also a BFS of (P) since it satisfies $A_{2 \times 3}\mathbf{x} = \mathbf{b}_{3 \times 1}$, $\mathbf{x} \geq \mathbf{0}$,
the columns of A corresponding to the **positive components** of \mathbf{x} is $[1, 1]^T$, is **LI**.
- \mathbf{x} is a **degenerate** BFS since the number of **positive components** of \mathbf{x} is strictly less 2.
- There are **two** different basis matrices B for \mathbf{x} .
 - (i) B with columns $[1, 1]^T, [2, 1]^T$, **basic variables** x_1, x_2 .
 - (ii) B' with columns $[1, 1]^T, [-1, 0]^T$, **basic variables** x_1, x_3 .

- Depending on the choice of the $(m - k)$ columns of A which are added, the same **degenerate BFS** \mathbf{x} will correspond to **different basis matrices** B ,

- Depending on the choice of the $(m - k)$ columns of A which are added, the same **degenerate BFS** \mathbf{x} will correspond to **different basis matrices** B , and will have different **basic** and **nonbasic** variables.

- Depending on the choice of the $(m - k)$ columns of A which are added, the same **degenerate BFS** \mathbf{x} will correspond to **different basis matrices** B , and will have different **basic** and **nonbasic** variables.
- If \mathbf{x} is a **BFS** then $\mathbf{x}_B = B^{-1}\mathbf{b}$,

- Depending on the choice of the $(m - k)$ columns of A which are added, the same **degenerate BFS** \mathbf{x} will correspond to **different basis matrices** B , and will have different **basic** and **nonbasic** variables.
- If \mathbf{x} is a **BFS** then $\mathbf{x}_B = B^{-1}\mathbf{b}$, where \mathbf{x}_B are the components of \mathbf{x} corresponding to the **basic variables**.

- Depending on the choice of the $(m - k)$ columns of A which are added, the same **degenerate BFS** \mathbf{x} will correspond to **different basis matrices** B , and will have different **basic** and **nonbasic** variables.
- If \mathbf{x} is a **BFS** then $\mathbf{x}_B = B^{-1}\mathbf{b}$, where \mathbf{x}_B are the components of \mathbf{x} corresponding to the **basic variables**.
- **BFS** \mathbf{x} is of the form, $\mathbf{x} = \begin{bmatrix} B^{-1}\mathbf{b}_{m \times 1} \\ \mathbf{0}_{(n-m) \times 1} \end{bmatrix}$.

- Depending on the choice of the $(m - k)$ columns of A which are added, the same **degenerate BFS** \mathbf{x} will correspond to **different basis matrices** B , and will have different **basic** and **nonbasic** variables.
- If \mathbf{x} is a **BFS** then $\mathbf{x}_B = B^{-1}\mathbf{b}$, where \mathbf{x}_B are the components of \mathbf{x} corresponding to the **basic variables**.
- **BFS** \mathbf{x} is of the form, $\mathbf{x} = \begin{bmatrix} B^{-1}\mathbf{b}_{m \times 1} \\ \mathbf{0}_{(n-m) \times 1} \end{bmatrix}$.
- Corresponding to **BFS** \mathbf{x}

- Depending on the choice of the $(m - k)$ columns of A which are added, the same **degenerate BFS** \mathbf{x} will correspond to **different basis matrices** B , and will have different **basic** and **nonbasic** variables.
- If \mathbf{x} is a **BFS** then $\mathbf{x}_B = B^{-1}\mathbf{b}$,
where \mathbf{x}_B are the components of \mathbf{x} corresponding to the **basic variables**.
- **BFS** \mathbf{x} is of the form, $\mathbf{x} = \begin{bmatrix} B^{-1}\mathbf{b}_{m \times 1} \\ \mathbf{0}_{(n-m) \times 1} \end{bmatrix}$.
- Corresponding to **BFS** \mathbf{x}
 $\mathbf{c}^T \mathbf{x} = \mathbf{c}_B^T \mathbf{x}_B = \mathbf{c}_B^T B^{-1} \mathbf{b}$,

- Depending on the choice of the $(m - k)$ columns of A which are added, the same **degenerate BFS** \mathbf{x} will correspond to **different basis matrices** B , and will have different **basic** and **nonbasic** variables.
- If \mathbf{x} is a **BFS** then $\mathbf{x}_B = B^{-1}\mathbf{b}$, where \mathbf{x}_B are the components of \mathbf{x} corresponding to the **basic variables**.
- BFS** \mathbf{x} is of the form, $\mathbf{x} = \begin{bmatrix} B^{-1}\mathbf{b}_{m \times 1} \\ \mathbf{0}_{(n-m) \times 1} \end{bmatrix}$.
- Corresponding to **BFS** \mathbf{x}
 $\mathbf{c}^T \mathbf{x} = \mathbf{c}_B^T \mathbf{x}_B = \mathbf{c}_B^T B^{-1} \mathbf{b}$,
 where, \mathbf{c}_B^T are the components of \mathbf{c}^T corresponding to the basic variables.

- Depending on the choice of the $(m - k)$ columns of A which are added, the same **degenerate BFS** \mathbf{x} will correspond to **different basis matrices** B , and will have different **basic** and **nonbasic** variables.
- If \mathbf{x} is a **BFS** then $\mathbf{x}_B = B^{-1}\mathbf{b}$,
where \mathbf{x}_B are the components of \mathbf{x} corresponding to the **basic variables**.
- **BFS** \mathbf{x} is of the form, $\mathbf{x} = \begin{bmatrix} B^{-1}\mathbf{b}_{m \times 1} \\ \mathbf{0}_{(n-m) \times 1} \end{bmatrix}$.
- Corresponding to **BFS** \mathbf{x}
 $\mathbf{c}^T \mathbf{x} = \mathbf{c}_B^T \mathbf{x}_B = \mathbf{c}_B^T B^{-1} \mathbf{b}$,
where, \mathbf{c}_B^T are the components of \mathbf{c}^T corresponding to the basic variables.
- If LPP (P) is the **diet problem**, (with \geq inequalities changed to equalities in the original problem),
then \mathbf{x} gives the **quantities** of the food products F_j ,
 $j = 1, \dots, n$ in the diet.

- The food products F_1, \dots, F_m , which correspond to the basic variables x_1, x_2, \dots, x_m of \mathbf{x} , are the ones which are included in the diet in the quantities x_j .

- The food products F_1, \dots, F_m , which correspond to the **basic variables** x_1, x_2, \dots, x_m of \mathbf{x} , are the ones which are **included** in the diet in the quantities x_j .
- The other food products corresponding to the **nonbasic variables** of \mathbf{x} (they have zero values) are **not** consumed.

- The food products F_1, \dots, F_m , which correspond to the **basic variables** x_1, x_2, \dots, x_m of \mathbf{x} , are the ones which are **included** in the diet in the quantities x_j .
- The other food products corresponding to the **nonbasic variables** of \mathbf{x} (they have zero values) are **not** consumed.
- Let us assume that (P^*) has **atleast one feasible solution** and let \mathbf{x} be a BFS (degenerate or non degenerate) of the LPP (P^*) .

- The food products F_1, \dots, F_m , which correspond to the **basic variables** x_1, x_2, \dots, x_m of \mathbf{x} , are the ones which are **included** in the diet in the quantities x_j .
- The other food products corresponding to the **nonbasic variables** of \mathbf{x} (they have zero values) are **not** consumed.
- Let us assume that (P^*) has **atleast one feasible solution** and let \mathbf{x} be a BFS (degenerate or non degenerate) of the LPP (P^*) .
- Let $B = [\tilde{\mathbf{a}}_1 \dots \tilde{\mathbf{a}}_m]$ be **a** (or **the**) **basis matrix** corresponding to \mathbf{x} .

- The food products F_1, \dots, F_m , which correspond to the **basic variables** x_1, x_2, \dots, x_m of \mathbf{x} , are the ones which are **included** in the diet in the quantities x_j .
- The other food products corresponding to the **nonbasic variables** of \mathbf{x} (they have zero values) are **not** consumed.
- Let us assume that (P^*) has **atleast one feasible solution** and let \mathbf{x} be a BFS (degenerate or non degenerate) of the LPP (P^*) .
- Let $B = [\tilde{\mathbf{a}}_1 \dots \tilde{\mathbf{a}}_m]$ be **a (or the) basis matrix** corresponding to \mathbf{x} .
- For all $k = 1, 2, \dots, n$, let

$$\tilde{\mathbf{a}}_k = \sum_{i=1}^m u_{ik} \tilde{\mathbf{a}}_i = [\tilde{\mathbf{a}}_1 \dots \tilde{\mathbf{a}}_m] \begin{bmatrix} u_{1k} \\ \vdots \\ u_{mk} \end{bmatrix} \quad (*)$$

for some real u_{ik} 's, $i = 1, \dots, m$, $k = 1, \dots, n$,

- The food products F_1, \dots, F_m , which correspond to the **basic variables** x_1, x_2, \dots, x_m of \mathbf{x} , are the ones which are **included** in the diet in the quantities x_j .
- The other food products corresponding to the **nonbasic variables** of \mathbf{x} (they have zero values) are **not** consumed.
- Let us assume that (P^*) has **atleast one feasible solution** and let \mathbf{x} be a BFS (degenerate or non degenerate) of the LPP (P^*) .
- Let $B = [\tilde{\mathbf{a}}_1 \dots \tilde{\mathbf{a}}_m]$ be **a (or the) basis matrix** corresponding to \mathbf{x} .
- For all $k = 1, 2, \dots, n$, let

$$\tilde{\mathbf{a}}_k = \sum_{i=1}^m u_{ik} \tilde{\mathbf{a}}_i = [\tilde{\mathbf{a}}_1 \dots \tilde{\mathbf{a}}_m] \begin{bmatrix} u_{1k} \\ \vdots \\ u_{mk} \end{bmatrix} \quad (*)$$

for some real u_{ik} 's, $i = 1, \dots, m$, $k = 1, \dots, n$,

- Then $\begin{bmatrix} u_{1k} \\ \vdots \\ u_{mk} \end{bmatrix} = B^{-1} \tilde{\mathbf{a}}_k$ for all $k = 1, 2, \dots, n$.

- In order to obtain the same amount of nutrient as unit amount of F_k , $k = 1, \dots, n$, one needs to consume (u_{1k} amount of F_1) + (u_{2k} amount of F_2) + \dots + (u_{mk} amount of F_m).

- In order to obtain the same amount of nutrient as unit amount of F_k , $k = 1, \dots, n$, one needs to consume (u_{1k} amount of F_1) + (u_{2k} amount of F_2) + \dots + (u_{mk} amount of F_m).
- Let z_k be the value of unit amount of F_k , if we include **only** $F_i, i = 1, \dots, m$ in the diet.

- In order to obtain the same amount of nutrient as unit amount of F_k , $k = 1, \dots, n$, one needs to consume (u_{1k} amount of F_1) + (u_{2k} amount of F_2) + ... + (u_{mk} amount of F_m).
- Let z_k be the value of unit amount of F_k , if we include **only** F_i , $i = 1, \dots, m$ in the diet.
- Then $z_k = u_{1k}c_1 + u_{2k}c_2 + \dots + u_{mk}c_m = \sum_{i=1}^m u_{ik}c_i$

$$= [c_1, \dots, c_m] \begin{bmatrix} u_{1k} \\ \vdots \\ u_{mk} \end{bmatrix} = \mathbf{c}_B^T B^{-1} \tilde{\mathbf{a}}_k.$$

- In order to obtain the same amount of nutrient as unit amount of F_k , $k = 1, \dots, n$, one needs to consume (u_{1k} amount of F_1) + (u_{2k} amount of F_2) + ... + (u_{mk} amount of F_m).
- Let z_k be the value of unit amount of F_k , if we include **only** F_i , $i = 1, \dots, m$ in the diet.
- Then $z_k = u_{1k}c_1 + u_{2k}c_2 + \dots + u_{mk}c_m = \sum_{i=1}^m u_{ik}c_i$

$$= [c_1, \dots, c_m] \begin{bmatrix} u_{1k} \\ \vdots \\ u_{mk} \end{bmatrix} = \mathbf{c}_B^T B^{-1} \tilde{\mathbf{a}}_k.$$
- For all $k = 1, \dots, m$, note that $z_k = c_k$.

- In order to obtain the same amount of nutrient as unit amount of F_k , $k = 1, \dots, n$, one needs to consume (u_{1k} amount of F_1) + (u_{2k} amount of F_2) + ... + (u_{mk} amount of F_m).
- Let z_k be the value of unit amount of F_k , if we include **only** F_i , $i = 1, \dots, m$ in the diet.
- Then $z_k = u_{1k}c_1 + u_{2k}c_2 + \dots + u_{mk}c_m = \sum_{i=1}^m u_{ik}c_i$

$$= [c_1, \dots, c_m] \begin{bmatrix} u_{1k} \\ \vdots \\ u_{mk} \end{bmatrix} = \mathbf{c}_B^T B^{-1} \tilde{\mathbf{a}}_k.$$
- For all $k = 1, \dots, m$, note that $z_k = c_k$.
- If $z_k > c_k$ then look for a better solution.

- the simplex table corresponding to BFS \mathbf{x} is given by

	0	..	0	..	$c_s - z_s$..	$c_k - z_k$..	
	$B^{-1}\tilde{\mathbf{a}}_1$..	$B^{-1}\tilde{\mathbf{a}}_m$..	$B^{-1}\tilde{\mathbf{a}}_s$..	$B^{-1}\tilde{\mathbf{a}}_k$..	$B^{-1}\mathbf{b}$
$\tilde{\mathbf{a}}_1$	1	..	0	..	u_{1s}	..	u_{1k}	..	x_1
$\tilde{\mathbf{a}}_2$	0	..	0	..	u_{2s}	..	u_{2k}	..	x_2
\vdots	0	..	0	..	\vdots	..	\vdots	..	\vdots
$\tilde{\mathbf{a}}_r$	\vdots	..	\vdots	..	u_{rs}	..	u_{rk}	..	x_r
\vdots	\vdots	..	\vdots	..	\vdots	..	\vdots	..	\vdots
$\tilde{\mathbf{a}}_m$	0	..	1	..	u_{ms}	..	u_{mk}	..	x_m

- **Case 1:** $c_k - z_k < 0$ for atleast one k , $k = m + 1, \dots, n$.

- **Case 1:** $c_k - z_k < 0$ for atleast one k , $k = m + 1, \dots, n$.

Let $c_s - z_s = \min\{c_k - z_k : c_k - z_k < 0, k = m + 1, \dots, n\}$.

- **Case 1:** $c_k - z_k < 0$ for atleast one k , $k = m + 1, \dots, n$.
Let $c_s - z_s = \min\{c_k - z_k : c_k - z_k < 0, k = m + 1, \dots, n\}$.
Simplex algorithm says that, include x_s in the diet or
make x_s a **basic variable**.

- **Case 1:** $c_k - z_k < 0$ for atleast one k , $k = m + 1, \dots, n$.
Let $c_s - z_s = \min\{c_k - z_k : c_k - z_k < 0, k = m + 1, \dots, n\}$.
Simplex algorithm says that, include x_s in the diet or
make x_s a **basic variable**.
Column \tilde{a}_s will be included in the basis matrix.

- Case 1:** $c_k - z_k < 0$ for atleast one k , $k = m + 1, \dots, n$.
 Let $c_s - z_s = \min\{c_k - z_k : c_k - z_k < 0, k = m + 1, \dots, n\}$.
Simplex algorithm says that, include x_s in the diet or make x_s a **basic variable**.
 Column \tilde{a}_s will be included in the basis matrix.
- If there exists, s, l , such that for both s, l ,
 $c_s - z_s = c_l - z_l = \min\{c_k - z_k : c_k - z_k < 0, k = m + 1, \dots, n\}$,

- **Case 1:** $c_k - z_k < 0$ for atleast one k , $k = m + 1, \dots, n$.
 Let $c_s - z_s = \min\{c_k - z_k : c_k - z_k < 0, k = m + 1, \dots, n\}$.
Simplex algorithm says that, include x_s in the diet or make x_s a **basic variable**.
 Column \tilde{a}_s will be included in the basis matrix.
- If there exists, s, l , such that for both s, l ,
 $c_s - z_s = c_l - z_l = \min\{c_k - z_k : c_k - z_k < 0, k = m + 1, \dots, n\}$,
 then include any **one** of these food products in the diet(as a basic variable).

- **Case 1:** $c_k - z_k < 0$ for atleast one k , $k = m + 1, \dots, n$.
 Let $c_s - z_s = \min\{c_k - z_k : c_k - z_k < 0, k = m + 1, \dots, n\}$.
Simplex algorithm says that, include x_s in the diet or make x_s a **basic variable**.
 Column \tilde{a}_s will be included in the basis matrix.
- If there exists, s, l , such that for both s, l ,
 $c_s - z_s = c_l - z_l = \min\{c_k - z_k : c_k - z_k < 0, k = m + 1, \dots, n\}$,
 then include any **one** of these food products in the diet(as a basic variable).
- Let \mathbf{x}' be the new feasible solution given by
 $x'_i = x_i - u_{is}x'_s$ for $i = 1, \dots, m$,

- **Case 1:** $c_k - z_k < 0$ for atleast one k , $k = m + 1, \dots, n$.
 Let $c_s - z_s = \min\{c_k - z_k : c_k - z_k < 0, k = m + 1, \dots, n\}$.
Simplex algorithm says that, include x_s in the diet or make x_s a **basic variable**.
 Column \tilde{a}_s will be included in the basis matrix.
- If there exists, s, l , such that for both s, l ,
 $c_s - z_s = c_l - z_l = \min\{c_k - z_k : c_k - z_k < 0, k = m + 1, \dots, n\}$,
 then include any **one** of these food products in the diet(as a basic variable).
- Let \mathbf{x}' be the new feasible solution given by
 $x'_i = x_i - u_{is}x'_s$ for $i = 1, \dots, m$,
 $x'_s \geq 0$ and
 $x'_i = 0$ for $i = m + 1, \dots, n, i \neq s$.

- If \mathbf{x}' is feasible for the LPP, then $x'_s \geq 0$ is such that $x'_i \geq 0$ for all $i = 1, \dots, m$.

- If \mathbf{x}' is feasible for the LPP, then $x'_s \geq 0$ is such that $x'_i \geq 0$ for all $i = 1, \dots, m$.
- If $u_{is} \leq 0$, for some $i = 1, \dots, m$,

- If \mathbf{x}' is feasible for the LPP, then $x'_s \geq 0$ is such that $x'_i \geq 0$ for all $i = 1, \dots, m$.
- If $u_{is} \leq 0$, for some $i = 1, \dots, m$, then for all $x'_s \geq 0$, $x'_i \geq 0$ for that i .

- If \mathbf{x}' is feasible for the LPP, then $x'_s \geq 0$ is such that $x'_i \geq 0$ for all $i = 1, \dots, m$.
- If $u_{is} \leq 0$, for some $i = 1, \dots, m$, then for all $x'_s \geq 0$, $x'_i \geq 0$ for that i .
- If $u_{is} > 0$ for some $i = 1, \dots, m$,

- If \mathbf{x}' is feasible for the LPP, then $x'_s \geq 0$ is such that $x'_i \geq 0$ for all $i = 1, \dots, m$.
- If $u_{is} \leq 0$, for some $i = 1, \dots, m$, then for all $x'_s \geq 0$, $x'_i \geq 0$ for that i .
- If $u_{is} > 0$ for some $i = 1, \dots, m$, $x'_s \geq 0$, should be such that $x'_s \leq \frac{x_i}{u_{is}}$.

- If \mathbf{x}' is feasible for the LPP, then $x'_s \geq 0$ is such that $x'_i \geq 0$ for all $i = 1, \dots, m$.
- If $u_{is} \leq 0$, for some $i = 1, \dots, m$, then for all $x'_s \geq 0$, $x'_i \geq 0$ for that i .
- If $u_{is} > 0$ for some $i = 1, \dots, m$, $x'_s \geq 0$, should be such that $x'_s \leq \frac{x_i}{u_{is}}$.
- **Case 1a:** For atleast one $i = 1, \dots, m$, $u_{is} > 0$ (where s is as defined in **Case 1**).

- If \mathbf{x}' is feasible for the LPP, then $x'_s \geq 0$ is such that $x'_i \geq 0$ for all $i = 1, \dots, m$.
- If $u_{is} \leq 0$, for some $i = 1, \dots, m$, then for all $x'_s \geq 0$, $x'_i \geq 0$ for that i .
- If $u_{is} > 0$ for some $i = 1, \dots, m$, $x'_s \geq 0$, should be such that $x'_s \leq \frac{x_i}{u_{is}}$.
- **Case 1a:** For atleast one $i = 1, \dots, m$, $u_{is} > 0$ (where s is as defined in **Case 1**).
Let $\frac{x_r}{u_{rs}} = \min\{\frac{x_i}{u_{is}} : u_{is} > 0\}$.

- If \mathbf{x}' is feasible for the LPP, then $x'_s \geq 0$ is such that $x'_i \geq 0$ for all $i = 1, \dots, m$.
- If $u_{is} \leq 0$, for some $i = 1, \dots, m$, then for all $x'_s \geq 0$, $x'_i \geq 0$ for that i .
- If $u_{is} > 0$ for some $i = 1, \dots, m$, $x'_s \geq 0$, should be such that $x'_s \leq \frac{x_i}{u_{is}}$.
- **Case 1a:** For atleast one $i = 1, \dots, m$, $u_{is} > 0$ (where s is as defined in **Case 1**).
Let $\frac{x_r}{u_{rs}} = \min\{\frac{x_i}{u_{is}} : u_{is} > 0\}$.
- $\frac{x_r}{u_{rs}}$ is called the **minimum ratio**.

- If \mathbf{x}' is feasible for the LPP, then $x'_s \geq 0$ is such that $x'_i \geq 0$ for all $i = 1, \dots, m$.
- If $u_{is} \leq 0$, for some $i = 1, \dots, m$, then for all $x'_s \geq 0$, $x'_i \geq 0$ for that i .
- If $u_{is} > 0$ for some $i = 1, \dots, m$, $x'_s \geq 0$, should be such that $x'_s \leq \frac{x_i}{u_{is}}$.
- **Case 1a:** For atleast one $i = 1, \dots, m$, $u_{is} > 0$ (where s is as defined in **Case 1**).
Let $\frac{x_r}{u_{rs}} = \min\{\frac{x_i}{u_{is}} : u_{is} > 0\}$.
- $\frac{x_r}{u_{rs}}$ is called the **minimum ratio**.
- $\mathbf{x}' \geq 0 \Rightarrow \mathbf{x}'_s \leq \frac{x_r}{u_{rs}}$.

- If \mathbf{x}' is feasible for the LPP, then $x'_s \geq 0$ is such that $x'_i \geq 0$ for all $i = 1, \dots, m$.
- If $u_{is} \leq 0$, for some $i = 1, \dots, m$, then for all $x'_s \geq 0$, $x'_i \geq 0$ for that i .
- If $u_{is} > 0$ for some $i = 1, \dots, m$, $x'_s \geq 0$, should be such that $x'_s \leq \frac{x_i}{u_{is}}$.
- **Case 1a:** For atleast one $i = 1, \dots, m$, $u_{is} > 0$ (where s is as defined in **Case 1**).
Let $\frac{x_r}{u_{rs}} = \min\{\frac{x_i}{u_{is}} : u_{is} > 0\}$.
- $\frac{x_r}{u_{rs}}$ is called the **minimum ratio**.
- $\mathbf{x}' \geq \mathbf{0} \Rightarrow \mathbf{x}'_s \leq \frac{x_r}{u_{rs}}$.
- Also $A\mathbf{x}' = \mathbf{b}$ which implies $\mathbf{x}' \in \text{Fea}(LPP)$.

- $\mathbf{c}^T \mathbf{x}' = \sum_{i=1}^m c_i x'_i + c_s x'_s$

- $$\begin{aligned}\mathbf{c}^T \mathbf{x}' &= \sum_{i=1}^m c_i x'_i + c_s x'_s \\ &= \mathbf{c}^T \mathbf{x} + x'_s (c_s - z_s),\end{aligned}$$

- $$\begin{aligned}
 \bullet \quad \mathbf{c}^T \mathbf{x}' &= \sum_{i=1}^m c_i x'_i + c_s x'_s \\
 &= \mathbf{c}^T \mathbf{x} + x'_s (c_s - z_s), \\
 &\Rightarrow \mathbf{c}^T \mathbf{x}' \leq \mathbf{c}^T \mathbf{x}.
 \end{aligned}$$

- $\mathbf{c}^T \mathbf{x}' = \sum_{i=1}^m c_i x'_i + c_s x'_s$
 $= \mathbf{c}^T \mathbf{x} + x'_s (c_s - z_s),$
 $\Rightarrow \mathbf{c}^T \mathbf{x}' \leq \mathbf{c}^T \mathbf{x}.$
- $\mathbf{c}^T \mathbf{x}' < \mathbf{c}^T \mathbf{x}$ if $x'_s > 0,$

- $\bullet \mathbf{c}^T \mathbf{x}' = \sum_{i=1}^m c_i x'_i + c_s x'_s$
 $= \mathbf{c}^T \mathbf{x} + x'_s (c_s - z_s),$
 $\Rightarrow \mathbf{c}^T \mathbf{x}' \leq \mathbf{c}^T \mathbf{x}.$
- $\bullet \mathbf{c}^T \mathbf{x}' < \mathbf{c}^T \mathbf{x}$ if $x'_s > 0$, which is when the **minimum ratio**
 $\frac{x_r}{u_{rs}} > 0.$

- $\bullet \mathbf{c}^T \mathbf{x}' = \sum_{i=1}^m c_i x'_i + c_s x'_s$
 $= \mathbf{c}^T \mathbf{x} + x'_s (c_s - z_s),$
 $\Rightarrow \mathbf{c}^T \mathbf{x}' \leq \mathbf{c}^T \mathbf{x}.$
- $\bullet \mathbf{c}^T \mathbf{x}' < \mathbf{c}^T \mathbf{x}$ if $x'_s > 0$, which is when the **minimum ratio**
 $\frac{x_r}{u_{rs}} > 0.$
- \bullet Let $x'_s = \frac{x_r}{u_{rs}}.$

- $\bullet \mathbf{c}^T \mathbf{x}' = \sum_{i=1}^m c_i x'_i + c_s x'_s$
 $= \mathbf{c}^T \mathbf{x} + x'_s (c_s - z_s),$
 $\Rightarrow \mathbf{c}^T \mathbf{x}' \leq \mathbf{c}^T \mathbf{x}.$
- $\bullet \mathbf{c}^T \mathbf{x}' < \mathbf{c}^T \mathbf{x}$ if $x'_s > 0$, which is when the **minimum ratio**
 $\frac{x_r}{u_{rs}} > 0.$
- \bullet Let $x'_s = \frac{x_r}{u_{rs}}.$
- \bullet Then $x'_r = x_r - u_{rs} \frac{x_r}{u_{rs}} = 0$

- $\bullet \mathbf{c}^T \mathbf{x}' = \sum_{i=1}^m c_i x'_i + c_s x'_s$
 $= \mathbf{c}^T \mathbf{x} + x'_s (c_s - z_s),$
 $\Rightarrow \mathbf{c}^T \mathbf{x}' \leq \mathbf{c}^T \mathbf{x}.$
- $\bullet \mathbf{c}^T \mathbf{x}' < \mathbf{c}^T \mathbf{x}$ if $x'_s > 0$, which is when the **minimum ratio**
 $\frac{x_r}{u_{rs}} > 0.$
- \bullet Let $x'_s = \frac{x_r}{u_{rs}}.$
- \bullet Then $x'_r = x_r - u_{rs} \frac{x_r}{u_{rs}} = 0$
- $\bullet x_s$ is called the **entering variable**,

- $\bullet \mathbf{c}^T \mathbf{x}' = \sum_{i=1}^m c_i x'_i + c_s x'_s$
 $= \mathbf{c}^T \mathbf{x} + x'_s (c_s - z_s),$
 $\Rightarrow \mathbf{c}^T \mathbf{x}' \leq \mathbf{c}^T \mathbf{x}.$
- $\bullet \mathbf{c}^T \mathbf{x}' < \mathbf{c}^T \mathbf{x}$ if $x'_s > 0$, which is when the **minimum ratio**
 $\frac{x_r}{u_{rs}} > 0.$
- \bullet Let $x'_s = \frac{x_r}{u_{rs}}.$
- \bullet Then $x'_r = x_r - u_{rs} \frac{x_r}{u_{rs}} = 0$
- $\bullet x_s$ is called the **entering variable**, and x_r is called a **leaving variable**.

- $\bullet \mathbf{c}^T \mathbf{x}' = \sum_{i=1}^m c_i x'_i + c_s x'_s$
 $= \mathbf{c}^T \mathbf{x} + x'_s (c_s - z_s),$
 $\Rightarrow \mathbf{c}^T \mathbf{x}' \leq \mathbf{c}^T \mathbf{x}.$
- $\bullet \mathbf{c}^T \mathbf{x}' < \mathbf{c}^T \mathbf{x}$ if $x'_s > 0$, which is when the **minimum ratio**
 $\frac{x_r}{u_{rs}} > 0.$
- \bullet Let $x'_s = \frac{x_r}{u_{rs}}.$
- \bullet Then $x'_r = x_r - u_{rs} \frac{x_r}{u_{rs}} = 0$
- $\bullet x_s$ is called the **entering variable**, and x_r is called a **leaving variable**.
- \bullet If there exists $r, t \in \{m+1, \dots, n\}$
 $\frac{x_r}{u_{rs}} = \frac{x_t}{u_{ts}} = \min \left\{ \frac{x_i}{u_{is}} : u_{is} > 0 \right\},$
 then take any **one** of r, t as the **leaving variable**.

- x' is a **BFS** of the LPP.

- \mathbf{x}' is a **BFS** of the LPP.
- The basis matrix corresponding to \mathbf{x}' is
$$B' = [\tilde{\mathbf{a}}_1 \dots \tilde{\mathbf{a}}_{r-1} \tilde{\mathbf{a}}_s \tilde{\mathbf{a}}_{r+1} \dots \tilde{\mathbf{a}}_m].$$

- \mathbf{x}' is a **BFS** of the LPP.
- The basis matrix corresponding to \mathbf{x}' is

$$B' = [\tilde{\mathbf{a}}_1 \dots \tilde{\mathbf{a}}_{r-1} \tilde{\mathbf{a}}_s \tilde{\mathbf{a}}_{r+1} \dots \tilde{\mathbf{a}}_m].$$
- $\tilde{\mathbf{a}}_k = \sum_{i=1, i \neq r}^m (u_{ik} - \frac{u_{is}}{u_{rs}} u_{rk}) \tilde{\mathbf{a}}_i + \tilde{\mathbf{a}}_s \frac{u_{rk}}{u_{rs}}.$

- \mathbf{x}' is a **BFS** of the LPP.
- The basis matrix corresponding to \mathbf{x}' is

$$B' = [\tilde{\mathbf{a}}_1 \dots \tilde{\mathbf{a}}_{r-1} \tilde{\mathbf{a}}_s \tilde{\mathbf{a}}_{r+1} \dots \tilde{\mathbf{a}}_m].$$
- $\tilde{\mathbf{a}}_k = \sum_{i=1, i \neq r}^m (u_{ik} - \frac{u_{is}}{u_{rs}} u_{rk}) \tilde{\mathbf{a}}_i + \tilde{\mathbf{a}}_s \frac{u_{rk}}{u_{rs}}.$
- For all $k = 1, \dots, n$, the new u_{ik} 's are given by,

- \mathbf{x}' is a **BFS** of the LPP.
- The basis matrix corresponding to \mathbf{x}' is

$$B' = [\tilde{\mathbf{a}}_1 \dots \tilde{\mathbf{a}}_{r-1} \tilde{\mathbf{a}}_s \tilde{\mathbf{a}}_{r+1} \dots \tilde{\mathbf{a}}_m].$$
- $\tilde{\mathbf{a}}_k = \sum_{i=1, i \neq r}^m (u_{ik} - \frac{u_{is}}{u_{rs}} u_{rk}) \tilde{\mathbf{a}}_i + \tilde{\mathbf{a}}_s \frac{u_{rk}}{u_{rs}}.$
- For all $k = 1, \dots, n$, the new u_{ik} 's are given by,

$$u'_{ik} = u_{ik} - \frac{u_{is}}{u_{rs}} u_{rk} \text{ for } i = 1, \dots, r-1, r+1, m$$

- \mathbf{x}' is a **BFS** of the LPP.
- The basis matrix corresponding to \mathbf{x}' is

$$B' = [\tilde{\mathbf{a}}_1 \dots \tilde{\mathbf{a}}_{r-1} \tilde{\mathbf{a}}_s \tilde{\mathbf{a}}_{r+1} \dots \tilde{\mathbf{a}}_m].$$
- $\tilde{\mathbf{a}}_k = \sum_{i=1, i \neq r}^m (u_{ik} - \frac{u_{is}}{u_{rs}} u_{rk}) \tilde{\mathbf{a}}_i + \tilde{\mathbf{a}}_s \frac{u_{rk}}{u_{rs}}.$
- For all $k = 1, \dots, n$, the new u_{ik} 's are given by,

$$u'_{ik} = u_{ik} - \frac{u_{is}}{u_{rs}} u_{rk} \text{ for } i = 1, \dots, r-1, r+1, m$$
and $u'_{sk} = \frac{u_{rk}}{u_{rs}}.$

- \mathbf{x}' is a **BFS** of the LPP.
- The basis matrix corresponding to \mathbf{x}' is

$$B' = [\tilde{\mathbf{a}}_1 \dots \tilde{\mathbf{a}}_{r-1} \tilde{\mathbf{a}}_s \tilde{\mathbf{a}}_{r+1} \dots \tilde{\mathbf{a}}_m].$$
- $\tilde{\mathbf{a}}_k = \sum_{i=1, i \neq r}^m (u_{ik} - \frac{u_{is}}{u_{rs}} u_{rk}) \tilde{\mathbf{a}}_i + \tilde{\mathbf{a}}_s \frac{u_{rk}}{u_{rs}}.$
- For all $k = 1, \dots, n$, the new u_{ik} 's are given by,

$$u'_{ik} = u_{ik} - \frac{u_{is}}{u_{rs}} u_{rk} \text{ for } i = 1, \dots, r-1, r+1, m$$
and $u'_{sk} = \frac{u_{rk}}{u_{rs}}.$
- $\mathbf{b} = \sum_{i=1, i \neq r}^m (x_i - \frac{u_{is}}{u_{rs}} x_r) \tilde{\mathbf{a}}_i + \tilde{\mathbf{a}}_s (\frac{x_r}{u_{rs}})$

- \mathbf{x}' is a **BFS** of the LPP.
- The basis matrix corresponding to \mathbf{x}' is

$$B' = [\tilde{\mathbf{a}}_1 \dots \tilde{\mathbf{a}}_{r-1} \tilde{\mathbf{a}}_s \tilde{\mathbf{a}}_{r+1} \dots \tilde{\mathbf{a}}_m].$$
- $\tilde{\mathbf{a}}_k = \sum_{i=1, i \neq r}^m (u_{ik} - \frac{u_{is}}{u_{rs}} u_{rk}) \tilde{\mathbf{a}}_i + \tilde{\mathbf{a}}_s \frac{u_{rk}}{u_{rs}}.$
- For all $k = 1, \dots, n$, the new u_{ik} 's are given by,

$$u'_{ik} = u_{ik} - \frac{u_{is}}{u_{rs}} u_{rk} \text{ for } i = 1, \dots, r-1, r+1, m$$
and $u'_{sk} = \frac{u_{rk}}{u_{rs}}.$
- $\mathbf{b} = \sum_{i=1, i \neq r}^m (x_i - \frac{u_{is}}{u_{rs}} x_r) \tilde{\mathbf{a}}_i + \tilde{\mathbf{a}}_s (\frac{x_r}{u_{rs}})$
- If z'_k denotes the new values of z_k then

- \mathbf{x}' is a **BFS** of the LPP.
- The basis matrix corresponding to \mathbf{x}' is

$$B' = [\tilde{\mathbf{a}}_1 \dots \tilde{\mathbf{a}}_{r-1} \tilde{\mathbf{a}}_s \tilde{\mathbf{a}}_{r+1} \dots \tilde{\mathbf{a}}_m].$$
- $\tilde{\mathbf{a}}_k = \sum_{i=1, i \neq r}^m (u_{ik} - \frac{u_{is}}{u_{rs}} u_{rk}) \tilde{\mathbf{a}}_i + \tilde{\mathbf{a}}_s \frac{u_{rk}}{u_{rs}}.$
- For all $k = 1, \dots, n$, the new u_{ik} 's are given by,

$$u'_{ik} = u_{ik} - \frac{u_{is}}{u_{rs}} u_{rk} \text{ for } i = 1, \dots, r-1, r+1, m$$
and $u'_{sk} = \frac{u_{rk}}{u_{rs}}.$
- $\mathbf{b} = \sum_{i=1, i \neq r}^m (x_i - \frac{u_{is}}{u_{rs}} x_r) \tilde{\mathbf{a}}_i + \tilde{\mathbf{a}}_s (\frac{x_r}{u_{rs}})$
- If z'_k denotes the new values of z_k then

$$z'_k = z_k + \frac{(c_s - z_s)}{u_{rs}} u_{rk}$$

- \mathbf{x}' is a **BFS** of the LPP.
- The basis matrix corresponding to \mathbf{x}' is

$$B' = [\tilde{\mathbf{a}}_1 \dots \tilde{\mathbf{a}}_{r-1} \tilde{\mathbf{a}}_s \tilde{\mathbf{a}}_{r+1} \dots \tilde{\mathbf{a}}_m].$$
- $\tilde{\mathbf{a}}_k = \sum_{i=1, i \neq r}^m (u_{ik} - \frac{u_{is}}{u_{rs}} u_{rk}) \tilde{\mathbf{a}}_i + \tilde{\mathbf{a}}_s \frac{u_{rk}}{u_{rs}}.$
- For all $k = 1, \dots, n$, the new u_{ik} 's are given by,

$$u'_{ik} = u_{ik} - \frac{u_{is}}{u_{rs}} u_{rk} \text{ for } i = 1, \dots, r-1, r+1, m$$
and $u'_{sk} = \frac{u_{rk}}{u_{rs}}.$
- $\mathbf{b} = \sum_{i=1, i \neq r}^m (x_i - \frac{u_{is}}{u_{rs}} x_r) \tilde{\mathbf{a}}_i + \tilde{\mathbf{a}}_s (\frac{x_r}{u_{rs}})$
- If z'_k denotes the new values of z_k then

$$z'_k = z_k + \frac{(c_s - z_s)}{u_{rs}} u_{rk}$$
and $c_k - z'_k = (c_k - z_k) - \frac{(c_s - z_s)}{u_{rs}} u_{rk}.$

- The simplex table corresponding to the new BFS \mathbf{x}' is given by

	$C_s - Z'_s$ $C_s - Z_s - \frac{(C_s - Z_s)}{u_{rs}} u_{rs} = 0$	$C_k - Z'_k$ $(C_k - Z_k) - \frac{(C_s - Z_s)}{u_{rs}} u_{rk}$	
	$B'^{-1} \tilde{\mathbf{a}}_s$	$B'^{-1} \tilde{\mathbf{a}}_k$	$B'^{-1} \mathbf{b}$
$\tilde{\mathbf{a}}_1$	$u_{1s} - \frac{u_{1s}}{u_{rs}} u_{rs} = 0$	$u_{1k} - \frac{u_{1s}}{u_{rs}} u_{rk}$	$x_1 - \frac{u_{1s}}{u_{rs}} x_r$
$\tilde{\mathbf{a}}_2$	$u_{2s} - \frac{u_{2s}}{u_{rs}} u_{rs} = 0$	$u_{2k} - \frac{u_{2s}}{u_{rs}} u_{rk}$	$x_2 - \frac{u_{2s}}{u_{rs}} x_r$
\vdots	0	\vdots	\vdots
$\tilde{\mathbf{a}}_r$	$\frac{u_{rs}}{u_{rs}} = 1$	$\frac{u_{rk}}{u_{rs}}$	$\frac{x_r}{u_{rs}}$
\vdots	\vdots	\vdots	\vdots
$\tilde{\mathbf{a}}_m$	$u_{ms} - \frac{u_{ms}}{u_{rs}} u_{rs} = 0$	$u_{mk} - \frac{u_{ms}}{u_{rs}} u_{rk}$	$x_m - \frac{u_{ms}}{u_{rs}} x_r$

- The simplex table corresponding to the new BFS \mathbf{x}' is given by

	$C_s - Z'_s$ $C_s - Z_s - \frac{(C_s - Z_s)}{u_{rs}} u_{rs} = 0$	$C_k - Z'_k$ $(C_k - Z_k) - \frac{(C_s - Z_s)}{u_{rs}} u_{rk}$	
	$B'^{-1} \tilde{\mathbf{a}}_s$	$B'^{-1} \tilde{\mathbf{a}}_k$	$B'^{-1} \mathbf{b}$
$\tilde{\mathbf{a}}_1$	$u_{1s} - \frac{u_{1s}}{u_{rs}} u_{rs} = 0$	$u_{1k} - \frac{u_{1s}}{u_{rs}} u_{rk}$	$x_1 - \frac{u_{1s}}{u_{rs}} x_r$
$\tilde{\mathbf{a}}_2$	$u_{2s} - \frac{u_{2s}}{u_{rs}} u_{rs} = 0$	$u_{2k} - \frac{u_{2s}}{u_{rs}} u_{rk}$	$x_2 - \frac{u_{2s}}{u_{rs}} x_r$
\vdots	0	\vdots	\vdots
$\tilde{\mathbf{a}}_r$	$\frac{u_{rs}}{u_{rs}} = 1$	$\frac{u_{rk}}{u_{rs}}$	$\frac{x_r}{u_{rs}}$
\vdots	\vdots	\vdots	\vdots
$\tilde{\mathbf{a}}_m$	$u_{ms} - \frac{u_{ms}}{u_{rs}} u_{rs} = 0$	$u_{mk} - \frac{u_{ms}}{u_{rs}} u_{rk}$	$x_m - \frac{u_{ms}}{u_{rs}} x_r$

- The entry u_{rs} of the previous table which is made 1 (by dividing) in this table is called the **pivot element**.

- **Case 1b:** For all $i = 1, \dots, m$, $u_{is} \leq 0$ (where s is as defined in **Case 1**).

- **Case 1b:** For all $i = 1, \dots, m$, $u_{is} \leq 0$ (where s is as defined in **Case 1**).
- Then $x' \geq 0$, for all $x'_s \geq 0$,

- **Case 1b:** For all $i = 1, \dots, m$, $u_{is} \leq 0$ (where s is as defined in **Case 1**).
- Then $\mathbf{x}' \geq 0$, for all $x'_s \geq 0$,
 $\Rightarrow \mathbf{x}' \in \text{Fea}(LPP)$ for all $x'_s \geq 0$.

- **Case 1b:** For all $i = 1, \dots, m$, $u_{is} \leq 0$ (where s is as defined in **Case 1**).
- Then $\mathbf{x}' \geq 0$, for all $x'_s \geq 0$,
 $\Rightarrow \mathbf{x}' \in \text{Fea}(LPP)$ for all $x'_s \geq 0$.
- Since $\mathbf{c}^T \mathbf{x}' = \mathbf{c}^T \mathbf{x} + x'_s(c_s - z_s)$, $(c_s - z_s) < 0$.

- **Case 1b:** For all $i = 1, \dots, m$, $u_{is} \leq 0$ (where s is as defined in **Case 1**).
- Then $\mathbf{x}' \geq \mathbf{0}$, for all $x'_s \geq 0$,
 $\Rightarrow \mathbf{x}' \in \text{Fea}(LPP)$ for all $x'_s \geq 0$.
- Since $\mathbf{c}^T \mathbf{x}' = \mathbf{c}^T \mathbf{x} + x'_s(\mathbf{c}_s - \mathbf{z}_s)$, $(\mathbf{c}_s - \mathbf{z}_s) < 0$.
 So given any $M \in \mathbb{R}$, by taking x'_s sufficiently large we can make $\mathbf{c}^T \mathbf{x}'$ smaller than M .

- **Case 1b:** For all $i = 1, \dots, m$, $u_{is} \leq 0$ (where s is as defined in **Case 1**).
- Then $\mathbf{x}' \geq \mathbf{0}$, for all $x'_s \geq 0$,
 $\Rightarrow \mathbf{x}' \in \text{Fea}(LPP)$ for all $x'_s \geq 0$.
- Since $\mathbf{c}^T \mathbf{x}' = \mathbf{c}^T \mathbf{x} + x'_s(\mathbf{c}_s - z_s)$, $(\mathbf{c}_s - z_s) < 0$.
 So given any $M \in \mathbb{R}$, by taking x'_s sufficiently large we can make $\mathbf{c}^T \mathbf{x}'$ smaller than M .
 So the LPP does **not** have an optimal solution.

- $D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, A_{m \times n} \mathbf{d} = \mathbf{0}, \mathbf{d} \geq \mathbf{0}\}$ is the set of all directions of $S = \text{Fea}(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}, \text{rank}(A) = m\}$

- $D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, A_{m \times n} \mathbf{d} = \mathbf{0}, \mathbf{d} \geq \mathbf{0}\}$ is the set of all directions of $S = \text{Fea}(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}, \text{rank}(A) = m\}$
- If for some basis matrix B and a column $\tilde{\mathbf{a}}_s$ of A , $B^{-1} \tilde{\mathbf{a}}_s \leq \mathbf{0}$

$$\mathbf{d} = \begin{bmatrix} -B^{-1} \tilde{\mathbf{a}}_s \\ 0 \\ \vdots \\ 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

is an extreme direction of S ,

- $D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, A_{m \times n} \mathbf{d} = \mathbf{0}, \mathbf{d} \geq \mathbf{0}\}$ is the set of all directions of $S = \text{Fea}(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}, \text{rank}(A) = m\}$
- If for some basis matrix B and a column $\tilde{\mathbf{a}}_s$ of A , $B^{-1} \tilde{\mathbf{a}}_s \leq \mathbf{0}$

$$\mathbf{d} = \begin{bmatrix} -B^{-1} \tilde{\mathbf{a}}_s \\ 0 \\ \vdots \\ 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

is an extreme direction of S , where the entry **1** in the above vector is at the s th position.

- $D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, A_{m \times n} \mathbf{d} = \mathbf{0}, \mathbf{d} \geq \mathbf{0}\}$ is the set of all directions of $S = \text{Fea}(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}, \text{rank}(A) = m\}$
- If for some basis matrix B and a column $\tilde{\mathbf{a}}_s$ of A , $B^{-1} \tilde{\mathbf{a}}_s \leq \mathbf{0}$

$$\mathbf{d} = \begin{bmatrix} -B^{-1} \tilde{\mathbf{a}}_s \\ 0 \\ \vdots \\ 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

is an extreme direction of S , where the entry **1** in the above vector is at the s th position.

- If \mathbf{d} is as above then $\mathbf{c}^T \mathbf{d} < 0$.

- $D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, A_{m \times n} \mathbf{d} = \mathbf{0}, \mathbf{d} \geq \mathbf{0}\}$ is the set of all directions of $S = \text{Fea}(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}, \text{rank}(A) = m\}$
- If for some basis matrix B and a column $\tilde{\mathbf{a}}_s$ of A , $B^{-1} \tilde{\mathbf{a}}_s \leq \mathbf{0}$

$$\mathbf{d} = \begin{bmatrix} -B^{-1} \tilde{\mathbf{a}}_s \\ 0 \\ \vdots \\ 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

is an extreme direction of S , where the entry **1** in the above vector is at the s th position.

- If \mathbf{d} is as above then $\mathbf{c}^T \mathbf{d} < 0$.
- **Case 2: (Optimality Condition)** (sufficient condition)
 $c_k - z_k \geq 0$ for all $k = 1, \dots, n$.

- $D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, A_{m \times n} \mathbf{d} = \mathbf{0}, \mathbf{d} \geq \mathbf{0}\}$ is the set of all directions of $S = \text{Fea}(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}, \text{rank}(A) = m\}$
- If for some basis matrix B and a column $\tilde{\mathbf{a}}_s$ of A , $B^{-1} \tilde{\mathbf{a}}_s \leq \mathbf{0}$

$$\mathbf{d} = \begin{bmatrix} -B^{-1} \tilde{\mathbf{a}}_s \\ 0 \\ \vdots \\ 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

is an extreme direction of S , where the entry **1** in the above vector is at the s th position.

- If \mathbf{d} is as above then $\mathbf{c}^T \mathbf{d} < 0$.
- **Case 2: (Optimality Condition)** (sufficient condition)
 $c_k - z_k \geq 0$ for all $k = 1, \dots, n$.
 Then \mathbf{x} is optimal for the LPP.

- If instead the LPP would have been a **maximization problem** as given below:

- If instead the LPP would have been a **maximization problem** as given below:

$$\text{Max } \mathbf{c}^T \mathbf{x}$$

subject to

$$\mathbf{A}_{m \times n} \mathbf{x} = \mathbf{b}, \quad \mathbf{x} \geq \mathbf{0}, \quad \text{rank}(\mathbf{A}) = m$$

- If instead the LPP would have been a **maximization problem** as given below:
Max $\mathbf{c}^T \mathbf{x}$
subject to
 $\mathbf{A}_{m \times n} \mathbf{x} = \mathbf{b}$, $\mathbf{x} \geq \mathbf{0}$, $\text{rank}(\mathbf{A}) = m$
- **Case 1'**: $c_k - z_k > 0$ for at least one k , $k = m + 1, \dots, n$.

- If instead the LPP would have been a **maximization problem** as given below:

$$\text{Max } \mathbf{c}^T \mathbf{x}$$

subject to

$$\mathbf{A}_{m \times n} \mathbf{x} = \mathbf{b}, \quad \mathbf{x} \geq \mathbf{0}, \quad \text{rank}(\mathbf{A}) = m$$

- **Case 1'**: $c_k - z_k > 0$ for at least one $k, k = m + 1, \dots, n$.
s th variable will be the **entering variable** if

$$c_s - z_s = \max\{c_k - z_k : c_k - z_k > 0, \quad k = m + 1, \dots, n\}.$$

- If instead the LPP would have been a **maximization problem** as given below:

$$\text{Max } \mathbf{c}^T \mathbf{x}$$

subject to

$$\mathbf{A}_{m \times n} \mathbf{x} = \mathbf{b}, \quad \mathbf{x} \geq \mathbf{0}, \quad \text{rank}(\mathbf{A}) = m$$

- **Case 1'**: $c_k - z_k > 0$ for at least one $k, k = m + 1, \dots, n$.
 s th variable will be the **entering variable** if

$$c_s - z_s = \max\{c_k - z_k : c_k - z_k > 0, \quad k = m + 1, \dots, n\}.$$

- **Case 1'a**:

$$c_s - z_s = \max\{c_k - z_k : c_k - z_k > 0, \quad k = m + 1, \dots, n\},$$

and for atleast one $i, u_{is} > 0$.

- If instead the LPP would have been a **maximization problem** as given below:

$$\text{Max } \mathbf{c}^T \mathbf{x}$$

subject to

$$\mathbf{A}_{m \times n} \mathbf{x} = \mathbf{b}, \quad \mathbf{x} \geq \mathbf{0}, \quad \text{rank}(\mathbf{A}) = m$$

- **Case 1'**: $\mathbf{c}_k - \mathbf{z}_k > 0$ for at least one k , $k = m + 1, \dots, n$.
s th variable will be the **entering variable** if

$$\mathbf{c}_s - \mathbf{z}_s = \max\{\mathbf{c}_k - \mathbf{z}_k : \mathbf{c}_k - \mathbf{z}_k > 0, \quad k = m + 1, \dots, n\}.$$

- **Case 1'a**:

$$\mathbf{c}_s - \mathbf{z}_s = \max\{\mathbf{c}_k - \mathbf{z}_k : \mathbf{c}_k - \mathbf{z}_k > 0, \quad k = m + 1, \dots, n\},$$

and for atleast one i , $\mathbf{u}_{is} > 0$.

- **Case 1'b**:

$$\mathbf{c}_s - \mathbf{z}_s = \max\{\mathbf{c}_k - \mathbf{z}_k : \mathbf{c}_k - \mathbf{z}_k > 0, \quad k = m + 1, \dots, n\},$$

and for all i , $\mathbf{u}_{is} \leq 0$.

- If instead the LPP would have been a **maximization problem** as given below:

$$\text{Max } \mathbf{c}^T \mathbf{x}$$

subject to

$$\mathbf{A}_{m \times n} \mathbf{x} = \mathbf{b}, \quad \mathbf{x} \geq \mathbf{0}, \quad \text{rank}(\mathbf{A}) = m$$

- **Case 1'**: $\mathbf{c}_k - \mathbf{z}_k > 0$ for at least one k , $k = m + 1, \dots, n$.
s th variable will be the **entering variable** if

$$\mathbf{c}_s - \mathbf{z}_s = \max\{\mathbf{c}_k - \mathbf{z}_k : \mathbf{c}_k - \mathbf{z}_k > 0, \quad k = m + 1, \dots, n\}.$$

- **Case 1'a**:

$$\mathbf{c}_s - \mathbf{z}_s = \max\{\mathbf{c}_k - \mathbf{z}_k : \mathbf{c}_k - \mathbf{z}_k > 0, \quad k = m + 1, \dots, n\},$$

and for atleast one i , $u_{is} > 0$.

- **Case 1'b**:

$$\mathbf{c}_s - \mathbf{z}_s = \max\{\mathbf{c}_k - \mathbf{z}_k : \mathbf{c}_k - \mathbf{z}_k > 0, \quad k = m + 1, \dots, n\},$$

and for all i , $u_{is} \leq 0$.

- **Case 2'**: (**Optimality condition** (sufficient condition))

$$\mathbf{c}_k - \mathbf{z}_k \leq 0 \text{ for all } k = 1, \dots, n.$$

- If instead the LPP would have been a **maximization problem** as given below:

$$\text{Max } \mathbf{c}^T \mathbf{x}$$

subject to

$$\mathbf{A}_{m \times n} \mathbf{x} = \mathbf{b}, \quad \mathbf{x} \geq \mathbf{0}, \quad \text{rank}(\mathbf{A}) = m$$

- **Case 1'**: $\mathbf{c}_k - \mathbf{z}_k > 0$ for at least one k , $k = m + 1, \dots, n$.
s th variable will be the **entering variable** if

$$\mathbf{c}_s - \mathbf{z}_s = \max\{\mathbf{c}_k - \mathbf{z}_k : \mathbf{c}_k - \mathbf{z}_k > 0, \quad k = m + 1, \dots, n\}.$$

- **Case 1'a**:

$$\mathbf{c}_s - \mathbf{z}_s = \max\{\mathbf{c}_k - \mathbf{z}_k : \mathbf{c}_k - \mathbf{z}_k > 0, \quad k = m + 1, \dots, n\},$$

and for atleast one i , $u_{is} > 0$.

- **Case 1'b**:

$$\mathbf{c}_s - \mathbf{z}_s = \max\{\mathbf{c}_k - \mathbf{z}_k : \mathbf{c}_k - \mathbf{z}_k > 0, \quad k = m + 1, \dots, n\},$$

and for all i , $u_{is} \leq 0$.

- **Case 2'**: (**Optimality condition** (sufficient condition))

$$\mathbf{c}_k - \mathbf{z}_k \leq 0 \text{ for all } k = 1, \dots, n.$$

- The above **optimality conditions** (for max and min problems) are **sufficient** but not **necessary** for an optimal solution.

- Consider the Problem (P): Min $x_1 + x_2$ subject to

- Consider the Problem (P): Min $x_1 + x_2$ subject to
 $-x_1 + 2x_2 \leq 4$

- Consider the Problem (P): Min $x_1 + x_2$ subject to

$$-x_1 + 2x_2 \leq 4$$

$$-x_1 + x_2 \leq 1$$

- Consider the Problem (P): Min $x_1 + x_2$ subject to
$$-x_1 + 2x_2 \leq 4$$
$$-x_1 + x_2 \leq 1$$
$$x_1 \geq 0, x_2 \geq 0.$$

- Consider the Problem (P): Min $x_1 + x_2$ subject to
 $-x_1 + 2x_2 \leq 4$
 $-x_1 + x_2 \leq 1$
 $x_1 \geq 0, x_2 \geq 0$.

The simplex table corresponding to the BFS with basic variables x_2 and s_1 is given by:

$c_j - z_j$	2	0	0	-1	
	x_1	x_2	s_1	s_2	$B^{-1}\mathbf{b}$
x_2	-1	1	0	1	1
s_1	1	0	1	-2	1

- Consider the Problem (P): Min $x_1 + x_2$ subject to
 $-x_1 + 2x_2 \leq 4$
 $-x_1 + x_2 \leq 1$
 $x_1 \geq 0, x_2 \geq 0$.

The simplex table corresponding to the BFS with basic variables x_2 and s_1 is given by:

$c_j - z_j$	2	0	0	-1	
	x_1	x_2	s_1	s_2	$B^{-1}\mathbf{b}$
x_2	-1	1	0	1	1
s_1	1	0	1	-2	1

The entering variable is s_2 and the leaving variable is x_2 for the next BFS.

- Consider the Problem (P): Min $x_1 + x_2$ subject to
 $-x_1 + 2x_2 \leq 4$
 $-x_1 + x_2 \leq 1$
 $x_1 \geq 0, x_2 \geq 0$.

The simplex table corresponding to the BFS with basic variables x_2 and s_1 is given by:

$c_j - z_j$	2	0	0	-1	
	x_1	x_2	s_1	s_2	$B^{-1}\mathbf{b}$
x_2	-1	1	0	1	1
s_1	1	0	1	-2	1

The entering variable is s_2 and the leaving variable is x_2 for the next BFS.

$c_j - z_j$	1	1	0	0	
	x_1	x_2	s_1	s_2	$B^{-1}\mathbf{b}$
s_2	-1	1	0	1	1
s_1	-1	2	1	0	4

Note that the above table is optimal.

- Suppose if we change the objective function to Min

$$x_1 - 2x_2$$

- | $c_j - z_j$ | -1 | 0 | 0 | 2 | |
|-------------|-------|-------|-------|-------|--------------------|
| | x_1 | x_2 | s_1 | s_2 | $B^{-1}\mathbf{b}$ |
| x_2 | -1 | 1 | 0 | 1 | 1 |
| s_1 | 1 | 0 | 1 | -2 | 1 |

- Suppose if we change the objective function to Min

$$x_1 - 2x_2$$

- | $c_j - z_j$ | -1 | 0 | 0 | 2 | |
|-------------|-------|-------|-------|-------|--------------------|
| | x_1 | x_2 | s_1 | s_2 | $B^{-1}\mathbf{b}$ |
| x_2 | -1 | 1 | 0 | 1 | 1 |
| s_1 | 1 | 0 | 1 | -2 | 1 |

The entering variable is x_1 and the leaving variable is s_1 for the next BFS.

- Suppose if we change the objective function to Min

$$x_1 - 2x_2$$

- | $c_j - z_j$ | -1 | 0 | 0 | 2 | |
|-------------|-------|-------|-------|-------|--------------------|
| | x_1 | x_2 | s_1 | s_2 | $B^{-1}\mathbf{b}$ |
| x_2 | -1 | 1 | 0 | 1 | 1 |
| s_1 | 1 | 0 | 1 | -2 | 1 |

The entering variable is x_1 and the leaving variable is s_1 for the next BFS.

Then the next table is

- Suppose if we change the objective function to Min

$$x_1 - 2x_2$$

- | $c_j - z_j$ | -1 | 0 | 0 | 2 | |
|-------------|-------|-------|-------|-------|--------------------|
| | x_1 | x_2 | s_1 | s_2 | $B^{-1}\mathbf{b}$ |
| x_2 | -1 | 1 | 0 | 1 | 1 |
| s_1 | 1 | 0 | 1 | -2 | 1 |

The entering variable is x_1 and the leaving variable is s_1 for the next BFS.

Then the next table is

- | $c_j - z_j$ | 0 | 0 | 1 | 0 | |
|-------------|-------|-------|-------|-------|--------------------|
| | x_1 | x_2 | s_1 | s_2 | $B^{-1}\mathbf{b}$ |
| x_2 | 0 | 1 | 1 | -1 | 3 |
| x_1 | 1 | 0 | 1 | -2 | 2 |

The above table is optimal.