

# Plan

- Examples of optimization problems
- Examples of Linear Programming Problems (LPP)
- Solution of an LPP by the Graphical Method
- Extreme points and Corner points
- Exercises and Questions

# Optimization Problems

# Optimization Problems

- Nothing takes place in the world whose meaning is not that of some minimum or maximum: [L. Euler](#)

# Optimization Problems

- Nothing takes place in the world whose meaning is not that of some minimum or maximum: [L. Euler](#)
- **Heron's problem:** 'On Mirrors' ( book related to laws of reflection of light, approx 1st century AD).

# Optimization Problems

- Nothing takes place in the world whose meaning is not that of some minimum or maximum: L. Euler
- Heron's problem: 'On Mirrors' ( book related to laws of reflection of light, approx 1st century AD).  
A and B are two given points on the same side of a line L.  
Find a point D on L such that the sum of the distances from A to D and from B to D is a minimum.

# Optimization Problems

- Nothing takes place in the world whose meaning is not that of some minimum or maximum: L. Euler
- Heron's problem: 'On Mirrors' ( book related to laws of reflection of light, approx 1st century AD).  
A and B are two given points on the same side of a line L.  
Find a point D on L such that the sum of the distances from A to D and from B to D is a minimum.
- Extremal principles of nature:

# Optimization Problems

- Nothing takes place in the world whose meaning is not that of some minimum or maximum: L. Euler
- Heron's problem: 'On Mirrors' ( book related to laws of reflection of light, approx 1st century AD).  
A and B are two given points on the same side of a line L. Find a point D on L such that the sum of the distances from A to D and from B to D is a minimum.
- Extremal principles of nature:
- Laws of reflection of light.  
Nature breaks a ray of light at equal angles, if it does not unnecessarily want it to meander to no purpose.

# Optimization Problems

- Nothing takes place in the world whose meaning is not that of some minimum or maximum: L. Euler
- Heron's problem: 'On Mirrors' ( book related to laws of reflection of light, approx 1st century AD).  
A and B are two given points on the same side of a line L. Find a point D on L such that the sum of the distances from A to D and from B to D is a minimum.
- Extremal principles of nature:
- Laws of reflection of light.  
Nature breaks a ray of light at equal angles, if it does not unnecessarily want it to meander to no purpose.
- Laws of refraction of light.  
What characterizes the trajectory of light moving from one point to another in a non homogeneous medium is that it is traversed in a minimum time.



- Isoperimetric problem, Oldest version: 'Aenid' of Vergil.

- Isoperimetric problem, Oldest version: 'Aenid' of Vergil.  
Escaping from her brother, the Phoenician princess  
(Phoenicia is Lebanon, with parts of Syria, Israel) Dido set  
off westward in search of a safe place to settle down.

- **Isoperimetric problem**, **Oldest version**: 'Aenid' of Vergil.  
Escaping from her brother, the Phoenician princess  
(Phoenicia is Lebanon, with parts of Syria, Israel) Dido set  
off westward in search of a safe place to settle down.  
She liked a certain place now known as 'Bay of Tunis'.

- **Isoperimetric problem**, **Oldest version**: 'Aenid' of Vergil.  
Escaping from her brother, the Phoenician princess  
(Phoenicia is Lebanon, with parts of Syria, Israel) Dido set  
off westward in search of a safe place to settle down.  
She liked a certain place now known as 'Bay of Tunis'.  
Dido asked the local leader Yarb for as much land as could  
be 'encircled with a bull's hide' ( 9th century BC).

- **Isoperimetric problem**, **Oldest version**: 'Aenid' of Vergil.  
Escaping from her brother, the Phoenician princess (Phoenicia is Lebanon, with parts of Syria, Israel) Dido set off westward in search of a safe place to settle down. She liked a certain place now known as 'Bay of Tunis'. Dido asked the local leader Yarb for as much land as could be 'encircled with a bull's hide' ( 9th century BC).
- **Isoperimetric problem**: Among all closed plane curves of a given length( perimeter), find the one that encloses the maximum area.

- **Isoperimetric problem**, **Oldest version**: 'Aenid' of Vergil.  
Escaping from her brother, the Phoenician princess (Phoenicia is Lebanon, with parts of Syria, Israel) Dido set off westward in search of a safe place to settle down. She liked a certain place now known as 'Bay of Tunis'. Dido asked the local leader Yarb for as much land as could be 'encircled with a bull's hide' ( 9th century BC).
- **Isoperimetric problem**: Among all closed plane curves of a given length( perimeter), find the one that encloses the **maximum area**. **Answer**: Circle

- **Isoperimetric problem**, **Oldest version**: 'Aenid' of Vergil.  
Escaping from her brother, the Phoenician princess (Phoenicia is Lebanon, with parts of Syria, Israel) Dido set off westward in search of a safe place to settle down. She liked a certain place now known as 'Bay of Tunis'. Dido asked the local leader Yarb for as much land as could be 'encircled with a bull's hide' ( 9th century BC).
- **Isoperimetric problem**: Among all closed plane curves of a given length( perimeter), find the one that encloses the **maximum area**. **Answer**: Circle
- **Isoepiphanic property of the Sphere**: The Sphere encloses the largest volume among all closed surfaces with the same surface area.

- **Isoperimetric problem**, **Oldest version**: 'Aenid' of Vergil.  
Escaping from her brother, the Phoenician princess (Phoenicia is Lebanon, with parts of Syria, Israel) Dido set off westward in search of a safe place to settle down. She liked a certain place now known as 'Bay of Tunis'. Dido asked the local leader Yarb for as much land as could be 'encircled with a bull's hide' ( 9th century BC).
- **Isoperimetric problem**: Among all closed plane curves of a given length( perimeter), find the one that encloses the **maximum area**. **Answer**: Circle
- **Isoepiphanic property of the Sphere**: The Sphere encloses the largest volume among all closed surfaces with the same surface area.
- **Easier problem**: Among all rectangles of a given perimeter (say 20), find the one that encloses the **maximum area**.  
Maximize *ac*



- **Isoperimetric problem**, **Oldest version**: ‘Aenid’ of Vergil.  
Escaping from her brother, the Phoenician princess (Phoenicia is Lebanon, with parts of Syria, Israel) Dido set off westward in search of a safe place to settle down. She liked a certain place now known as ‘Bay of Tunis’. Dido asked the local leader Yarb for as much land as could be ‘**encircled with a bull’s hide**’ ( 9th century BC).
- **Isoperimetric problem**: Among all closed plane curves of a given length( perimeter), find the one that encloses the **maximum area**. **Answer**: Circle
- **Isoepiphanic property of the Sphere**: The Sphere encloses the largest volume among all closed surfaces with the same surface area.
- **Easier problem**: Among all rectangles of a given perimeter (say 20), find the one that encloses the **maximum area**.  
Maximize ***ac***  
subject to,  $a + c = 10$ ,  
 $a \geq 0, c \geq 0$ .

- **Isoperimetric problem**, **Oldest version**: 'Aenid' of Vergil.  
Escaping from her brother, the Phoenician princess (Phoenicia is Lebanon, with parts of Syria, Israel) Dido set off westward in search of a safe place to settle down. She liked a certain place now known as 'Bay of Tunis'. Dido asked the local leader Yarb for as much land as could be 'encircled with a bull's hide' (9th century BC).
- **Isoperimetric problem**: Among all closed plane curves of a given length (perimeter), find the one that encloses the **maximum area**. **Answer**: Circle
- **Isoepiphanic property of the Sphere**: The Sphere encloses the largest volume among all closed surfaces with the same surface area.
- **Easier problem**: Among all rectangles of a given perimeter (say 20), find the one that encloses the **maximum area**.  
Maximize  **$ac$**   
subject to,  **$a + c = 10$** ,  
 **$a \geq 0, c \geq 0$** .  
**Answer**: Square

# Linear Programming Problems

Diet Problem: For U.S soldiers, World War II

- Let there be  $m$  nutrients  $N_1, N_2, \dots, N_m$  and  $n$  food products,  $F_1, F_2, \dots, F_n$ , available in the market which can supply these nutrients.

# Linear Programming Problems

Diet Problem: For U.S soldiers, World War II

- Let there be  $m$  nutrients  $N_1, N_2, \dots, N_m$  and  $n$  food products,  $F_1, F_2, \dots, F_n$ , available in the market which can supply these nutrients.

For healthy survival a human being requires say atleast,  $b_i$  units of the  $i$  th nutrient,  $i = 1, 2, \dots, m$ , respectively.

# Linear Programming Problems

## Diet Problem: For U.S soldiers, World War II

- Let there be  $m$  nutrients  $N_1, N_2, \dots, N_m$  and  $n$  food products,  $F_1, F_2, \dots, F_n$ , available in the market which can supply these nutrients.

For healthy survival a human being requires say atleast,  $b_i$  units of the  $i$  th nutrient,  $i = 1, 2, \dots, m$ , respectively.

Let  $a_{ij}$  be the amount of the  $i$  th nutrient ( $N_i$ ) present in unit amount of the  $j$  th food product ( $F_j$ ), and let  $c_j, j = 1, 2, \dots, n$  be the cost of unit amount of  $F_j$ .

# Linear Programming Problems

## Diet Problem: For U.S soldiers, World War II

- Let there be  $m$  nutrients  $N_1, N_2, \dots, N_m$  and  $n$  food products,  $F_1, F_2, \dots, F_n$ , available in the market which can supply these nutrients.

For healthy survival a human being requires say atleast,  $b_i$  units of the  $i$  th nutrient,  $i = 1, 2, \dots, m$ , respectively.

Let  $a_{ij}$  be the amount of the  $i$  th nutrient ( $N_i$ ) present in unit amount of the  $j$  th food product ( $F_j$ ), and let  $c_j, j = 1, 2, \dots, n$  be the cost of unit amount of  $F_j$ .

So the problem is to decide on a diet of minimum cost consisting of the  $n$  food products ( in various quantities) so that one gets the required amount of each of the nutrients.

# Formulation of the Diet Problem

# Formulation of the Diet Problem

Let  $x_j$  be the amount of the  $j$ th food product to be used in the diet (so  $x_j \geq 0$  for  $j = 1, 2, \dots, n$ ,) and the problem reduces to (under certain simplifying assumptions):



# Formulation of the Diet Problem

Let  $x_j$  be the amount of the  $j$ th food product to be used in the diet (so  $x_j \geq 0$  for  $j = 1, 2, \dots, n$ ,) and the problem reduces to (under certain simplifying assumptions):

$$\text{Min } \sum_{j=1}^n c_j x_j = \mathbf{c}^T \mathbf{x}$$

# Formulation of the Diet Problem

Let  $x_j$  be the amount of the  $j$ th food product to be used in the diet (so  $x_j \geq 0$  for  $j = 1, 2, \dots, n$ ), and the problem reduces to (under certain simplifying assumptions):

$$\text{Min } \sum_{j=1}^n c_j x_j = \mathbf{c}^T \mathbf{x}$$

subject to

$$\sum_{j=1}^n a_{ij} x_j \geq b_i, \text{ for } i = 1, 2, \dots, m,$$

# Formulation of the Diet Problem

Let  $x_j$  be the amount of the  $j$ th food product to be used in the diet (so  $x_j \geq 0$  for  $j = 1, 2, \dots, n$ ), and the problem reduces to (under certain simplifying assumptions):

$$\text{Min } \sum_{j=1}^n c_j x_j = \mathbf{c}^T \mathbf{x}$$

subject to

$$\sum_{j=1}^n a_{ij} x_j \geq b_i, \text{ for } i = 1, 2, \dots, m,$$

$$x_j \geq 0 \text{ for all } j = 1, 2, \dots, n.$$

## Formulation of the Diet Problem

Let  $x_j$  be the amount of the  $j$ th food product to be used in the diet (so  $x_j \geq 0$  for  $j = 1, 2, \dots, n$ ), and the problem reduces to (under certain simplifying assumptions):

$$\text{Min } \sum_{j=1}^n c_j x_j = \mathbf{c}^T \mathbf{x}$$

subject to

$$\sum_{j=1}^n a_{ij} x_j \geq b_i, \text{ for } i = 1, 2, \dots, m,$$

$$x_j \geq 0 \text{ for all } j = 1, 2, \dots, n.$$

or as

$$\mathbf{Ax} \geq \mathbf{b}, \text{ ( or alternatively as } -\mathbf{Ax} \leq -\mathbf{b} ), \mathbf{x} \geq \mathbf{0},$$

## Formulation of the Diet Problem

Let  $x_j$  be the amount of the  $j$ th food product to be used in the diet (so  $x_j \geq 0$  for  $j = 1, 2, \dots, n$ ), and the problem reduces to (under certain simplifying assumptions):

$$\text{Min } \sum_{j=1}^n c_j x_j = \mathbf{c}^T \mathbf{x}$$

subject to

$$\sum_{j=1}^n a_{ij} x_j \geq b_i, \text{ for } i = 1, 2, \dots, m,$$

$$x_j \geq 0 \text{ for all } j = 1, 2, \dots, n.$$

or as

$$\mathbf{Ax} \geq \mathbf{b}, \text{ ( or alternatively as } -\mathbf{Ax} \leq -\mathbf{b} ), \mathbf{x} \geq \mathbf{0},$$

where  $\mathbf{A}$  is an  $m \times n$  matrix (a matrix with  $m$  rows and  $n$  columns), the  $(i, j)$  th entry of  $\mathbf{A}$  is given by  $a_{ij}$ ,

# Formulation of the Diet Problem

Let  $x_j$  be the amount of the  $j$ th food product to be used in the diet (so  $x_j \geq 0$  for  $j = 1, 2, \dots, n$ ), and the problem reduces to (under certain simplifying assumptions):

$$\text{Min } \sum_{j=1}^n c_j x_j = \mathbf{c}^T \mathbf{x}$$

subject to

$$\sum_{j=1}^n a_{ij} x_j \geq b_i, \text{ for } i = 1, 2, \dots, m,$$

$$x_j \geq 0 \text{ for all } j = 1, 2, \dots, n.$$

or as

$$\mathbf{Ax} \geq \mathbf{b}, \text{ ( or alternatively as } -\mathbf{Ax} \leq -\mathbf{b} \text{ ), } \mathbf{x} \geq \mathbf{0},$$

where  $\mathbf{A}$  is an  $m \times n$  matrix (a matrix with  $m$  rows and  $n$  columns), the  $(i, j)$  th entry of  $\mathbf{A}$  is given by  $a_{ij}$ ,

$\mathbf{b} = [b_1, b_2, \dots, b_m]^T$  and  $\mathbf{x} = [x_1, x_2, \dots, x_n]^T$ ,  $\mathbf{0}$  is the zero vector with  $n$  components.

## Transportation Problem: Soviet Union 1940's

## Transportation Problem: Soviet Union 1940's

Let there be  $m$  **supply stations**,  $S_1, S_2, \dots, S_m$  for a particular **product** (P) and  $n$  **destination stations**,  $D_1, D_2, \dots, D_n$  where the product is to be transported.



## Transportation Problem: Soviet Union 1940's

Let there be  $m$  **supply stations**,  $S_1, S_2, \dots, S_m$  for a particular **product** (P) and  $n$  **destination stations**,  $D_1, D_2, \dots, D_n$  where the product is to be transported.

Let  $c_{ij}$  be the **cost of transportation** of unit amount of the product (P) from  $S_i$  to  $D_j$ .

## Transportation Problem: Soviet Union 1940's

Let there be  $m$  **supply stations**,  $S_1, S_2, \dots, S_m$  for a particular **product** (P) and  $n$  **destination stations**,  $D_1, D_2, \dots, D_n$  where the product is to be transported.

Let  $c_{ij}$  be the **cost of transportation** of unit amount of the product (P) from  $S_i$  to  $D_j$ .

Let  $s_i$  be the amount of the product **available** at  $S_i$  and let  $d_j$  be the corresponding **demand** at  $D_j$ .

## Transportation Problem: Soviet Union 1940's

Let there be  $m$  **supply stations**,  $S_1, S_2, \dots, S_m$  for a particular **product** (P) and  $n$  **destination stations**,  $D_1, D_2, \dots, D_n$  where the product is to be transported.

Let  $c_{ij}$  be the **cost of transportation** of unit amount of the product (P) from  $S_i$  to  $D_j$ .

Let  $s_i$  be the amount of the product **available** at  $S_i$  and let  $d_j$  be the corresponding **demand** at  $D_j$ .

The problem is to find  $x_{ij}$ ,  $i = 1, 2, \dots, m$ ,  $j = 1, 2, \dots, n$ , where  $x_{ij}$  is the amount of the product to be transported from  $S_i$  to  $D_j$  such that the demands  $d_j$  are met and the cost of transportation is **minimum**.

## Transportation Problem: Soviet Union 1940's

Let there be  $m$  **supply stations**,  $S_1, S_2, \dots, S_m$  for a particular **product** (P) and  $n$  **destination stations**,  $D_1, D_2, \dots, D_n$  where the product is to be transported.

Let  $c_{ij}$  be the **cost of transportation** of unit amount of the product (P) from  $S_i$  to  $D_j$ .

Let  $s_i$  be the amount of the product **available** at  $S_i$  and let  $d_j$  be the corresponding **demand** at  $D_j$ .

The problem is to find  $x_{ij}$ ,  $i = 1, 2, \dots, m$ ,  $j = 1, 2, \dots, n$ , where  $x_{ij}$  is the amount of the product to be transported from  $S_i$  to  $D_j$  such that the demands  $d_j$  are met and the cost of transportation is **minimum**.

# Formulation of the Transportation Problem

# Formulation of the Transportation Problem

The problem can be modelled as (under certain simplifying assumptions )

$$\text{Min } \sum_{i,j} c_{ij} x_{ij}$$

# Formulation of the Transportation Problem

The problem can be modelled as (under certain simplifying assumptions )

$$\text{Min } \sum_{i,j} c_{ij} x_{ij}$$

subject to

$$\sum_{j=1}^n x_{ij} \leq s_i, \text{ for } i = 1, 2, \dots, m,$$

# Formulation of the Transportation Problem

The problem can be modelled as (under certain simplifying assumptions )

$$\text{Min } \sum_{i,j} c_{ij} x_{ij}$$

subject to

$$\sum_{j=1}^n x_{ij} \leq s_i, \text{ for } i = 1, 2, \dots, m,$$

$$\sum_{i=1}^m x_{ij} \geq d_j, \text{ for } j = 1, 2, \dots, n,$$



# Formulation of the Transportation Problem

The problem can be modelled as (under certain simplifying assumptions )

$$\text{Min } \sum_{i,j} c_{ij} x_{ij}$$

subject to

$$\sum_{j=1}^n x_{ij} \leq s_i, \text{ for } i = 1, 2, \dots, m,$$

$$\sum_{i=1}^m x_{ij} \geq d_j, \text{ for } j = 1, 2, \dots, n,$$

$$x_{ij} \geq 0 \text{ for all } i = 1, 2, \dots, m, \quad j = 1, 2, \dots, n.$$

# Formulation of the Transportation Problem

The problem can be modelled as (under certain simplifying assumptions )

$$\text{Min } \sum_{i,j} c_{ij} x_{ij}$$

subject to

$$\sum_{j=1}^n x_{ij} \leq s_i, \text{ for } i = 1, 2, \dots, m,$$

$$\sum_{i=1}^m x_{ij} \geq d_j, \text{ for } j = 1, 2, \dots, n,$$

$$x_{ij} \geq 0 \text{ for all } i = 1, 2, \dots, m, \quad j = 1, 2, \dots, n.$$

Note that the constraints of the above LPP can again be written as:

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

# Formulation of the Transportation Problem

The problem can be modelled as (under certain simplifying assumptions )

$$\text{Min } \sum_{i,j} c_{ij} x_{ij}$$

subject to

$$\sum_{j=1}^n x_{ij} \leq s_i, \text{ for } i = 1, 2, \dots, m,$$

$$\sum_{i=1}^m x_{ij} \geq d_j, \text{ for } j = 1, 2, \dots, n,$$

$$x_{ij} \geq 0 \text{ for all } i = 1, 2, \dots, m, \quad j = 1, 2, \dots, n.$$

Note that the constraints of the above LPP can again be written as:

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

$$\text{subject to } \mathbf{Ax} \leq \mathbf{b},$$

# Formulation of the Transportation Problem

The problem can be modelled as (under certain simplifying assumptions )

$$\text{Min } \sum_{i,j} c_{ij} x_{ij}$$

subject to

$$\sum_{j=1}^n x_{ij} \leq s_i, \text{ for } i = 1, 2, \dots, m,$$

$$\sum_{i=1}^m x_{ij} \geq d_j, \text{ for } j = 1, 2, \dots, n,$$

$$x_{ij} \geq 0 \text{ for all } i = 1, 2, \dots, m, \quad j = 1, 2, \dots, n.$$

Note that the constraints of the above LPP can again be written as:

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

$$\text{subject to } \mathbf{Ax} \leq \mathbf{b},$$

$$\mathbf{x} \geq \mathbf{0},$$

# Formulation of the Transportation Problem

The problem can be modelled as (under certain simplifying assumptions )

$$\text{Min } \sum_{i,j} c_{ij} x_{ij}$$

subject to

$$\sum_{j=1}^n x_{ij} \leq s_i, \text{ for } i = 1, 2, \dots, m,$$

$$\sum_{i=1}^m x_{ij} \geq d_j, \text{ for } j = 1, 2, \dots, n,$$

$$x_{ij} \geq 0 \text{ for all } i = 1, 2, \dots, m, \quad j = 1, 2, \dots, n.$$

Note that the constraints of the above LPP can again be written as:

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

$$\text{subject to } \mathbf{Ax} \leq \mathbf{b},$$

$$\mathbf{x} \geq \mathbf{0},$$

where  $\mathbf{A}$  is a matrix with  $(m + n)$  rows and  $(m \times n)$  columns,

# Formulation of the Transportation Problem

The problem can be modelled as (under certain simplifying assumptions )

$$\text{Min } \sum_{i,j} c_{ij} x_{ij}$$

subject to

$$\sum_{j=1}^n x_{ij} \leq s_i, \text{ for } i = 1, 2, \dots, m,$$

$$\sum_{i=1}^m x_{ij} \geq d_j, \text{ for } j = 1, 2, \dots, n,$$

$$x_{ij} \geq 0 \text{ for all } i = 1, 2, \dots, m, \quad j = 1, 2, \dots, n.$$

Note that the constraints of the above LPP can again be written as:

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

$$\text{subject to } \mathbf{Ax} \leq \mathbf{b},$$

$$\mathbf{x} \geq \mathbf{0},$$

where  $\mathbf{A}$  is a matrix with  $(m + n)$  rows and  $(m \times n)$  columns,

$\mathbf{x}, \mathbf{0}$  are vectors with  $m \times n$  components

# Formulation of the Transportation Problem

The problem can be modelled as (under certain simplifying assumptions )

$$\text{Min } \sum_{i,j} c_{ij} x_{ij}$$

subject to

$$\sum_{j=1}^n x_{ij} \leq s_i, \text{ for } i = 1, 2, \dots, m,$$

$$\sum_{i=1}^m x_{ij} \geq d_j, \text{ for } j = 1, 2, \dots, n,$$

$$x_{ij} \geq 0 \text{ for all } i = 1, 2, \dots, m, \quad j = 1, 2, \dots, n.$$

Note that the constraints of the above LPP can again be written as:

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

$$\text{subject to } \mathbf{Ax} \leq \mathbf{b},$$

$$\mathbf{x} \geq \mathbf{0},$$

where  $\mathbf{A}$  is a matrix with  $(m + n)$  rows and  $(m \times n)$  columns,

$\mathbf{x}, \mathbf{0}$  are vectors with  $m \times n$  components

and  $\mathbf{b} = [s_1, \dots, s_m, -d_1, \dots, -d_n]^T$ .

For example the 1st row of  $A$  (the row corresponding to the first supply constraint) is given by

$$[1, 1, \dots, 1, 0, \dots, 0]^T$$



For example the 1st row of  $A$  (the row corresponding to the first supply constraint) is given by

$$[1, 1, \dots, 1, 0, \dots, 0]^T$$

that is 1 in the first  $n$  positions and 0's elsewhere.

For example the 1st row of  $A$  (the row corresponding to the first supply constraint) is given by

$$[1, 1, \dots, 1, 0, \dots, 0]^T$$

that is 1 in the first  $n$  positions and 0's elsewhere.

The second row of  $A$  (the row corresponding to the second supply constraint) is given by

$$[0, \dots, 0, 1, 1, \dots, 1, 0, \dots, 0]^T$$

For example the 1st row of  $A$  (the row corresponding to the first supply constraint) is given by

$$[1, 1, \dots, 1, 0, \dots, 0]^T$$

that is 1 in the first  $n$  positions and 0's elsewhere.

The second row of  $A$  (the row corresponding to the second supply constraint) is given by

$$[0, \dots, 0, 1, 1, \dots, 1, 0, \dots, 0]^T$$

that is 1 in the  $(n + 1)$  th position to the  $2n$  th position and 0's elsewhere.

For example the 1st row of  $A$  (the row corresponding to the first supply constraint) is given by

$$[1, 1, \dots, 1, 0, \dots, 0]^T$$

that is 1 in the first  $n$  positions and 0's elsewhere.

The second row of  $A$  (the row corresponding to the second supply constraint) is given by

$$[0, \dots, 0, 1, 1, \dots, 1, 0, \dots, 0]^T$$

that is 1 in the  $(n+1)$  th position to the  $2n$  th position and 0's elsewhere.

The  $m$ th row of  $A$  (the row corresponding to the  $m$  th supply constraint) is given by

$$[0, \dots, 0, 1, 1, \dots, 1]^T$$

For example the 1st row of  $A$  (the row corresponding to the first supply constraint) is given by

$$[1, 1, \dots, 1, 0, \dots, 0]^T$$

that is 1 in the first  $n$  positions and 0's elsewhere.

The second row of  $A$  (the row corresponding to the second supply constraint) is given by

$$[0, \dots, 0, 1, 1, \dots, 1, 0, \dots, 0]^T$$

that is 1 in the  $(n+1)$  th position to the  $2n$  th position and 0's elsewhere.

The  $m$ th row of  $A$  (the row corresponding to the  $m$  th supply constraint) is given by

$$[0, \dots, 0, 1, 1, \dots, 1]^T$$

that is 1 in the  $(m-1)n+1$  th position to the  $mn$  th position and 0's elsewhere.

For example the 1st row of  $A$  (the row corresponding to the first supply constraint) is given by

$$[1, 1, \dots, 1, 0, \dots, 0]^T$$

that is 1 in the first  $n$  positions and 0's elsewhere.

The second row of  $A$  (the row corresponding to the second supply constraint) is given by

$$[0, \dots, 0, 1, 1, \dots, 1, 0, \dots, 0]^T$$

that is 1 in the  $(n+1)$  th position to the  $2n$  th position and 0's elsewhere.

The  $m$ th row of  $A$  (the row corresponding to the  $m$  th supply constraint) is given by

$$[0, \dots, 0, 1, 1, \dots, 1]^T$$

that is 1 in the  $(m-1)n+1$  th position to the  $mn$  th position and 0's elsewhere.

The  $(m + 1)$  th row of  $\mathbf{A}$  (the row corresponding to the first destination constraint) is given by

$$[-1, 0, \dots, 0, -1, 0, \dots, 0, -1, 0, \dots, 0],$$

The  $(m + 1)$  th row of  $A$  (the row corresponding to the first destination constraint) is given by

$$[-1, 0, \dots, 0, -1, 0, \dots, 0, -1, 0, \dots, 0],$$

that is  $-1$  at the first position,  $-1$  at the  $(n+1)$ th position,  $-1$  at the  $(2n+1)$ th position, ...,  $-1$  at the  $((m-1)n + 1)$  th position, etc and 0's elsewhere.



The  $(m + 1)$  th row of  $A$  (the row corresponding to the first destination constraint) is given by

$$[-1, 0, \dots, 0, -1, 0, \dots, 0, -1, 0, \dots, 0],$$

that is  $-1$  at the first position,  $-1$  at the  $(n+1)$ th position,  $-1$  at the  $(2n+1)$ th position, ...,  $-1$  at the  $((m-1)n + 1)$  th position, etc and 0's elsewhere.

The  $(m + n)$  th row of  $A$  (the row corresponding to the  $n$ th (last) destination constraint) is given by

$$[0, \dots, -1, 0, \dots, -1, 0, \dots, -1, 0, \dots, -1, \dots, 0, \dots, -1],$$

The  $(m + 1)$  th row of  $A$  (the row corresponding to the first destination constraint) is given by

$$[-1, 0, \dots, 0, -1, 0, \dots, 0, -1, 0, \dots, 0],$$

that is  $-1$  at the first position,  $-1$  at the  $(n+1)$ th position,  $-1$  at the  $(2n+1)$ th position, ...,  $-1$  at the  $((m-1)n + 1)$  th position, etc and 0's elsewhere.

The  $(m + n)$  th row of  $A$  (the row corresponding to the  $n$ th (last) destination constraint) is given by

$$[0, \dots, -1, 0, \dots, -1, 0, \dots, -1, 0, \dots, -1, \dots, 0, \dots, -1],$$

that is  $-1$  at the  $n$ th position,  $-1$  at the  $2n$  th position,  $-1$  at the  $3n$  th position, ...,  $-1$  at the  $(m \times n)$  th position, etc and 0's elsewhere.

# Linear Programming Problem

Given  $\mathbf{c} \in \mathbb{R}^n$ , a column vector with  $n$  components,  $\mathbf{b} \in \mathbb{R}^m$ , a column vector with  $m$  components, and an  $\mathbf{A} \in \mathbb{R}^{m \times n}$ , a matrix with  $m$  rows and  $n$  columns

# Linear Programming Problem

Given  $\mathbf{c} \in \mathbb{R}^n$ , a column vector with  $n$  components,  $\mathbf{b} \in \mathbb{R}^m$ , a column vector with  $m$  components, and an  $\mathbf{A} \in \mathbb{R}^{m \times n}$ , a matrix with  $m$  rows and  $n$  columns

A **linear programming problem**(LPP) is given by :

Max or Min  $\mathbf{c}^T \mathbf{x}$

# Linear Programming Problem

Given  $\mathbf{c} \in \mathbb{R}^n$ , a column vector with  $n$  components,  $\mathbf{b} \in \mathbb{R}^m$ , a column vector with  $m$  components, and an  $\mathbf{A} \in \mathbb{R}^{m \times n}$ , a matrix with  $m$  rows and  $n$  columns

A **linear programming problem**(LPP) is given by :

Max or Min  $\mathbf{c}^T \mathbf{x}$

subject to  $\mathbf{Ax} \leq \mathbf{b}$  (or  $\mathbf{Ax} \geq \mathbf{b}$ ),

# Linear Programming Problem

Given  $\mathbf{c} \in \mathbb{R}^n$ , a column vector with  $n$  components,  $\mathbf{b} \in \mathbb{R}^m$ , a column vector with  $m$  components, and an  $\mathbf{A} \in \mathbb{R}^{m \times n}$ , a matrix with  $m$  rows and  $n$  columns

A **linear programming problem**(LPP) is given by :

Max or Min  $\mathbf{c}^T \mathbf{x}$

subject to  $\mathbf{Ax} \leq \mathbf{b}$  (or  $\mathbf{Ax} \geq \mathbf{b}$ ),

$\mathbf{x} \geq \mathbf{0}$ .

# Linear Programming Problem

Given  $\mathbf{c} \in \mathbb{R}^n$ , a column vector with  $n$  components,  $\mathbf{b} \in \mathbb{R}^m$ , a column vector with  $m$  components, and an  $\mathbf{A} \in \mathbb{R}^{m \times n}$ , a matrix with  $m$  rows and  $n$  columns

A **linear programming problem**(LPP) is given by :

Max or Min  $\mathbf{c}^T \mathbf{x}$

subject to  $\mathbf{Ax} \leq \mathbf{b}$  (or  $\mathbf{Ax} \geq \mathbf{b}$ ),

$\mathbf{x} \geq \mathbf{0}$ .

The function  $f(\mathbf{x}) = \mathbf{c}^T \mathbf{x}$  is called the **objective function**, the constraints  $\mathbf{x} \geq \mathbf{0}$  are called the **non negativity constraints**.

Note that the above problem can also be written as:

Max or Min  $\mathbf{c}^T \mathbf{x}$



Note that the above problem can also be written as:

Max or Min  $\mathbf{c}^T \mathbf{x}$

$\mathbf{a}_i^T \mathbf{x} \leq b_i$  for  $i = 1, 2, \dots, m$ ,

Note that the above problem can also be written as:

Max or Min  $\mathbf{c}^T \mathbf{x}$

$$\mathbf{a}_i^T \mathbf{x} \leq b_i \text{ for } i = 1, 2, \dots, m,$$

$$x_j \geq 0 \text{ for } j = 1, 2, \dots, n,$$

$$\text{or } -\mathbf{e}_j^T \mathbf{x} \leq 0 \text{ for } j = 1, 2, \dots, n,$$

Note that the above problem can also be written as:

Max or Min  $\mathbf{c}^T \mathbf{x}$

$\mathbf{a}_i^T \mathbf{x} \leq b_i$  for  $i = 1, 2, \dots, m$ ,

$x_j \geq 0$  for  $j = 1, 2, \dots, n$ ,

or  $-\mathbf{e}_j^T \mathbf{x} \leq 0$  for  $j = 1, 2, \dots, n$ ,

where  $\mathbf{a}_i^T$  is the  $i$  th row of the matrix  $\mathbf{A}$ , and  $\mathbf{e}_j$  is the  $j$  th column of the identity matrix of order  $n$ ,  $\mathbf{I}_n$ .

Note that the above problem can also be written as:

Max or Min  $\mathbf{c}^T \mathbf{x}$

$\mathbf{a}_i^T \mathbf{x} \leq b_i$  for  $i = 1, 2, \dots, m$ ,

$x_j \geq 0$  for  $j = 1, 2, \dots, n$ ,

or  $-\mathbf{e}_j^T \mathbf{x} \leq 0$  for  $j = 1, 2, \dots, n$ ,

where  $\mathbf{a}_i^T$  is the  $i$  th row of the matrix  $\mathbf{A}$ , and  $\mathbf{e}_j$  is the  $j$  th column of the identity matrix of order  $n$ ,  $\mathbf{I}_n$ .

Note that each of the functions

$\mathbf{a}_i^T \mathbf{x}$ , for  $i = 1, 2, \dots, m$ ,

Note that the above problem can also be written as:

Max or Min  $\mathbf{c}^T \mathbf{x}$

$\mathbf{a}_i^T \mathbf{x} \leq b_i$  for  $i = 1, 2, \dots, m$ ,

$x_j \geq 0$  for  $j = 1, 2, \dots, n$ ,

or  $-\mathbf{e}_j^T \mathbf{x} \leq 0$  for  $j = 1, 2, \dots, n$ ,

where  $\mathbf{a}_i^T$  is the  $i$ th row of the matrix  $\mathbf{A}$ , and  $\mathbf{e}_j$  is the  $j$ th column of the identity matrix of order  $n$ ,  $\mathbf{I}_n$ .

Note that each of the functions

$\mathbf{a}_i^T \mathbf{x}$ , for  $i = 1, 2, \dots, m$ ,

$-\mathbf{e}_j^T \mathbf{x}$ , for  $j = 1, 2, \dots, n$ ,

Note that the above problem can also be written as:

Max or Min  $\mathbf{c}^T \mathbf{x}$

$\mathbf{a}_i^T \mathbf{x} \leq b_i$  for  $i = 1, 2, \dots, m$ ,

$x_j \geq 0$  for  $j = 1, 2, \dots, n$ ,

or  $-\mathbf{e}_j^T \mathbf{x} \leq 0$  for  $j = 1, 2, \dots, n$ ,

where  $\mathbf{a}_i^T$  is the  $i$  th row of the matrix  $\mathbf{A}$ , and  $\mathbf{e}_j$  is the  $j$  th column of the identity matrix of order  $n$ ,  $\mathbf{I}_n$ .

Note that each of the functions

$\mathbf{a}_i^T \mathbf{x}$ , for  $i = 1, 2, \dots, m$ ,

$-\mathbf{e}_j^T \mathbf{x}$ , for  $j = 1, 2, \dots, n$ ,

and  $\mathbf{c}^T \mathbf{x}$  are all **linear functions** from  $\mathbb{R}^n \rightarrow \mathbb{R}$ , hence the name **linear programming problem**.

# Linear function, Feasible solution, Optimal Solution

- A function  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is called a **linear map** (linear function, linear transformation) if it satisfies the following:

# Linear function, Feasible solution, Optimal Solution

- A function  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is called a **linear map** (linear function, linear transformation) if it satisfies the following:

$$T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y}) \text{ for all } \mathbf{x}, \mathbf{y} \in \mathbb{R}^n,$$



# Linear function, Feasible solution, Optimal Solution

- A function  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is called a **linear map** (linear function, linear transformation) if it satisfies the following:

$$T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y}) \text{ for all } \mathbf{x}, \mathbf{y} \in \mathbb{R}^n,$$

$$\text{and } T(\alpha \mathbf{x}) = \alpha T(\mathbf{x}) \text{ for all } \alpha \in \mathbb{R} \text{ and all } \mathbf{x} \in \mathbb{R}^n.$$

# Linear function, Feasible solution, Optimal Solution

- A function  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is called a **linear map** (linear function, linear transformation) if it satisfies the following:  
 $T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y})$  for all  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ ,  
and  $T(\alpha \mathbf{x}) = \alpha T(\mathbf{x})$  for all  $\alpha \in \mathbb{R}$  and all  $\mathbf{x} \in \mathbb{R}^n$ .
- An  $\mathbf{x} \geq \mathbf{0}$  satisfying the constraints  $A\mathbf{x} \leq \mathbf{b}$  (or  $A\mathbf{x} \geq \mathbf{b}$ ) is called a **feasible solution** of the linear programming problem ( LPP ).

# Linear function, Feasible solution, Optimal Solution

- A function  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is called a **linear map** (linear function, linear transformation) if it satisfies the following:  
 $T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y})$  for all  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ ,  
and  $T(\alpha \mathbf{x}) = \alpha T(\mathbf{x})$  for all  $\alpha \in \mathbb{R}$  and all  $\mathbf{x} \in \mathbb{R}^n$ .
- An  $\mathbf{x} \geq \mathbf{0}$  satisfying the constraints  $A\mathbf{x} \leq \mathbf{b}$  (or  $A\mathbf{x} \geq \mathbf{b}$ ) is called a **feasible solution** of the linear programming problem ( LPP ).
- The set of all feasible solutions of a LPP is called the **feasible region** of the LPP.

# Linear function, Feasible solution, Optimal Solution

- A function  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is called a **linear map** (linear function, linear transformation) if it satisfies the following:

$$T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y}) \text{ for all } \mathbf{x}, \mathbf{y} \in \mathbb{R}^n,$$

$$\text{and } T(\alpha \mathbf{x}) = \alpha T(\mathbf{x}) \text{ for all } \alpha \in \mathbb{R} \text{ and all } \mathbf{x} \in \mathbb{R}^n.$$

- An  $\mathbf{x} \geq \mathbf{0}$  satisfying the constraints  $A\mathbf{x} \leq \mathbf{b}$  (or  $A\mathbf{x} \geq \mathbf{b}$ ) is called a **feasible solution** of the linear programming problem ( LPP ).
- The set of all feasible solutions of a LPP is called the **feasible region** of the LPP.

Hence the **feasible region** of a LPP, denoted by  $\text{Fea}(\text{LPP})$  is given by,

$$\text{Fea}(\text{LPP}) = \{ \mathbf{x} \in \mathbb{R}^n : \mathbf{x} \geq \mathbf{0}, A\mathbf{x} \leq \mathbf{b} \}.$$

# Linear function, Feasible solution, Optimal Solution

- A function  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is called a **linear map** (linear function, linear transformation) if it satisfies the following:

$$T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y}) \text{ for all } \mathbf{x}, \mathbf{y} \in \mathbb{R}^n,$$

$$\text{and } T(\alpha \mathbf{x}) = \alpha T(\mathbf{x}) \text{ for all } \alpha \in \mathbb{R} \text{ and all } \mathbf{x} \in \mathbb{R}^n.$$

- An  $\mathbf{x} \geq \mathbf{0}$  satisfying the constraints  $A\mathbf{x} \leq \mathbf{b}$  (or  $A\mathbf{x} \geq \mathbf{b}$ ) is called a **feasible solution** of the linear programming problem ( LPP ).
- The set of all feasible solutions of a LPP is called the **feasible region** of the LPP.

Hence the **feasible region** of a LPP, denoted by  $\text{Fea}(\text{LPP})$  is given by,

$$\text{Fea}(\text{LPP}) = \{ \mathbf{x} \in \mathbb{R}^n : \mathbf{x} \geq \mathbf{0}, A\mathbf{x} \leq \mathbf{b} \}.$$

- Fea(LPP) =  $\{\mathbf{x} \in \mathbb{R}^n : x_j \geq 0, j = 1, \dots, n, \mathbf{a}_i^T \mathbf{x} \leq b_i, i = 1, \dots, m\}$

- $\text{Fea(LPP)} = \{\mathbf{x} \in \mathbb{R}^n : x_j \geq 0, j = 1, \dots, n, \mathbf{a}_i^T \mathbf{x} \leq b_i, i = 1, \dots, m\}$   
 $= \{\mathbf{x} \in \mathbb{R}^n : -\mathbf{e}_j^T \mathbf{x} \leq 0, j = 1, \dots, n, \mathbf{a}_i^T \mathbf{x} \leq b_i, i = 1, \dots, m\},$

- $\text{Fea(LPP)} = \{\mathbf{x} \in \mathbb{R}^n : x_j \geq 0, j = 1, \dots, n, \mathbf{a}_i^T \mathbf{x} \leq b_i, i = 1, \dots, m\}$   
 $= \{\mathbf{x} \in \mathbb{R}^n : -\mathbf{e}_j^T \mathbf{x} \leq 0, j = 1, \dots, n, \mathbf{a}_i^T \mathbf{x} \leq b_i, i = 1, \dots, m\},$

where  $\mathbf{a}_i^T$  is the  $i$ th row of the matrix  $\mathbf{A}$ , and  $\mathbf{e}_j$  is the  $j$ th standard unit vector, or the  $j$ th column of the identity matrix  $I_n$ .



- $\text{Fea}(\text{LPP}) = \{\mathbf{x} \in \mathbb{R}^n : x_j \geq 0, j = 1, \dots, n, \mathbf{a}_i^T \mathbf{x} \leq b_i, i = 1, \dots, m\}$   
 $= \{\mathbf{x} \in \mathbb{R}^n : -\mathbf{e}_j^T \mathbf{x} \leq 0, j = 1, \dots, n, \mathbf{a}_i^T \mathbf{x} \leq b_i, i = 1, \dots, m\},$

where  $\mathbf{a}_i^T$  is the  $i$ th row of the matrix  $\mathbf{A}$ , and  $\mathbf{e}_j$  is the  $j$ th standard unit vector, or the  $j$ th column of the identity matrix  $I_n$ .

- A **feasible solution** of an LPP is called an **optimal solution** if it **minimizes** or **maximizes** the objective function (depending on the nature of the problem).

- $\text{Fea(LPP)} = \{\mathbf{x} \in \mathbb{R}^n : x_j \geq 0, j = 1, \dots, n, \mathbf{a}_i^T \mathbf{x} \leq b_i, i = 1, \dots, m\}$   
 $= \{\mathbf{x} \in \mathbb{R}^n : -\mathbf{e}_j^T \mathbf{x} \leq 0, j = 1, \dots, n, \mathbf{a}_i^T \mathbf{x} \leq b_i, i = 1, \dots, m\},$

where  $\mathbf{a}_i^T$  is the  $i$ th row of the matrix  $\mathbf{A}$ , and  $\mathbf{e}_j$  is the  $j$ th standard unit vector, or the  $j$ th column of the identity matrix  $I_n$ .

- A **feasible solution** of an LPP is called an **optimal solution** if it **minimizes** or **maximizes** the objective function (depending on the nature of the problem).
- If the LPP has an **optimal solution**, then the value of the objective function  $\mathbf{c}^T \mathbf{x}$  where  $\mathbf{x}$  is an **optimal solution** of the LPP is called the **optimal value** of the LPP.

- **Example 1:** Given the linear programming problem

Max  $5x + 2y$

subject to

$$3x + 2y \leq 6$$

- **Example 1:** Given the linear programming problem

Max  $5x + 2y$

subject to

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

- **Example 1:** Given the linear programming problem

Max  $5x + 2y$

subject to

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$

- **Example 1:** Given the linear programming problem

$$\text{Max } 5x + 2y$$

subject to

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$

$[3, 0]^T$  is not a feasible solution.

- **Example 1:** Given the linear programming problem

$$\text{Max } 5x + 2y$$

subject to

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$

$[3, 0]^T$  is not a feasible solution.

$[0, 0]^T, [1, 0]^T$  are feasible solutions which are **not** optimal.

- **Example 1:** Given the linear programming problem

$$\text{Max } 5x + 2y$$

subject to

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$

$[3, 0]^T$  is not a feasible solution.

$[0, 0]^T, [1, 0]^T$  are feasible solutions which are **not** optimal.

Optimal solution =  $[2, 0]^T$ , Optimal value=10.



- **Example 1:** Given the linear programming problem

$$\text{Max } 5x + 2y$$

subject to

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$

$[3, 0]^T$  is not a feasible solution.

$[0, 0]^T, [1, 0]^T$  are feasible solutions which are **not** optimal.

Optimal solution =  $[2, 0]^T$ , Optimal value=10.

- **Example 2:** Consider the problem,

$$\text{Min } -x + 2y,$$

- **Example 1:** Given the linear programming problem

$$\text{Max } 5x + 2y$$

subject to

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$

$[3, 0]^T$  is not a feasible solution.

$[0, 0]^T, [1, 0]^T$  are feasible solutions which are **not** optimal.

Optimal solution =  $[2, 0]^T$ , Optimal value=10.

- **Example 2:** Consider the problem,

$$\text{Min } -x + 2y,$$

subject to

- **Example 1:** Given the linear programming problem

$$\text{Max } 5x + 2y$$

subject to

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$

$[3, 0]^T$  is not a feasible solution.

$[0, 0]^T, [1, 0]^T$  are feasible solutions which are **not** optimal.

Optimal solution =  $[2, 0]^T$ , Optimal value=10.

- **Example 2:** Consider the problem,

$$\text{Min } -x + 2y,$$

subject to

$$x + 2y \geq 1,$$

- **Example 1:** Given the linear programming problem

$$\text{Max } 5x + 2y$$

subject to

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$

$[3, 0]^T$  is not a feasible solution.

$[0, 0]^T, [1, 0]^T$  are feasible solutions which are **not** optimal.

Optimal solution =  $[2, 0]^T$ , Optimal value=10.

- **Example 2:** Consider the problem,

$$\text{Min } -x + 2y,$$

subject to

$$x + 2y \geq 1, -x + y \leq 1,$$

- **Example 1:** Given the linear programming problem

$$\text{Max } 5x + 2y$$

subject to

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$

$[3, 0]^T$  is not a feasible solution.

$[0, 0]^T, [1, 0]^T$  are feasible solutions which are **not** optimal.

Optimal solution =  $[2, 0]^T$ , Optimal value=10.

- **Example 2:** Consider the problem,

$$\text{Min } -x + 2y,$$

subject to

$$x + 2y \geq 1, -x + y \leq 1, 2x + 4y \leq 4,$$

- **Example 1:** Given the linear programming problem

$$\text{Max } 5x + 2y$$

subject to

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$

$[3, 0]^T$  is not a feasible solution.

$[0, 0]^T, [1, 0]^T$  are feasible solutions which are **not** optimal.

Optimal solution =  $[2, 0]^T$ , Optimal value=10.

- **Example 2:** Consider the problem,

$$\text{Min } -x + 2y,$$

subject to

$$x + 2y \geq 1, -x + y \leq 1, 2x + 4y \leq 4, \quad x \geq 0, y \geq 0.$$

- **Example 1:** Given the linear programming problem

$$\text{Max } 5x + 2y$$

subject to

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$

$[3, 0]^T$  is not a feasible solution.

$[0, 0]^T, [1, 0]^T$  are feasible solutions which are **not** optimal.

Optimal solution =  $[2, 0]^T$ , Optimal value=10.

- **Example 2:** Consider the problem,

$$\text{Min } -x + 2y,$$

subject to

$$x + 2y \geq 1, -x + y \leq 1, 2x + 4y \leq 4, \quad x \geq 0, y \geq 0.$$

$[0, 0]^T$  is not a feasible solution.

- **Example 1:** Given the linear programming problem

$$\text{Max } 5x + 2y$$

subject to

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$

$[3, 0]^T$  is not a feasible solution.

$[0, 0]^T, [1, 0]^T$  are feasible solutions which are **not** optimal.

Optimal solution =  $[2, 0]^T$ , Optimal value=10.

- **Example 2:** Consider the problem,

$$\text{Min } -x + 2y,$$

subject to

$$x + 2y \geq 1, -x + y \leq 1, 2x + 4y \leq 4, \quad x \geq 0, y \geq 0.$$

$[0, 0]^T$  is not a feasible solution.

$[1, 0]^T, [0, 1]^T$  are feasible solutions which are **not** optimal.



- **Example 1:** Given the linear programming problem

$$\text{Max } 5x + 2y$$

subject to

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$

$[3, 0]^T$  is not a feasible solution.

$[0, 0]^T, [1, 0]^T$  are feasible solutions which are **not** optimal.

Optimal solution =  $[2, 0]^T$ , Optimal value=10.

- **Example 2:** Consider the problem,

$$\text{Min } -x + 2y,$$

subject to

$$x + 2y \geq 1, -x + y \leq 1, 2x + 4y \leq 4, \quad x \geq 0, y \geq 0.$$

$[0, 0]^T$  is not a feasible solution.

$[1, 0]^T, [0, 1]^T$  are feasible solutions which are **not** optimal.

Optimal solution =  $[2, 0]^T$ , Optimal value=-2.

# Hyperplanes, Normals, Closed Half Spaces, Polyhedral set

- A subset  $H$  of  $\mathbb{R}^n$  is called a **hyperplane** if it can be written as:

$H = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T \mathbf{x} = d\}$  for some  $\mathbf{a} \in \mathbb{R}^n$  and  $d \in \mathbb{R}$ , or equivalently as

# Hyperplanes, Normals, Closed Half Spaces, Polyhedral set

- A subset  $H$  of  $\mathbb{R}^n$  is called a **hyperplane** if it can be written as:

$H = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T \mathbf{x} = d\}$  for some  $\mathbf{a} \in \mathbb{R}^n$  and  $d \in \mathbb{R}$ , or equivalently as

$H = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T (\mathbf{x} - \mathbf{x}_0) = 0\}$  for some  $\mathbf{a} \in \mathbb{R}^n$ ,  $d \in \mathbb{R}$ , and  $\mathbf{x}_0$  satisfying  $\mathbf{a}^T \mathbf{x}_0 = d$ .

# Hyperplanes, Normals, Closed Half Spaces, Polyhedral set

- A subset  $H$  of  $\mathbb{R}^n$  is called a **hyperplane** if it can be written as:

$H = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T \mathbf{x} = d\}$  for some  $\mathbf{a} \in \mathbb{R}^n$  and  $d \in \mathbb{R}$ , or equivalently as

$H = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T (\mathbf{x} - \mathbf{x}_0) = 0\}$  for some  $\mathbf{a} \in \mathbb{R}^n$ ,  $d \in \mathbb{R}$ , and  $\mathbf{x}_0$  satisfying  $\mathbf{a}^T \mathbf{x}_0 = d$ .

- So geometrically a **hyperplane** in  $\mathbb{R}$  is just an element of  $\mathbb{R}$  (a **single point**),

# Hyperplanes, Normals, Closed Half Spaces, Polyhedral set

- A subset  $H$  of  $\mathbb{R}^n$  is called a **hyperplane** if it can be written as:

$H = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T \mathbf{x} = d\}$  for some  $\mathbf{a} \in \mathbb{R}^n$  and  $d \in \mathbb{R}$ , or equivalently as

$H = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T (\mathbf{x} - \mathbf{x}_0) = 0\}$  for some  $\mathbf{a} \in \mathbb{R}^n$ ,  $d \in \mathbb{R}$ , and  $\mathbf{x}_0$  satisfying  $\mathbf{a}^T \mathbf{x}_0 = d$ .

- So geometrically a **hyperplane** in  $\mathbb{R}$  is just an element of  $\mathbb{R}$  (a **single point**), in  $\mathbb{R}^2$  it is just a **straight line**,

# Hyperplanes, Normals, Closed Half Spaces, Polyhedral set

- A subset  $H$  of  $\mathbb{R}^n$  is called a **hyperplane** if it can be written as:  
 $H = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T \mathbf{x} = d\}$  for some  $\mathbf{a} \in \mathbb{R}^n$  and  $d \in \mathbb{R}$ , or equivalently as  
 $H = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T (\mathbf{x} - \mathbf{x}_0) = 0\}$  for some  $\mathbf{a} \in \mathbb{R}^n$ ,  $d \in \mathbb{R}$ , and  $\mathbf{x}_0$  satisfying  $\mathbf{a}^T \mathbf{x}_0 = d$ .
- So geometrically a **hyperplane** in  $\mathbb{R}$  is just an element of  $\mathbb{R}$  (a **single point**), in  $\mathbb{R}^2$  it is just a **straight line**, in  $\mathbb{R}^3$  it is just the usual **plane** we are familiar with.

# Hyperplanes, Normals, Closed Half Spaces, Polyhedral set

- A subset  $H$  of  $\mathbb{R}^n$  is called a **hyperplane** if it can be written as:  
 $H = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T \mathbf{x} = d\}$  for some  $\mathbf{a} \in \mathbb{R}^n$  and  $d \in \mathbb{R}$ , or equivalently as  
 $H = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T (\mathbf{x} - \mathbf{x}_0) = 0\}$  for some  $\mathbf{a} \in \mathbb{R}^n$ ,  $d \in \mathbb{R}$ , and  $\mathbf{x}_0$  satisfying  $\mathbf{a}^T \mathbf{x}_0 = d$ .
- So geometrically a **hyperplane** in  $\mathbb{R}$  is just an element of  $\mathbb{R}$  (a **single point**), in  $\mathbb{R}^2$  it is just a **straight line**, in  $\mathbb{R}^3$  it is just the usual **plane** we are familiar with.
- The vector  $\mathbf{a}$  is called a **normal** to the hyperplane  $H$ , since it is **orthogonal** (or perpendicular) to each of the vectors  $\mathbf{x} - \mathbf{x}_0$  on the hyperplane with tail at  $\mathbf{x}_0$ .

- A collection of hyperplanes  $H_1, \dots, H_k$  in  $\mathbb{R}^n$  is said to be **Linearly Independent (LI)** if the corresponding normal vectors  $\mathbf{a}_1, \dots, \mathbf{a}_k$  are linearly independent as vectors in  $\mathbb{R}^n$ .



- A collection of hyperplanes  $H_1, \dots, H_k$  in  $\mathbb{R}^n$  is said to be **Linearly Independent (LI)** if the corresponding normal vectors  $\mathbf{a}_1, \dots, \mathbf{a}_k$  are linearly independent as vectors in  $\mathbb{R}^n$ .
- Otherwise the collection of hyperplanes is said to be **Linearly Dependent (LD)**.

- A collection of hyperplanes  $H_1, \dots, H_k$  in  $\mathbb{R}^n$  is said to be **Linearly Independent (LI)** if the corresponding normal vectors  $\mathbf{a}_1, \dots, \mathbf{a}_k$  are linearly independent as vectors in  $\mathbb{R}^n$ .
- Otherwise the collection of hyperplanes is said to be **Linearly Dependent (LD)**.
- **Definition:** Vectors  $\mathbf{a}_1, \dots, \mathbf{a}_k$  in  $\mathbb{R}^n$  are said to be **LD** if there exists real numbers  $c_1, \dots, c_k$ , not all zeros such that  $c_1 \mathbf{a}_1 + \dots + c_k \mathbf{a}_k = \mathbf{O}$ ,  $(^{**})$  where  $\mathbf{O}$  is the zero vector.

- A collection of hyperplanes  $H_1, \dots, H_k$  in  $\mathbb{R}^n$  is said to be **Linearly Independent (LI)** if the corresponding normal vectors  $\mathbf{a}_1, \dots, \mathbf{a}_k$  are linearly independent as vectors in  $\mathbb{R}^n$ .
- Otherwise the collection of hyperplanes is said to be **Linearly Dependent (LD)**.
- **Definition:** Vectors  $\mathbf{a}_1, \dots, \mathbf{a}_k$  in  $\mathbb{R}^n$  are said to be **LD** if there exists real numbers  $c_1, \dots, c_k$ , not all zeros such that  $c_1 \mathbf{a}_1 + \dots + c_k \mathbf{a}_k = \mathbf{O}$ ,  $(^{**})$  where  $\mathbf{O}$  is the zero vector.
- If they are not **LD** then the vectors are called **LI**, that is, the only solution to  $(^{**})$  is  $c_1 = \dots = c_k = 0$ .

- For  $k = 2$ ,  $\mathbf{a}_1, \mathbf{a}_2$  is **LD** if and only if either  $c_1 \neq 0$  or  $c_2 \neq 0$ ,  
 $\mathbf{a}_1 = -\frac{c_2}{c_1}\mathbf{a}_2$  (if  $c_1 \neq 0$ )

- For  $k = 2$ ,  $\mathbf{a}_1, \mathbf{a}_2$  is **LD** if and only if either  $c_1 \neq 0$  or  $c_2 \neq 0$ ,  
 $\mathbf{a}_1 = -\frac{c_2}{c_1}\mathbf{a}_2$  (if  $c_1 \neq 0$ )       $\mathbf{a}_2 = -\frac{c_1}{c_2}\mathbf{a}_1$  (if  $c_2 \neq 0$ ).  
[.1cm]

- For  $k = 2$ ,  $\mathbf{a}_1, \mathbf{a}_2$  is **LD** if and only if either  $c_1 \neq 0$  or  $c_2 \neq 0$ ,  
 $\mathbf{a}_1 = -\frac{c_2}{c_1}\mathbf{a}_2$  (if  $c_1 \neq 0$ )       $\mathbf{a}_2 = -\frac{c_1}{c_2}\mathbf{a}_1$  (if  $c_2 \neq 0$ ).  
 [.1cm]
- Any set of vectors containing the **zero vector** is **LD**.

- For  $k = 2$ ,  $\mathbf{a}_1, \mathbf{a}_2$  is **LD** if and only if either  $c_1 \neq 0$  or  $c_2 \neq 0$ ,  
 $\mathbf{a}_1 = -\frac{c_2}{c_1}\mathbf{a}_2$  (if  $c_1 \neq 0$ )       $\mathbf{a}_2 = -\frac{c_1}{c_2}\mathbf{a}_1$  (if  $c_2 \neq 0$ ).  
 [.1cm]
- Any set of vectors containing the **zero vector** is **LD**.  
 For example if say  $\mathbf{a}_1 = \mathbf{O}$  then  
 $1\mathbf{a}_1 + 0\mathbf{a}_2 \dots + 0\mathbf{a}_k = \mathbf{O}$ ,

- For  $k = 2$ ,  $\mathbf{a}_1, \mathbf{a}_2$  is **LD** if and only if either  $c_1 \neq 0$  or  $c_2 \neq 0$ ,  
 $\mathbf{a}_1 = -\frac{c_2}{c_1}\mathbf{a}_2$  (if  $c_1 \neq 0$ )       $\mathbf{a}_2 = -\frac{c_1}{c_2}\mathbf{a}_1$  (if  $c_2 \neq 0$ ).  
 [.1cm]
- Any set of vectors containing the **zero vector** is **LD**.  
 For example if say  $\mathbf{a}_1 = \mathbf{O}$  then  
 $1\mathbf{a}_1 + 0\mathbf{a}_2 \dots + 0\mathbf{a}_k = \mathbf{O}$ ,  
 there is a solution to  $c_1\mathbf{a}_1 + \dots + c_k\mathbf{a}_k = \mathbf{O}$ , with  
 $c_1 = 1, c_2 = 0, \dots, c_k = 0$  not all of which are zeros.



- **Example 2 revisited:** Consider the problem,  
Min  $-x + 2y$ ,

- **Example 2 revisited:** Consider the problem,  
Min  $-x + 2y$ ,  
subject to

- **Example 2 revisited:** Consider the problem,  
Min  $-x + 2y$ ,  
subject to  
 $x + 2y \geq 1$ ,

- **Example 2 revisited:** Consider the problem,

Min  $-x + 2y$ ,

subject to

$$x + 2y \geq 1, \quad -x + y \leq 1,$$

- **Example 2 revisited:** Consider the problem,

Min  $-x + 2y$ ,

subject to

$$x + 2y \geq 1, \quad -x + y \leq 1, \quad 2x + 4y \leq 4,$$

- **Example 2 revisited:** Consider the problem,

Min  $-x + 2y$ ,

subject to

$$x + 2y \geq 1, \quad -x + y \leq 1, \quad 2x + 4y \leq 4, \quad x \geq 0, y \geq 0.$$

- **Example 2 revisited:** Consider the problem,

Min  $-x + 2y$ ,

subject to

$$x + 2y \geq 1, -x + y \leq 1, 2x + 4y \leq 4, \quad x \geq 0, y \geq 0.$$

$H_1 = \{[x, y]^T : x = 0\}$  ,  $H_2 = \{[x, y]^T : y = 0\}$  is **LI**.

- **Example 2 revisited:** Consider the problem,

Min  $-x + 2y$ ,

subject to

$$x + 2y \geq 1, -x + y \leq 1, 2x + 4y \leq 4, \quad x \geq 0, y \geq 0.$$

$$H_1 = \{[x, y]^T : x = 0\}, H_2 = \{[x, y]^T : y = 0\} \text{ is LI.}$$

$$H_1 = \{[x, y]^T : x = 0\}, H_3 = \{[x, y]^T : x + 2y = 1\} \text{ is LI.}$$



- **Example 2 revisited:** Consider the problem,

Min  $-x + 2y$ ,

subject to

$$x + 2y \geq 1, -x + y \leq 1, 2x + 4y \leq 4, \quad x \geq 0, y \geq 0.$$

$$H_1 = \{[x, y]^T : x = 0\}, H_2 = \{[x, y]^T : y = 0\} \text{ is LI.}$$

$$H_1 = \{[x, y]^T : x = 0\}, H_3 = \{[x, y]^T : x + 2y = 1\} \text{ is LI.}$$

- A set of normals to  $H_1$  is  $\{[1, 0]^T, [-1, 0]^T\}$  whereas a set of normals to  $H_2$  is  $\{[0, 1]^T, [0, -1]^T\}$  and  $\{[1, 0]^T, [0, 1]^T\}$  is LI.

- **Example 2 revisited:** Consider the problem,

Min  $-x + 2y$ ,

subject to

$$x + 2y \geq 1, -x + y \leq 1, 2x + 4y \leq 4, \quad x \geq 0, y \geq 0.$$

$$H_1 = \{[x, y]^T : x = 0\}, H_2 = \{[x, y]^T : y = 0\} \text{ is LI.}$$

$$H_1 = \{[x, y]^T : x = 0\}, H_3 = \{[x, y]^T : x + 2y = 1\} \text{ is LI.}$$

- A set of normals to  $H_1$  is  $\{[1, 0]^T, [-1, 0]^T\}$  whereas a set of normals to  $H_2$  is  $\{[0, 1]^T, [0, -1]^T\}$  and  $\{[1, 0]^T, [0, 1]^T\}$  is LI.

$$H_3 = \{[x, y]^T : x + 2y = 1\}, H_4 = \{[x, y]^T : 2x + 4y = 4\} \text{ is LD.}$$

- **Example 2 revisited:** Consider the problem,

Min  $-x + 2y$ ,

subject to

$$x + 2y \geq 1, -x + y \leq 1, 2x + 4y \leq 4, \quad x \geq 0, y \geq 0.$$

$H_1 = \{[x, y]^T : x = 0\}$  ,  $H_2 = \{[x, y]^T : y = 0\}$  is **LI**.

$H_1 = \{[x, y]^T : x = 0\}$  ,  $H_3 = \{[x, y]^T : x + 2y = 1\}$  is **LI**.

- A set of normals to  $H_1$  is  $\{[1, 0]^T, [-1, 0]^T\}$  whereas a set of normals to  $H_2$  is  $\{[0, 1]^T, [0, -1]^T\}$  and  $\{[1, 0]^T, [0, 1]^T\}$  is **LI**.

$H_3 = \{[x, y]^T : x + 2y = 1\}$  ,  $H_4 = \{[x, y]^T : 2x + 4y = 4\}$  is **LD**.

- A set of normals to  $H_3$  is  $\{[1, 2]^T, [-1, -2]^T\}$  whereas a set of normals to  $H_2$  is  $\{[2, 4]^T, [-2, -4]^T\}$  and  $\{[1, 2]^T, [2, 4]^T\}$  is **LD** since  $2[1, 2]^T - 1[2, 4]^T = \mathbf{0}$ .

- **Example 2 revisited:** Consider the problem,

Min  $-x + 2y$ ,

subject to

$$x + 2y \geq 1, -x + y \leq 1, 2x + 4y \leq 4, \quad x \geq 0, y \geq 0.$$

$$H_1 = \{[x, y]^T : x = 0\}, H_2 = \{[x, y]^T : y = 0\} \text{ is LI.}$$

$$H_1 = \{[x, y]^T : x = 0\}, H_3 = \{[x, y]^T : x + 2y = 1\} \text{ is LI.}$$

- A set of normals to  $H_1$  is  $\{[1, 0]^T, [-1, 0]^T\}$  whereas a set of normals to  $H_2$  is  $\{[0, 1]^T, [0, -1]^T\}$  and  $\{[1, 0]^T, [0, 1]^T\}$  is LI.

$$H_3 = \{[x, y]^T : x + 2y = 1\}, H_4 = \{[x, y]^T : 2x + 4y = 4\} \text{ is LD.}$$

- A set of normals to  $H_3$  is  $\{[1, 2]^T, [-1, -2]^T\}$  whereas a set of normals to  $H_2$  is  $\{[2, 4]^T, [-2, -4]^T\}$  and  $\{[1, 2]^T, [2, 4]^T\}$  is LD since  $2[1, 2]^T - 1[2, 4]^T = \mathbf{0}$ .
- What about  $H_1, H_2, H_3$ ?

- Any set of  $(n + 1)$  hyperplanes in  $\mathbb{R}^n$  is LD.

- Any set of  $(n + 1)$  hyperplanes in  $\mathbb{R}^n$  is LD.
- Associated with a hyperplane  $H$  are two closed half spaces

- Any set of  $(n + 1)$  hyperplanes in  $\mathbb{R}^n$  is **LD**.
- Associated with a hyperplane  $H$  are two **closed half spaces**  
 $H_1 = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T \mathbf{x} \leq d\}$  and

- Any set of  $(n + 1)$  hyperplanes in  $\mathbb{R}^n$  is LD.
- Associated with a hyperplane  $H$  are two closed half spaces

$$H_1 = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T \mathbf{x} \leq d\} \text{ and}$$

$$H_2 = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T \mathbf{x} \geq d\}.$$



- Any set of  $(n + 1)$  hyperplanes in  $\mathbb{R}^n$  is LD.
- Associated with a hyperplane  $H$  are two closed half spaces

$$H_1 = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T \mathbf{x} \leq d\} \text{ and}$$

$$H_2 = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T \mathbf{x} \geq d\}.$$

Note that the hyperplane  $H$ , and the two half spaces  $H_1, H_2$  are all closed subsets of  $\mathbb{R}^n$

- Any set of  $(n + 1)$  hyperplanes in  $\mathbb{R}^n$  is LD.
- Associated with a hyperplane  $H$  are two closed half spaces

$$H_1 = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T \mathbf{x} \leq d\} \text{ and}$$

$$H_2 = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T \mathbf{x} \geq d\}.$$

Note that the hyperplane  $H$ , and the two half spaces  $H_1, H_2$  are all closed subsets of  $\mathbb{R}^n$  (since each of these sets contains all its boundary points, the boundary points being  $\mathbf{x} \in \mathbb{R}^n$  satisfying  $\mathbf{a}^T \mathbf{x} = d$ ).

- Any set of  $(n + 1)$  hyperplanes in  $\mathbb{R}^n$  is **LD**.
- Associated with a hyperplane  $H$  are two **closed half spaces**

$$H_1 = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T \mathbf{x} \leq d\} \text{ and}$$

$$H_2 = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T \mathbf{x} \geq d\}.$$

Note that the **hyperplane**  $H$ , and the two **half spaces**  $H_1, H_2$  are all **closed subsets** of  $\mathbb{R}^n$  (since each of these sets contains all its boundary points, the boundary points being  $\mathbf{x} \in \mathbb{R}^n$  satisfying  $\mathbf{a}^T \mathbf{x} = d$ ).

For example for the hyperplane  $H = \{[x, y]^T : x + 2y = 1\}$ , as well as the half spaces  $H_1 = \{[x, y]^T : x + 2y \leq 1\}$  and  $H_2 = \{[x, y]^T : x + 2y \geq 1\}$ , the boundary points are

- Any set of  $(n + 1)$  hyperplanes in  $\mathbb{R}^n$  is **LD**.
- Associated with a hyperplane  $H$  are two **closed half spaces**

$$H_1 = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T \mathbf{x} \leq d\} \text{ and}$$

$$H_2 = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^T \mathbf{x} \geq d\}.$$

Note that the **hyperplane**  $H$ , and the two **half spaces**  $H_1, H_2$  are all **closed subsets** of  $\mathbb{R}^n$  (since each of these sets contains all its boundary points, the boundary points being  $\mathbf{x} \in \mathbb{R}^n$  satisfying  $\mathbf{a}^T \mathbf{x} = d$ ).

For example for the hyperplane  $H = \{[x, y]^T : x + 2y = 1\}$ , as well as the half spaces  $H_1 = \{[x, y]^T : x + 2y \leq 1\}$  and  $H_2 = \{[x, y]^T : x + 2y \geq 1\}$ , the boundary points are  $\{[x, y]^T : x + 2y = 1\}$ .

- A set which is the intersection of a **finite number** of **closed half spaces** is called a **polyhedral set**.

- A set which is the intersection of a **finite number** of **closed half spaces** is called a **polyhedral set**.

For example the **feasible region** in **Example 2** is

- A set which is the intersection of a **finite number** of **closed half spaces** is called a **polyhedral set**.

For example the **feasible region** in **Example 2** is

$S = \{[x, y]^T : x \geq 0, y \geq 0, 2x + 4y \leq 4, -x + y \leq 1, x + 2y \geq 1\}$  which is the intersection of the closed half spaces

- A set which is the intersection of a **finite number** of **closed half spaces** is called a **polyhedral set**.

For example the **feasible region** in **Example 2** is

$S = \{[x, y]^T : x \geq 0, y \geq 0, 2x + 4y \leq 4, -x + y \leq 1, x + 2y \geq 1\}$  which is the intersection of the closed half spaces

$$H_1 = \{[x, y]^T : x \geq 0\},$$



- A set which is the intersection of a **finite number** of **closed half spaces** is called a **polyhedral set**.

For example the **feasible region** in **Example 2** is

$S = \{[x, y]^T : x \geq 0, y \geq 0, 2x + 4y \leq 4, -x + y \leq 1, x + 2y \geq 1\}$  which is the intersection of the closed half spaces

$H_1 = \{[x, y]^T : x \geq 0\}, H_2 = \{[x, y]^T : y \geq 0\},$

- A set which is the intersection of a **finite number** of **closed half spaces** is called a **polyhedral set**.

For example the **feasible region** in **Example 2** is

$S = \{[x, y]^T : x \geq 0, y \geq 0, 2x + 4y \leq 4, -x + y \leq 1, x + 2y \geq 1\}$  which is the intersection of the closed half spaces

$$H_1 = \{[x, y]^T : x \geq 0\}, H_2 = \{[x, y]^T : y \geq 0\},$$

$$H_3 = \{[x, y]^T : x + 2y \geq 1\},$$

- A set which is the intersection of a **finite number** of **closed half spaces** is called a **polyhedral set**.

For example the **feasible region** in **Example 2** is

$S = \{[x, y]^T : x \geq 0, y \geq 0, 2x + 4y \leq 4, -x + y \leq 1, x + 2y \geq 1\}$  which is the intersection of the closed half spaces

$$H_1 = \{[x, y]^T : x \geq 0\}, H_2 = \{[x, y]^T : y \geq 0\},$$

$$H_3 = \{[x, y]^T : x + 2y \geq 1\}, H_4 = \{[x, y]^T : 2x + 4y \leq 4\}$$

and

- A set which is the intersection of a **finite number** of **closed half spaces** is called a **polyhedral set**.

For example the **feasible region** in **Example 2** is

$S = \{[x, y]^T : x \geq 0, y \geq 0, 2x + 4y \leq 4, -x + y \leq 1, x + 2y \geq 1\}$  which is the intersection of the closed half spaces

$H_1 = \{[x, y]^T : x \geq 0\}$ ,  $H_2 = \{[x, y]^T : y \geq 0\}$ ,

$H_3 = \{[x, y]^T : x + 2y \geq 1\}$ ,  $H_4 = \{[x, y]^T : 2x + 4y \leq 4\}$

and  $H_5 = \{[x, y]^T : -x + y \leq 1\}$

is a **polyhedral set**.

- A set which is the intersection of a **finite number** of **closed half spaces** is called a **polyhedral set**.

For example the **feasible region** in **Example 2** is

$S = \{[x, y]^T : x \geq 0, y \geq 0, 2x + 4y \leq 4, -x + y \leq 1, x + 2y \geq 1\}$  which is the intersection of the closed half spaces

$H_1 = \{[x, y]^T : x \geq 0\}$ ,  $H_2 = \{[x, y]^T : y \geq 0\}$ ,

$H_3 = \{[x, y]^T : x + 2y \geq 1\}$ ,  $H_4 = \{[x, y]^T : 2x + 4y \leq 4\}$

and  $H_5 = \{[x, y]^T : -x + y \leq 1\}$

is a **polyhedral set**.

- The feasible region of a LPP is a **polyhedral set**.

- A set which is the intersection of a **finite number** of **closed half spaces** is called a **polyhedral set**.

For example the **feasible region** in **Example 2** is

$S = \{[x, y]^T : x \geq 0, y \geq 0, 2x + 4y \leq 4, -x + y \leq 1, x + 2y \geq 1\}$  which is the intersection of the closed half spaces

$H_1 = \{[x, y]^T : x \geq 0\}$ ,  $H_2 = \{[x, y]^T : y \geq 0\}$ ,

$H_3 = \{[x, y]^T : x + 2y \geq 1\}$ ,  $H_4 = \{[x, y]^T : 2x + 4y \leq 4\}$

and  $H_5 = \{[x, y]^T : -x + y \leq 1\}$

is a **polyhedral set**.

- The feasible region of a LPP is a **polyhedral set**.
- Since the intersection of any collection of **closed subsets** of  $\mathbb{R}^n$  is again a **closed subset** of  $\mathbb{R}^n$ , hence **Fea(LPP)** is a **closed** subset of  $\mathbb{R}^n$ , geometrically the **feasible region** of a LPP contains all its boundary points.

- The hyperplanes  $\mathbf{a}_i^T \mathbf{x} = b_i$  for  $i = 1, 2, \dots, m$ ,

- The hyperplanes  $\mathbf{a}_i^T \mathbf{x} = b_i$  for  $i = 1, 2, \dots, m$ ,  
 $x_j = 0$  for  $j = 1, 2, \dots, n$ , or  $-\mathbf{e}_j^T \mathbf{x} = 0$  for  $j = 1, 2, \dots, n$ ,



- The hyperplanes  $\mathbf{a}_i^T \mathbf{x} = b_i$  for  $i = 1, 2, \dots, m$ ,  
 $x_j = 0$  for  $j = 1, 2, \dots, n$ , or  $-\mathbf{e}_j^T \mathbf{x} = 0$  for  $j = 1, 2, \dots, n$ ,  
associated with an LPP are called its **defining hyperplanes**.

- The hyperplanes  $\mathbf{a}_i^T \mathbf{x} = b_i$  for  $i = 1, 2, \dots, m$ ,  
 $x_j = 0$  for  $j = 1, 2, \dots, n$ , or  $-\mathbf{e}_j^T \mathbf{x} = 0$  for  $j = 1, 2, \dots, n$ ,  
 associated with an LPP are called its **defining hyperplanes**.
- The associated **half spaces** are  
 $\mathbf{a}_i^T \mathbf{x} \leq b_i$ ,  $i = 1, 2, \dots, m$ ,

- The hyperplanes  $\mathbf{a}_i^T \mathbf{x} = b_i$  for  $i = 1, 2, \dots, m$ ,  
 $x_j = 0$  for  $j = 1, 2, \dots, n$ , or  $-\mathbf{e}_j^T \mathbf{x} = 0$  for  $j = 1, 2, \dots, n$ ,  
 associated with an LPP are called its **defining hyperplanes**.
- The associated **half spaces** are  
 $\mathbf{a}_i^T \mathbf{x} \leq b_i$ ,  $i = 1, 2, \dots, m$ ,  
 $x_j \geq 0$ ,  $j = 1, 2, \dots, n$ , or  $-\mathbf{e}_j^T \mathbf{x} \leq 0$ ,  $j = 1, 2, \dots, n$ .

- The hyperplanes  $\mathbf{a}_i^T \mathbf{x} = b_i$  for  $i = 1, 2, \dots, m$ ,  
 $x_j = 0$  for  $j = 1, 2, \dots, n$ , or  $-\mathbf{e}_j^T \mathbf{x} = 0$  for  $j = 1, 2, \dots, n$ ,  
 associated with an LPP are called its **defining hyperplanes**.
- The associated **half spaces** are  
 $\mathbf{a}_i^T \mathbf{x} \leq b_i$ ,  $i = 1, 2, \dots, m$ ,  
 $x_j \geq 0$ ,  $j = 1, 2, \dots, n$ , or  $-\mathbf{e}_j^T \mathbf{x} \leq 0$ ,  $j = 1, 2, \dots, n$ .

For example the **defining hyperplanes** in **Example 2** are  
 $H_1 = \{[x, y]^T : x = 0\}$ ,

- The hyperplanes  $\mathbf{a}_i^T \mathbf{x} = b_i$  for  $i = 1, 2, \dots, m$ ,  
 $x_j = 0$  for  $j = 1, 2, \dots, n$ , or  $-\mathbf{e}_j^T \mathbf{x} = 0$  for  $j = 1, 2, \dots, n$ ,  
 associated with an LPP are called its **defining hyperplanes**.
- The associated **half spaces** are  
 $\mathbf{a}_i^T \mathbf{x} \leq b_i$ ,  $i = 1, 2, \dots, m$ ,  
 $x_j \geq 0$ ,  $j = 1, 2, \dots, n$ , or  $-\mathbf{e}_j^T \mathbf{x} \leq 0$ ,  $j = 1, 2, \dots, n$ .  
 For example the **defining hyperplanes** in **Example 2** are  
 $H_1 = \{[x, y]^T : x = 0\}$ ,  $H_2 = \{[x, y]^T : y = 0\}$ ,

- The hyperplanes  $\mathbf{a}_i^T \mathbf{x} = b_i$  for  $i = 1, 2, \dots, m$ ,  
 $x_j = 0$  for  $j = 1, 2, \dots, n$ , or  $-\mathbf{e}_j^T \mathbf{x} = 0$  for  $j = 1, 2, \dots, n$ ,  
 associated with an LPP are called its **defining hyperplanes**.
- The associated **half spaces** are  
 $\mathbf{a}_i^T \mathbf{x} \leq b_i$ ,  $i = 1, 2, \dots, m$ ,  
 $x_j \geq 0$ ,  $j = 1, 2, \dots, n$ , or  $-\mathbf{e}_j^T \mathbf{x} \leq 0$ ,  $j = 1, 2, \dots, n$ .

For example the **defining hyperplanes** in **Example 2** are

$$H_1 = \{[x, y]^T : x = 0\}, H_2 = \{[x, y]^T : y = 0\},$$

$$H_3 = \{[x, y]^T : x + 2y = 1\},$$

- The hyperplanes  $\mathbf{a}_i^T \mathbf{x} = b_i$  for  $i = 1, 2, \dots, m$ ,  
 $x_j = 0$  for  $j = 1, 2, \dots, n$ , or  $-\mathbf{e}_j^T \mathbf{x} = 0$  for  $j = 1, 2, \dots, n$ ,  
 associated with an LPP are called its **defining hyperplanes**.
- The associated **half spaces** are  
 $\mathbf{a}_i^T \mathbf{x} \leq b_i$ ,  $i = 1, 2, \dots, m$ ,  
 $x_j \geq 0$ ,  $j = 1, 2, \dots, n$ , or  $-\mathbf{e}_j^T \mathbf{x} \leq 0$ ,  $j = 1, 2, \dots, n$ .

For example the **defining hyperplanes** in **Example 2** are  
 $H_1 = \{[x, y]^T : x = 0\}$ ,  $H_2 = \{[x, y]^T : y = 0\}$ ,  
 $H_3 = \{[x, y]^T : x + 2y = 1\}$ ,  $H_4 = \{[x, y]^T : 2x + 4y = 4\}$   
 and

- The hyperplanes  $\mathbf{a}_i^T \mathbf{x} = b_i$  for  $i = 1, 2, \dots, m$ ,  
 $x_j = 0$  for  $j = 1, 2, \dots, n$ , or  $-\mathbf{e}_j^T \mathbf{x} = 0$  for  $j = 1, 2, \dots, n$ ,  
 associated with an LPP are called its **defining hyperplanes**.
- The associated **half spaces** are  
 $\mathbf{a}_i^T \mathbf{x} \leq b_i$ ,  $i = 1, 2, \dots, m$ ,  
 $x_j \geq 0$ ,  $j = 1, 2, \dots, n$ , or  $-\mathbf{e}_j^T \mathbf{x} \leq 0$ ,  $j = 1, 2, \dots, n$ .

For example the **defining hyperplanes** in **Example 2** are  
 $H_1 = \{[x, y]^T : x = 0\}$ ,  $H_2 = \{[x, y]^T : y = 0\}$ ,  
 $H_3 = \{[x, y]^T : x + 2y = 1\}$ ,  $H_4 = \{[x, y]^T : 2x + 4y = 4\}$   
 and  $H_5 = \{[x, y]^T : -x + y = 1\}$ .



# Solution by graphical method of LPP's in two variables

- **Example 1:** Given the linear programming problem

Max  $5x + 2y$

subject to

$$3x + 2y \leq 6$$

# Solution by graphical method of LPP's in two variables

- **Example 1:** Given the linear programming problem

$$\text{Max } 5x + 2y$$

subject to

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

# Solution by graphical method of LPP's in two variables

- **Example 1:** Given the linear programming problem

$$\text{Max } 5x + 2y$$

subject to

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$

# Solution by graphical method of LPP's in two variables

- **Example 1:** Given the linear programming problem

$$\text{Max } 5x + 2y$$

subject to

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$

The **optimal solution** to be  $x = 2, y = 0$  that is  $[2, 0]^T$  which happened to be a **corner point** of the feasible region.

# Solution by graphical method of LPP's in two variables

- **Example 1:** Given the linear programming problem

$$\text{Max } 5x + 2y$$

subject to

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$

The **optimal solution** to be  $x = 2, y = 0$  that is  $[2, 0]^T$  which happened to be a **corner point** of the feasible region.

Later we will again convince ourselves ( by using more rigor) that this is indeed the **optimal solution**. The **optimal value** is **10**.

# Solution by graphical method of LPP's in two variables

- **Example 1:** Given the linear programming problem

$$\text{Max } 5x + 2y$$

subject to

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$

The **optimal solution** to be  $x = 2, y = 0$  that is  $[2, 0]^T$  which happened to be a **corner point** of the feasible region.

Later we will again convince ourselves ( by using more rigor) that this is indeed the **optimal solution**. The **optimal value** is **10**.

- **Example 2:** Consider the problem,  
Min  $-x + 2y$   
subject to

- **Example 2:** Consider the problem,  
Min  $-x + 2y$   
subject to  
 $x + 2y \geq 1$



- **Example 2:** Consider the problem,

Min  $-x + 2y$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

- **Example 2:** Consider the problem,

Min  $-x + 2y$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

- **Example 2:** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

The above linear programming problem **does not** have an optimal solution.

- **Example 2:** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

The above linear programming problem **does not** have an optimal solution.

- **Example 3:** Note that in the above problem keeping the feasible region same, if we just change the objective function to  $\text{Min } 2x + y$ , then the changed problem has a **unique** optimal solution.

- **Example 2:** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

The above linear programming problem **does not** have an optimal solution.

- **Example 3:** Note that in the above problem keeping the feasible region same, if we just change the objective function to  $\text{Min } 2x + y$ , then the changed problem has a **unique** optimal solution.

- **Example 4:** Also in **Example 2** if we change objective function to Min  $x + 2y$  then the changed problem has infinitely many optimal solutions, although the set of optimal solutions is bounded.

- **Example 4:** Also in **Example 2** if we change objective function to  $\text{Min } x + 2y$  then the changed problem has **infinitely many** optimal solutions, although the set of optimal solutions is **bounded**.
- **Example 5:** Note that in **Example 2** keeping the feasible region same, if we just change the objective function to  $\text{Min } y$ , then the changed problem has **infinitely many** optimal solutions and the set of optimal solutions is **unbounded**.

- **Example 4:** Also in **Example 2** if we change objective function to  $\text{Min } x + 2y$  then the changed problem has **infinitely many** optimal solutions, although the set of optimal solutions is **bounded**.
- **Example 5:** Note that in **Example 2** keeping the feasible region same, if we just change the objective function to  $\text{Min } y$ , then the changed problem has **infinitely many** optimal solutions and the set of optimal solutions is **unbounded**.
- **Example 6:** Max  $-x + 2y$   
subject to  
 $x + 2y \leq 1$



- **Example 4:** Also in **Example 2** if we change objective function to  $\text{Min } x + 2y$  then the changed problem has **infinitely many** optimal solutions, although the set of optimal solutions is **bounded**.
- **Example 5:** Note that in **Example 2** keeping the feasible region same, if we just change the objective function to  $\text{Min } y$ , then the changed problem has **infinitely many** optimal solutions and the set of optimal solutions is **unbounded**.
- **Example 6:** Max  $-x + 2y$   
subject to  
 $x + 2y \leq 1$   
 $-x + y \geq 1$ ,

- **Example 4:** Also in **Example 2** if we change objective function to  $\text{Min } x + 2y$  then the changed problem has **infinitely many** optimal solutions, although the set of optimal solutions is **bounded**.
- **Example 5:** Note that in **Example 2** keeping the feasible region same, if we just change the objective function to  $\text{Min } y$ , then the changed problem has **infinitely many** optimal solutions and the set of optimal solutions is **unbounded**.
- **Example 6:** Max  $-x + 2y$   
subject to  
 $x + 2y \leq 1$   
 $-x + y \geq 1$ ,  
 $x \geq 0, y \geq 0$ .

- **Example 4:** Also in **Example 2** if we change objective function to  $\text{Min } x + 2y$  then the changed problem has **infinitely many** optimal solutions, although the set of optimal solutions is **bounded**.
- **Example 5:** Note that in **Example 2** keeping the feasible region same, if we just change the objective function to  $\text{Min } y$ , then the changed problem has **infinitely many** optimal solutions and the set of optimal solutions is **unbounded**.
- **Example 6:** Max  $-x + 2y$   
subject to  
 $x + 2y \leq 1$   
 $-x + y \geq 1$ ,  
 $x \geq 0, y \geq 0$ .  
 Clearly the feasible region of this problem is the **empty set**. So this problem is called **infeasible**, and since it **does not** have a **feasible solution** it obviously **does not** have an **optimal solution**.

- Question 1: Can there be exactly 2, 5, or say exactly 100 optimal solutions of a LPP?

- Question 1: Can there be exactly 2, 5, or say exactly 100 optimal solutions of a LPP?
- If we have two optimal solutions then what about points in between and on the line segment joining these two solutions?

- Question 1: Can there be exactly 2, 5, or say exactly 100 optimal solutions of a LPP?
- If we have two optimal solutions then what about points in between and on the line segment joining these two solutions?
- Question 2: Is the set of optimal solutions of an LPP a convex set?

- Question 1: Can there be exactly 2, 5, or say exactly 100 optimal solutions of a LPP?
- If we have two optimal solutions then what about points in between and on the line segment joining these two solutions?
- Question 2: Is the set of optimal solutions of an LPP a convex set?

That is, if  $\mathbf{x}_1$  and  $\mathbf{x}_2$  are two optimal solutions of a LPP,

- **Question 1:** Can there be exactly 2, 5, or say exactly 100 optimal solutions of a LPP?
- If we have two optimal solutions then what about points in between and on the line segment joining these two solutions?
- **Question 2:** Is the set of optimal solutions of an LPP a convex set?

That is, if  $\mathbf{x}_1$  and  $\mathbf{x}_2$  are two optimal solutions of a LPP, then are  $\mathbf{y}$ 's of the form  $\mathbf{y} = \lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2$ ,  $0 \leq \lambda \leq 1$ , also optimal solutions of the LPP?



- A nonempty set,  $S \subseteq \mathbb{R}^n$  is said to be a **convex set** if for all  $\mathbf{x}_1, \mathbf{x}_2 \in S$ ,  $\lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2 \in S$ , for all  $0 \leq \lambda \leq 1$ .

- A nonempty set,  $S \subseteq R^n$  is said to be a **convex set** if for all  $\mathbf{x}_1, \mathbf{x}_2 \in S$ ,  $\lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2 \in S$ , for all  $0 \leq \lambda \leq 1$ .
- $\lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2$ ,  $0 \leq \lambda \leq 1$  is called a **convex combination** of  $\mathbf{x}_1$  and  $\mathbf{x}_2$ .

- A nonempty set,  $S \subseteq R^n$  is said to be a **convex set** if for all  $\mathbf{x}_1, \mathbf{x}_2 \in S$ ,  $\lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2 \in S$ , for all  $0 \leq \lambda \leq 1$ .
- $\lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2$ ,  $0 \leq \lambda \leq 1$  is called a **convex combination** of  $\mathbf{x}_1$  and  $\mathbf{x}_2$ .
- If in the above expression,  $0 < \lambda < 1$ , then the **convex combination** is said to be a **strict convex combination** of  $\mathbf{x}_1$  and  $\mathbf{x}_2$ .

- A nonempty set,  $S \subseteq R^n$  is said to be a **convex set** if for all  $\mathbf{x}_1, \mathbf{x}_2 \in S$ ,  $\lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2 \in S$ , for all  $0 \leq \lambda \leq 1$ .
- $\lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2$ ,  $0 \leq \lambda \leq 1$  is called a **convex combination** of  $\mathbf{x}_1$  and  $\mathbf{x}_2$ .
- If in the above expression,  $0 < \lambda < 1$ , then the **convex combination** is said to be a **strict convex combination** of  $\mathbf{x}_1$  and  $\mathbf{x}_2$ .

**Example:**  $H = \{[x, y]^T : x + 2y \leq 1\}$ , is a convex set.  
 Note that  $\mathbf{x}_1 = [0, 0]^T$ ,  $\mathbf{x}_2 = [1, 0]^T \in H$ .

- A nonempty set,  $S \subseteq R^n$  is said to be a **convex set** if for all  $\mathbf{x}_1, \mathbf{x}_2 \in S$ ,  $\lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2 \in S$ , for all  $0 \leq \lambda \leq 1$ .
- $\lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2$ ,  $0 \leq \lambda \leq 1$  is called a **convex combination** of  $\mathbf{x}_1$  and  $\mathbf{x}_2$ .
- If in the above expression,  $0 < \lambda < 1$ , then the **convex combination** is said to be a **strict convex combination** of  $\mathbf{x}_1$  and  $\mathbf{x}_2$ .

**Example:**  $H = \{[x, y]^T : x + 2y \leq 1\}$ , is a convex set.

Note that  $\mathbf{x}_1 = [0, 0]^T$ ,  $\mathbf{x}_2 = [1, 0]^T \in H$ .

$[0, 0]^T = 1[0, 0]^T + 0[1, 0]^T$  is a **convex combination** of  $\mathbf{x}_1$  and  $\mathbf{x}_2$ , but **not** a **strict convex combination** of  $\mathbf{x}_1$  and  $\mathbf{x}_2$  whereas

- A nonempty set,  $S \subseteq R^n$  is said to be a **convex set** if for all  $\mathbf{x}_1, \mathbf{x}_2 \in S$ ,  $\lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2 \in S$ , for all  $0 \leq \lambda \leq 1$ .
- $\lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2$ ,  $0 \leq \lambda \leq 1$  is called a **convex combination** of  $\mathbf{x}_1$  and  $\mathbf{x}_2$ .
- If in the above expression,  $0 < \lambda < 1$ , then the **convex combination** is said to be a **strict convex combination** of  $\mathbf{x}_1$  and  $\mathbf{x}_2$ .

**Example:**  $H = \{[x, y]^T : x + 2y \leq 1\}$ , is a convex set.

Note that  $\mathbf{x}_1 = [0, 0]^T$ ,  $\mathbf{x}_2 = [1, 0]^T \in H$ .

$[0, 0]^T = 1[0, 0]^T + 0[1, 0]^T$  is a **convex combination** of  $\mathbf{x}_1$  and  $\mathbf{x}_2$ , but **not** a **strict convex combination** of  $\mathbf{x}_1$  and  $\mathbf{x}_2$  whereas

$[\frac{2}{3}, 0]^T = \frac{1}{3}[0, 0]^T + \frac{2}{3}[1, 0]^T$  is a **convex combination** as well as a **strict convex combination** of  $\mathbf{x}_1$  and  $\mathbf{x}_2$ .

- Let us first try to answer **Question 2**.

- Let us first try to answer **Question 2**.

If the answer to this question is a **YES** then that would imply that if a LPP has **more than one optimal solution** then it should have **infinitely many optimal solutions**, so the answer to **Question 1** would be a **NO**.



- Let us first try to answer **Question 2**.

If the answer to this question is a **YES** then that would imply that if a LPP has **more than one optimal solution** then it should have **infinitely many optimal solutions**, so the answer to **Question 1** would be a **NO**.

Answer to **Question 2** is **YES**, that is the set of all **optimal solutions** of an LPP is indeed a **convex set**.

- Let us first try to answer **Question 2**.

If the answer to this question is a **YES** then that would imply that if a LPP has **more than one optimal solution** then it should have **infinitely many optimal solutions**, so the answer to **Question 1** would be a **NO**.

Answer to **Question 2** is **YES**, that is the set of all **optimal solutions** of an LPP is indeed a **convex set**.

- **Question 3:** If the **feasible region** of a LPP is a **nonempty, bounded set** then does the LPP **always** have an **optimal solution**?

- Let us first try to answer **Question 2**.

If the answer to this question is a **YES** then that would imply that if a LPP has **more than one optimal solution** then it should have **infinitely many optimal solutions**, so the answer to **Question 1** would be a **NO**.

Answer to **Question 2** is **YES**, that is the set of all **optimal solutions** of an LPP is indeed a **convex set**.

- **Question 3:** If the **feasible region** of a LPP is a **nonempty, bounded set** then does the LPP **always** have an **optimal solution**?

The answer to this question is **YES**, due to **Weierstrass**, called the **Extreme Value Theorem**:

- **Extreme Value Theorem:** If  $S$  is a nonempty, closed, bounded subset of  $\mathbb{R}^n$  and  $f : S \rightarrow \mathbb{R}$  is continuous, then  $f$  attains both its minimum and maximum value in  $S$ .

- **Extreme Value Theorem:** If  $S$  is a nonempty, closed, bounded subset of  $\mathbb{R}^n$  and  $f : S \rightarrow \mathbb{R}$  is continuous, then  $f$  attains both its minimum and maximum value in  $S$ .
- **Question 4:** Whenever a LPP has an optimal solution does there always exist **at least one corner point** (points lying at the point of intersection of at least two distinct lines in case  $n = 2$ ), at which the optimal value is attained?

- **Extreme Value Theorem:** If  $S$  is a nonempty, closed, bounded subset of  $\mathbb{R}^n$  and  $f : S \rightarrow \mathbb{R}$  is continuous, then  $f$  attains both its minimum and maximum value in  $S$ .
- **Question 4:** Whenever a LPP has an optimal solution does there always exist **at least one corner point** (points lying at the point of intersection of at least two distinct lines in case  $n = 2$ ), at which the optimal value is attained?
- Given a LPP with a nonempty feasible region,  $Fea(LPP) = S \subset \mathbb{R}^n$ , an  $\mathbf{x} \in S$  is called a corner point of  $S$ , if  $\mathbf{x}$  lies at the point of intersection of  $n$  linearly independent hyperplanes defining  $S$ .

- Consider the feasible region of **Example 2**,

- Consider the feasible region of **Example 2**,  
 $S = \{[x, y]^T : x \geq 0, y \geq 0, 2x + 4y \leq 4, -x + y \leq 1, x + 2y \geq 1\}$ .



- Consider the feasible region of **Example 2**,  
 $S = \{[x, y]^T : x \geq 0, y \geq 0, 2x + 4y \leq 4, -x + y \leq 1, x + 2y \geq 1\}$ .
- The defining hyperplanes are  $H_1 = \{[x, y]^T : x = 0\}$ ,

- Consider the feasible region of **Example 2**,  
 $S = \{[x, y]^T : x \geq 0, y \geq 0, 2x + 4y \leq 4, -x + y \leq 1, x + 2y \geq 1\}$ .
- The defining hyperplanes are  $H_1 = \{[x, y]^T : x = 0\}$ ,  
 $H_2 = \{[x, y]^T : y = 0\}$ ,

- Consider the feasible region of **Example 2**,  
 $S = \{[x, y]^T : x \geq 0, y \geq 0, 2x + 4y \leq 4, -x + y \leq 1, x + 2y \geq 1\}$ .
- The defining hyperplanes are  $H_1 = \{[x, y]^T : x = 0\}$ ,  
 $H_2 = \{[x, y]^T : y = 0\}$ ,  $H_3 = \{[x, y]^T : x + 2y = 1\}$ ,

- Consider the feasible region of **Example 2**,  
 $S = \{[x, y]^T : x \geq 0, y \geq 0, 2x + 4y \leq 4, -x + y \leq 1, x + 2y \geq 1\}$ .
- The defining hyperplanes are  $H_1 = \{[x, y]^T : x = 0\}$ ,  
 $H_2 = \{[x, y]^T : y = 0\}$ ,  $H_3 = \{[x, y]^T : x + 2y = 1\}$ ,  
 $H_4 = \{[x, y]^T : 2x + 4y = 4\}$  and

- Consider the feasible region of **Example 2**,  
 $S = \{[x, y]^T : x \geq 0, y \geq 0, 2x + 4y \leq 4, -x + y \leq 1, x + 2y \geq 1\}$ .
- The defining hyperplanes are  $H_1 = \{[x, y]^T : x = 0\}$ ,  
 $H_2 = \{[x, y]^T : y = 0\}$ ,  $H_3 = \{[x, y]^T : x + 2y = 1\}$ ,  
 $H_4 = \{[x, y]^T : 2x + 4y = 4\}$  and  
 $H_5 = \{[x, y]^T : -x + y = 1\}$ .
- $[1, 0]^T, [0, 1]^T, [0, \frac{1}{2}]^T, [2, 0]^T$  are the **corner points** of  $S$ .
- $[1, 0]^T$  lies on  $H_2, H_3$ , which are **LI**.

- Consider the feasible region of **Example 2**,  
 $S = \{[x, y]^T : x \geq 0, y \geq 0, 2x + 4y \leq 4, -x + y \leq 1, x + 2y \geq 1\}$ .
- The defining hyperplanes are  $H_1 = \{[x, y]^T : x = 0\}$ ,  
 $H_2 = \{[x, y]^T : y = 0\}$ ,  $H_3 = \{[x, y]^T : x + 2y = 1\}$ ,  
 $H_4 = \{[x, y]^T : 2x + 4y = 4\}$  and  
 $H_5 = \{[x, y]^T : -x + y = 1\}$ .
- $[1, 0]^T, [0, 1]^T, [0, \frac{1}{2}]^T, [2, 0]^T$  are the **corner points** of  $S$ .
- $[1, 0]^T$  lies on  $H_2, H_3$ , which are **LI**.
- $[0, 1]^T$  lies on  $H_1, H_4, H_5$ , of which each of  $\{H_1, H_4\}$ ,  
 $\{H_1, H_5\}$ ,  $\{H_4, H_5\}$  is **LI**.

- Consider the feasible region of **Example 2**,  
 $S = \{[x, y]^T : x \geq 0, y \geq 0, 2x + 4y \leq 4, -x + y \leq 1, x + 2y \geq 1\}$ .
- The defining hyperplanes are  $H_1 = \{[x, y]^T : x = 0\}$ ,  
 $H_2 = \{[x, y]^T : y = 0\}$ ,  $H_3 = \{[x, y]^T : x + 2y = 1\}$ ,  
 $H_4 = \{[x, y]^T : 2x + 4y = 4\}$  and  
 $H_5 = \{[x, y]^T : -x + y = 1\}$ .
- $[1, 0]^T, [0, 1]^T, [0, \frac{1}{2}]^T, [2, 0]^T$  are the **corner points** of  $S$ .
- $[1, 0]^T$  lies on  $H_2, H_3$ , which are **LI**.
- $[0, 1]^T$  lies on  $H_1, H_4, H_5$ , of which each of  $\{H_1, H_4\}$ ,  
 $\{H_1, H_5\}$ ,  $\{H_4, H_5\}$  is **LI**.
- $[0, \frac{1}{2}]^T$  lies on  $H_1, H_3$ , which are **LI**.

- Consider the feasible region of **Example 2**,  
 $S = \{[x, y]^T : x \geq 0, y \geq 0, 2x + 4y \leq 4, -x + y \leq 1, x + 2y \geq 1\}$ .
- The defining hyperplanes are  $H_1 = \{[x, y]^T : x = 0\}$ ,  
 $H_2 = \{[x, y]^T : y = 0\}$ ,  $H_3 = \{[x, y]^T : x + 2y = 1\}$ ,  
 $H_4 = \{[x, y]^T : 2x + 4y = 4\}$  and  
 $H_5 = \{[x, y]^T : -x + y = 1\}$ .
- $[1, 0]^T, [0, 1]^T, [0, \frac{1}{2}]^T, [2, 0]^T$  are the **corner points** of  $S$ .
- $[1, 0]^T$  lies on  $H_2, H_3$ , which are **LI**.
- $[0, 1]^T$  lies on  $H_1, H_4, H_5$ , of which each of  $\{H_1, H_4\}$ ,  
 $\{H_1, H_5\}$ ,  $\{H_4, H_5\}$  is **LI**.
- $[0, \frac{1}{2}]^T$  lies on  $H_1, H_3$ , which are **LI**.
- $[2, 0]^T$  lies on  $H_2, H_4$ , which are **LI**.



- An  $\mathbf{x} \in \text{Fea}(LPP)$  which lies in **atleast one defining hyperplane** of the  $\text{Fea}(LPP)$  is a **boundary point** of  $\text{Fea}(LPP)$ .

- An  $\mathbf{x} \in \text{Fea}(LPP)$  which lies in **atleast one defining hyperplane** of the  $\text{Fea}(LPP)$  is a **boundary point** of  $\text{Fea}(LPP)$ .

$[\frac{3}{2}, 0]^T$  is a **boundary point** but **not** a corner point of  $S$  since it lies on only **one** defining hyperplane of  $S$ .

- An  $\mathbf{x} \in \text{Fea}(LPP)$  which lies in **atleast one defining hyperplane** of the  $\text{Fea}(LPP)$  is a **boundary point** of  $\text{Fea}(LPP)$ .  
 $[\frac{3}{2}, 0]^T$  is a **boundary point** but **not** a corner point of  $S$  since it lies on only **one** defining hyperplane of  $S$ .
- An  $\mathbf{x} \in \text{Fea}(LPP)$  which does **not** lie on any of the **defining hyperplanes** of  $\text{Fea}(LPP)$  is an **interior point** of  $\text{Fea}(LPP)$

- An  $\mathbf{x} \in \text{Fea}(LPP)$  which lies in **atleast one defining hyperplane** of the  $\text{Fea}(LPP)$  is a **boundary point** of  $\text{Fea}(LPP)$ .

$[\frac{3}{2}, 0]^T$  is a **boundary point** but **not** a corner point of  $S$  since it lies on only **one** defining hyperplane of  $S$ .

- An  $\mathbf{x} \in \text{Fea}(LPP)$  which does **not** lie on any of the **defining hyperplanes** of  $\text{Fea}(LPP)$  is an **interior point** of  $\text{Fea}(LPP)$

$[\frac{3}{2}, \frac{1}{8}]^T$  is an **interior point** of  $S$  since it **does not** lie on any defining hyperplane of  $S$ .

- Note that our feasible region  $S$ , has  $(m + n)$  **defining hyperplanes**.

- Note that our feasible region  $S$ , has  $(m + n)$  **defining hyperplanes**.
- (Alternate definition of corner points): The **corner points** of the feasible region of a LPP **cannot** be written as a **strict convex combination** of **two distinct points** of the feasible region, in other words those are all **extreme points** of the feasible region.

- Note that our feasible region  $S$ , has  $(m + n)$  **defining hyperplanes**.
- (Alternate definition of corner points): The **corner points** of the feasible region of a LPP **cannot** be written as a **strict convex combination** of **two distinct points** of the feasible region, in other words those are all **extreme points** of the feasible region.
- Given a nonempty **convex set**,  $S \subset \mathbb{R}^n$ , an  $\mathbf{x} \in S$  is said to be an **extreme point** of  $S$  if  $\mathbf{x}$  **cannot** be written as a **strict convex combination** of **two distinct elements** of  $S$ .

- Note that our feasible region  $S$ , has  $(m + n)$  **defining hyperplanes**.
- (Alternate definition of corner points): The **corner points** of the feasible region of a LPP **cannot** be written as a **strict convex combination** of **two distinct points** of the feasible region, in other words those are all **extreme points** of the feasible region.
- Given a nonempty **convex set**,  $S \subset \mathbb{R}^n$ , an  $\mathbf{x} \in S$  is said to be an **extreme point** of  $S$  if  $\mathbf{x}$  **cannot** be written as a **strict convex combination** of **two distinct elements** of  $S$ .  
That is if  $\mathbf{x} = \lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2$ , for some  $0 < \lambda < 1$  and  $\mathbf{x}_1, \mathbf{x}_2 \in S$ , then



- Note that our feasible region  $S$ , has  $(m + n)$  **defining hyperplanes**.
- (Alternate definition of corner points): The **corner points** of the feasible region of a LPP **cannot** be written as a **strict convex combination** of **two distinct points** of the feasible region, in other words those are all **extreme points** of the feasible region.
- Given a nonempty **convex set**,  $S \subset \mathbb{R}^n$ , an  $\mathbf{x} \in S$  is said to be an **extreme point** of  $S$  if  $\mathbf{x}$  **cannot** be written as a **strict convex combination** of **two distinct elements** of  $S$ .  
That is if  $\mathbf{x} = \lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2$ , for some  $0 < \lambda < 1$  and  $\mathbf{x}_1, \mathbf{x}_2 \in S$ , then  $\mathbf{x}_1 = \mathbf{x}_2 = \mathbf{x}$ .

- Note that our feasible region  $S$ , has  $(m + n)$  **defining hyperplanes**.
- (Alternate definition of corner points): The **corner points** of the feasible region of a LPP **cannot** be written as a **strict convex combination** of **two distinct points** of the feasible region, in other words those are all **extreme points** of the feasible region.
- Given a nonempty **convex set**,  $S \subset \mathbb{R}^n$ , an  $\mathbf{x} \in S$  is said to be an **extreme point** of  $S$  if  $\mathbf{x}$  **cannot** be written as a **strict convex combination** of **two distinct elements** of  $S$ .  
That is if  $\mathbf{x} = \lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2$ , for some  $0 < \lambda < 1$  and  $\mathbf{x}_1, \mathbf{x}_2 \in S$ , then  $\mathbf{x}_1 = \mathbf{x}_2 = \mathbf{x}$ .
- All points on the **boundary** of a **disc** are extreme points of the disc.
- A **hyperplane**, **half space** does not have any extreme point.

- **Theorem:** If  $S = \text{Fea}(LPP)$  is nonempty, where  
 $S = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}_i^T \mathbf{x} \leq b_i, i = 1, \dots, m, -\mathbf{e}_j^T \mathbf{x} \leq 0, j = 1, \dots, n\}$   
then  $\mathbf{x} \in S$  is a **corner point** of  $S$  if and only if it is an **extreme point** of  $S$ .

- Theorem:** If  $S = \text{Fea}(LPP)$  is nonempty, where  

$$S = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}_i^T \mathbf{x} \leq b_i, i = 1, \dots, m, \quad -\mathbf{e}_j^T \mathbf{x} \leq 0, j = 1, \dots, n\}$$
 then  $\mathbf{x} \in S$  is a **corner point** of  $S$  if and only if it is an **extreme point** of  $S$ .
- The total number of **extreme points** of the feasible region of a LPP with  $(m + n)$  constraints (including the non negativity constraints) is  $\leq (m + n)C_n$ .

- Theorem:** If  $S = \text{Fea}(LPP)$  is nonempty, where  

$$S = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}_i^T \mathbf{x} \leq b_i, i = 1, \dots, m, \quad -\mathbf{e}_j^T \mathbf{x} \leq 0, j = 1, \dots, n\}$$
 then  $\mathbf{x} \in S$  is a **corner point** of  $S$  if and only if it is an **extreme point** of  $S$ .
- The total number of **extreme points** of the feasible region of a LPP with  $(m + n)$  constraints (including the non negativity constraints) is  $\leq (m + n)C_n$ .
- Exercise:** Think of a LPP  $(m + n)$  constraints such that the number of **extreme points** of the  $\text{Fea}(LPP)$  is equal to  $\leq (m + n)C_n$ .

- **Exercise:** If possible give an example of a LPP with  $(m + n)$  constraints such that the number of **extreme points** of the  $Fea(LPP)$  is strictly greater than  $(m + n)$  and is strictly less than  $(m + n)_{C_n}$ .

- **Exercise:** If possible give an example of a LPP with  $(m + n)$  constraints such that the number of **extreme points** of the  $Fea(LPP)$  is strictly greater than  $(m + n)$  and is strictly less than  $(m + n)_{C_n}$ .
- **Exercise:** If possible give an example of a LPP with  $(m + n)$  constraints such that the number of **extreme points** of the  $Fea(LPP)$  is strictly less than  $(m + n)$ .

- **Exercise:** If possible give an example of a LPP with  $(m + n)$  constraints such that the number of **extreme points** of the  $Fea(LPP)$  is strictly greater than  $(m + n)$  and is strictly less than  $(m + n)_{C_n}$ .
- **Exercise:** If possible give an example of a LPP with  $(m + n)$  constraints such that the number of **extreme points** of the  $Fea(LPP)$  is strictly less than  $(m + n)$ .
- **Question:** Does the feasible region of a LPP of the form given before (with non negativity constraints) always have an **extreme point**?



- **Exercise:** If possible give an example of a LPP with  $(m + n)$  constraints such that the number of **extreme points** of the  $Fea(LPP)$  is strictly greater than  $(m + n)$  and is strictly less than  $(m + n)_{C_n}$ .
- **Exercise:** If possible give an example of a LPP with  $(m + n)$  constraints such that the number of **extreme points** of the  $Fea(LPP)$  is strictly less than  $(m + n)$ .
- **Question:** Does the feasible region of a LPP of the form given before (with non negativity constraints) always have an **extreme point**?

Answer: **YES**, we will see this later.

- **Exercise:** If possible give an example of a LPP with  $(m + n)$  constraints such that the number of **extreme points** of the  $Fea(LPP)$  is strictly greater than  $(m + n)$  and is strictly less than  $(m + n)_{C_n}$ .
- **Exercise:** If possible give an example of a LPP with  $(m + n)$  constraints such that the number of **extreme points** of the  $Fea(LPP)$  is strictly less than  $(m + n)$ .
- **Question:** Does the feasible region of a LPP of the form given before (with non negativity constraints) always have an **extreme point**?

Answer: **YES**, we will see this later.

- **Exercise:** Think of a **convex set** defined by only one hyperplane. Will it have any **extreme point**?

- **Exercise:** If possible give an example of a LPP with  $(m + n)$  constraints such that the number of **extreme points** of the  $Fea(LPP)$  is strictly greater than  $(m + n)$  and is strictly less than  $(m + n)_{C_n}$ .
- **Exercise:** If possible give an example of a LPP with  $(m + n)$  constraints such that the number of **extreme points** of the  $Fea(LPP)$  is strictly less than  $(m + n)$ .
- **Question:** Does the feasible region of a LPP of the form given before (with non negativity constraints) always have an **extreme point**?

Answer: **YES**, we will see this later.

- **Exercise:** Think of a **convex set** defined by only one hyperplane. Will it have any **extreme point**?